

# THE STOCHASTIC RENORMALIZED MEAN CURVATURE FLOW FOR PLANAR CONVEX SETS

MARC ARNAUDON, KOLÉHÈ COULIBALY-PASQUIER, AND LAURENT MICLO

ABSTRACT. We investigate renormalized mean curvature flow (RMCF) and stochastic renormalized mean curvature flow (SRMCF) for convex sets in the plane. RMCF is the inverse gradient flow for logarithm of  $\sigma/\lambda^2$  where  $\sigma$  is the perimeter and  $\lambda$  is the volume. SRMCF is RMCF perturbed by some Brownian noise and has the remarkable property that it can be intertwined with Brownian motion, yielding a generalization of Pitman "2M - X" theorem. We prove that along RMCF, entropy  $\mathcal{E}_t$  for curvature as well as  $h_t := \sigma_t/\lambda_t$  are non-increasing. We deduce infinite lifetime and convergence to a disk after normalization. For SRMCF the situation is more complicated. As  $h_t$  is always a supermartingale, for  $\mathcal{E}_t$  to be a supermartingale, we need that the starting set is invariant by the isometry group  $G_n$  generated by the reflection with respect to the vertical line and the rotation of angle  $2\pi/n$ , for some  $n \geq 3$ . But for proving infinite lifetime, we need invariance of the starting set by  $G_n$  for some  $n \geq 7$ . We provide the first SRMCF with infinite lifetime which cannot be reduced to a finite dimensional flow. Gage inequality plays a major role in our study of the regularity of flows, as well as a careful investigation of morphological skeletons. We characterize symmetric convex sets with star shaped skeletons in terms of properties of their Gauss map. Finally, we establish a new isoperimetric estimate for these sets, of order  $1/n^4$  where  $n$  is the number of branches of the skeleton.

## 1. INTRODUCTION

The evolution of simple closed curves in  $\mathbb{R}^2$  by mean curvature flow (MCF) has been investigated for a long time. It can also be described as the inverse gradient flow for the perimeter, or as some kind of (nonlinear) geometrical heat equation. The motivation for the study of mean curvature flow comes from physics. In 1986, Gage and Hamilton ([6]) proved that starting from any convex smooth simple closed curve, the mean curvature flow converges in finite time to one point, and the form of the curve becomes circular. In 1987, Grayson ([7]) generalized this result to non necessarily convex starting curve. It is a remarkable fact that no intersection occurs during the evolution of the flow.

The renormalized mean curvature flow (RMCF) can roughly be defined as the solution to the evolution equation for curves by mean curvature, to which we add a constant normal field which prevents from implosion. More precisely we will prove in Lemma 2.4 that RMCF is the inverse gradient flow for logarithm of  $\sigma(\partial D)/\lambda(D)^2$  where the considered curve is the boundary  $\partial D$  of a bounded

---

*Date:* March 6, 2023 *File:* ACM2-230305.tex

LM acknowledges fundings from the grants ANR-17-EURE-0010 and AFOSR-22IOE016.

domain  $D$ ,  $\lambda(D)$  is the volume of the domain and  $\sigma(\partial D)$  is the perimeter of the curve. For this flow, intersection can occur when the starting curve is not convex. But when the starting curve is convex, we will prove in Theorem 2.3 that the lifetime of the flow  $(\partial D_t)_{t \geq 0}$  is infinite and the curve converges to a circle. Two quantities will be investigated for the convergence: the ratio  $h_t := \sigma(\partial D_t)/\lambda(D_t)$  and the entropy  $\text{Ent}_t := \int_{\partial D_t} \rho_t \log \rho_t$ ,  $\rho_t$  being the curvature at each point of  $\partial D_t$ . We will prove that these two quantities are non-increasing along the flow (Lemmas 2.7 and 2.16).

One of the main goals of this paper is the investigation of stochastic renormalized mean curvature flow (SRMCF) in  $\mathbb{R}^2$ . To define it, we add to the RMCF a one dimensional normal Brownian noise. The SRMCF can be considered as a noise model for the previous flow. The intensity of the noise is chosen so that the generator of the flow is intertwined with the one of the Brownian motion, via a Markov kernel (see e.g. [2] and [1]). Alternatively, very nice connections with Bessel 3 processes can be exhibited ([2]). In fact, when the intertwining can be realized with a pair of stochastic processes  $(X_t, D_t)_{t \geq 0}$  with a Brownian motion  $(X_t)_{t \geq 0}$  satisfying that at any time  $t \geq 0$ ,  $X_t$  is uniformly distributed inside  $D_t$  conditionally to  $(D_s)_{0 \leq s \leq t}$ , the construction is a generalization of Pitman "2M - X" theorem. An important object in the construction of the coupling  $(X_t, D_t)$  is the inner skeleton  $S_t$  of  $D_t$ , which is the singularity set of distance to boundary, inside  $D_t$ : the equation for  $\partial D_t$  has a component of the drift which is proportional to the local time of  $X_t$  at  $S_t$  ([1]). A very nice fact about the skeleton is that even if  $\partial D_t$  has a Brownian noise,  $S_t$  has finite variation. As we will see in the present paper, the inner skeleton  $S_t$  also plays a role in the lifetime of  $D_t$ . We will prove that starting with a convex subset  $D_0$  of  $\mathbb{R}^2$ , the explosion occurs only when  $\partial D_t$  meets  $S_t$  (Theorem 3.8). We will also prove that similarly to the deterministic situation, the process  $h_t$  is a supermartingale (Lemma 4.3). For the entropy being a supermartingale we will need that  $D_0$  is invariant by the linear group  $G_n$  generated by the rotation of angle  $2\pi/n$  with  $n \geq 3$ , and the symmetry with respect to an axis (we will choose the vertical one, see Proposition 4.8).  $G_n$ -invariance for any  $n \geq 2$  will be proved to be preserved by the flow. Finally we will prove that  $G_n$ -invariance of  $D_0$  with  $n \geq 7$  implies infinite lifetime for the stochastic renormalized mean curvature flow (Theorem 4.15).

In Section 5 we investigate some class of convex sets in  $\mathbb{R}^2$  which are symmetric with respect to  $G_n$ , and with star-shaped skeletons. We prove (Proposition 5.5) that they are preserved by all our flows. The last section is devoted to the proof of a new isoperimetric inequality for these classes of convex sets (Proposition 6.1). A bound of order  $1/n^4$  is observed.

**Definition 1.1.** *A simple closed curve is said to be strictly convex when its geodesic curvatures are positive.*

Note that the inside domain of such a curve is strictly convex in the usual sense i.e. it is strictly contained in one side of any tangent line, except for the contact

point, but the converse is not necessarily true, as the curvature may vanish at isolated points.

It is possible to parametrize a simple strictly convex closed curve in  $\mathbb{R}^2$  using the angle  $\theta$  between the tangent vector  $T := (\cos(\theta), \sin(\theta))$  and the oriented  $x$  axis. The coordinate  $\theta$  will make the equations of our flows simple to analyze, in particular since operators  $\partial_\theta$  and  $\partial_t$  will commute, contrarily to derivative with respect to curvilinear abscissa  $\partial_s$  and  $\partial_t$ , as shown in (12). We will essentially use the one-to-one correspondence in  $\mathbb{R}^2$  between simple strictly convex closed curves (up to translation) and positive functions  $\rho$  that satisfy

$$\int_0^{2\pi} \frac{\cos(\theta)}{\rho(\theta)} d\theta = \int_0^{2\pi} \frac{\sin(\theta)}{\rho(\theta)} d\theta = 0 \quad (1)$$

as in Lemma 4.1.1 of Gage and Hamilton [6]. The function  $\rho$  is in fact the curvature of the curve, see also Section 5.

Using the stochastic evolution of curves, such as stochastic mean curvature flow (SMCF) (20) and SRMCF (11), we will derive the equation of the curvature, and we will show in Theorem 3.8 that positivity of the curvature is preserved by the flows. Also Equation (1) is conserved along these equations, and so we will get an alternative definition of the stochastic evolution of a convex curve in term of the solution of some stochastic partial differential equation (Theorem 3.8).

To fix some notations used throughout the paper, let us present some notions associated to a simple  $C^2$  closed curve  $C : \mathbb{T} \ni u \mapsto C(u) \in \mathbb{R}^2$ , where  $\mathbb{T} := \mathbb{R}/(2\pi\mathbb{Z})$ . In this paper, all curves will be closed and immersed.

The bounded domain whose boundary is  $C$  is denoted  $D$ . The quantities  $\lambda(D)$  and  $\sigma(C)$  respectively stand for the volume of  $D$  and the perimeter of  $C = \partial D$ . We designate by  $h(D)$  the isoperimetric ratio  $\sigma(\partial D)/\lambda(D)$ , not to be confounded with the planar isoperimetric ratio  $\sigma(\partial D)^2/\lambda(D)$ . For any  $x \in C$ ,  $\nu_C(x)$  is the outer unit normal vector of the curve  $C$  at the point  $x$  and  $\rho_C(x)$  is the corresponding mean curvature.

## 2. THE RENORMALIZED MEAN CURVATURE FLOW (RMCF)

Let  $u \in \mathbb{T} \mapsto C_0(u)$  be a simple  $C^{2+\alpha}$  closed curve in  $\mathbb{R}^2$ .

**Definition 2.1.** *Let  $C(t, u) : [0, T_c) \times \mathbb{T} \rightarrow \mathbb{R}^2$  a family of simple closed curve, we say that  $C(t, \cdot)$  evolves under the renormalized mean curvature flow, shortly (RMCF), if it satisfy the following equation*

$$\begin{cases} \partial_t C(t, u) &= (-\rho_t(C(t, u)) + 2h(D(t)))\nu_{C(t, \cdot)}(C(t, u)) \\ C(0, u) &= C_0(u) \end{cases} \quad (2)$$

Sometimes we will omit the parameter  $D(t)$  or  $C(t, \cdot)$  in the notation, and even simply write for example  $h(t)$  for  $h(D(t))$ .

**Remark 2.2.** *Since the symbol of the equation (2) is same as in the mean curvature flow, we have the short time existence of the equation (2) (i.e.  $T_c > 0$ ) for simple initial closed  $C^\infty$  curve, see for example [6] or [5]. We need the simplicity of the curve to have no ambiguity for  $h$  (mainly for the interior volume, see Figure.2*

example a) ) and to have well-defined outer unit normal vector.

An alternative proof of the existence, using quasi-linear equations, can be found in Chapter 4 of [2], Theorem 40 with  $B_t = 0$ .

**2.1. The main result of this section.** As we will see later, the evolution of the curvature is easier to analyze. Also in the Euclidean plane we have some geometrical inequalities concerning convex closed curve: the Gage inequality (5), involves the non local term  $h$ , and permits to have some a priori estimate along the solution; also the usual isoperimetric inequality plays a important role in the study of this flow. The principal result of this section is the following:

**Theorem 2.3.** *If the starting closed curve  $C(0, \cdot)$  is strictly convex and simple, then the solution  $C(t, \cdot)$  of equation (2) is defined for all  $t \in [0, \infty[$ , it remains strictly convex and simple for all times and becomes circular, the isoperimetric ratio is decreasing (except for circular starting curves). After renormalization and translation, we have the convergence with respect to the Hausdorff metric*

$$\frac{1}{\sqrt{6t}}[C(t, \cdot) - c_{\text{int}}(t)] \xrightarrow{d_H} C(0, 1),$$

to the circle of center 0 and radius 1, where for all  $t$ ,  $c_{\text{int}}(t)$  is a center of an inscribed circle of  $C(t, \cdot)$ .

The rest of this section is devoted to the proof of Theorem 2.3.

**2.2. Gradient flow formulation, and evolution of geometric quantities.** To a solution  $(C(t, \cdot))_{t \in [0, T_c[}$  of (2), associate

$$\forall t \in [0, T_c[, \forall u \in \mathbb{T}, \quad v(t, u) := |\partial_u C(t, u)|$$

and  $s$  the arc-length parametrization,  $\partial_s := \frac{1}{v} \partial_u$  (equivalently  $ds = v du$ ), started at  $C(t, s)|_{s=0} = C(t, u)|_{u=0}$ . To prevent the domain of definition of  $s$  to depend on  $t$ , we decide that  $s$  is defined in  $\mathbb{R}$  and periodic with period  $\sigma(C(t))$ . Let  $T := \partial_s C(t, s)$ , the tangent vector of the curve  $C(t, \cdot)$  at the point  $C(t, s)$ . Let  $\nu(t, s)$  be the unit vector obtained by a rotation of  $T(t, s)$  by an angle of  $-\pi/2$ . To make a choice of orientation, we will always assume that  $\nu$  is the outer normal of the curve.

**Lemma 2.4.** *Equation (2) is the inverse gradient flow of the functional*

$$\Psi : D \mapsto \ln \frac{\sigma(\partial D)}{\lambda(D)^2}.$$

*Proof.* Let  $C(t, u) : [0, T) \times \mathbb{T} \rightarrow \mathbb{R}^2$  be a family of simple closed curves, such that  $\partial_t C(t, u) = X(t, u)$  for some smooth  $X : [0, T) \times \mathbb{T} \rightarrow \mathbb{R}^2$ . Then

$$\frac{d}{dt} \lambda(t) = \int_{C(t, \cdot)} \langle X, \nu \rangle ds$$

and

$$\frac{d}{dt} \sigma(t) = \int_{C(t, \cdot)} \langle X, \nu \rangle \rho ds,$$

so we have

$$\begin{aligned} \frac{d}{dt} \frac{\sigma(t)}{\lambda(t)^2} &= \frac{1}{\lambda(t)^2} \left( \int_{C(t, \cdot)} \langle X, \nu \rangle \rho ds - \frac{2\sigma(t)}{\lambda(t)} \int_{C(t, \cdot)} \langle X, \nu \rangle ds \right) \\ &= \frac{1}{\lambda(t)^2} \left( \int_{C(t, \cdot)} \langle X, (\rho - \frac{2\sigma(t)}{\lambda(t)}) \nu \rangle ds \right) \\ &= \frac{\sigma(t)}{\lambda(t)^2} \left( \int_{C(t, \cdot)} \langle X, (\rho - \frac{2\sigma(t)}{\lambda(t)}) \nu \rangle \frac{ds}{\sigma(t)} \right). \end{aligned}$$

Hence  $(-\rho + 2h)\nu$  is the  $\mathbb{L}^2 \left( \frac{1}{\sigma(t)} ds \right)$  inverse gradient flow of  $\Psi$ .  $\square$

**Proposition 2.5.** *Under the RMCF, we have the following evolution of geometric quantities:*

$$\begin{cases} \partial_t v &= -\rho(\rho - 2h)v \\ \partial_t \partial_s &= \partial_s \partial_t + \rho(\rho - 2h) \partial_s \\ \partial_t T &= -\partial_s(\rho)\nu \\ \partial_t \nu &= \partial_s(\rho)T. \end{cases} \quad (3)$$

*Proof.* We differentiate equation (2) in  $u$ , and we get:

$$\begin{aligned} \partial_t \partial_u C &= \partial_u \partial_t C \\ &= -(\partial_u \rho)\nu + (-\rho + 2h)\partial_u \nu. \end{aligned}$$

We deduce:

$$\begin{aligned} 2v\partial_t v &= \partial_t v^2 = \partial_t \langle \partial_u C, \partial_u C \rangle = 2\langle \partial_t \partial_u C, \partial_u C \rangle \\ &= 2\langle -(\partial_u \rho)\nu + (-\rho + 2h)\partial_u \nu, \partial_u C \rangle \\ &= 2(-\rho + 2h)\langle \partial_u \nu, \partial_u C \rangle = 2v^2 \rho(-\rho + 2h). \end{aligned}$$

So we get the first part by identification. Also by the first computation

$$\begin{aligned} \partial_t \partial_s &= \partial_t \left( \frac{1}{v} \partial_u \right) = \frac{\rho(\rho - 2h)v}{v^2} \partial_u + \frac{1}{v} \partial_t \partial_u, \\ &= \rho(\rho - 2h) \partial_s + \partial_s \partial_t \end{aligned}$$

and

$$\begin{aligned} \partial_t T &= \partial_t \partial_s C = \partial_s \partial_t C + \rho(\rho - 2h) \partial_s C \\ &= -(\partial_s \rho)\nu + (-\rho + 2h)\partial_s \nu + \rho(\rho - 2h)\partial_s C = -(\partial_s \rho)\nu \end{aligned}$$

since  $\partial_t \langle \nu, \nu \rangle = 0$ ,  $\partial_t \nu$  is tangential. Also  $\partial_t \langle T, \nu \rangle = 0$ , so we get the last point from the previous one.  $\square$

**Proposition 2.6.** *Under the RMCF, we have the following evolution of the curvature:*

$$\partial_t \rho = \partial_s^2 \rho + \rho^2(\rho - 2h). \quad (4)$$

*Proof.* It is a direct consequence of the previous proposition,

$$\begin{aligned} \partial_t \rho &= \partial_t \langle T, \partial_s \nu \rangle = \langle T, \partial_t \partial_s \nu \rangle = \langle T, \partial_s \partial_t \nu + \rho(\rho - 2h)\partial_s \nu \rangle \\ &= \langle T, \partial_s(\partial_s(\rho)T) + \rho^2(\rho - 2h)T \rangle \\ &= \partial_s^2 \rho + \rho^2(\rho - 2h). \end{aligned}$$

□

**2.3. A priori estimate of geometric quantities.** We get the following evolution of geometrics quantities:

**Lemma 2.7.** *If the family of curves  $C(t, \cdot)$ , solution of the flow (2), remain simple for all  $t \in [0, T_c]$ , then we have for all  $t \in [0, T_c]$ :*

$$\begin{aligned} (1) \quad & \frac{d}{dt}\sigma(C_t) = -\int \rho^2 ds + \frac{4\pi\sigma(C_t)}{\lambda(D_t)}; \\ (2) \quad & \frac{d}{dt}\lambda(D_t) = -2\pi + \frac{2\sigma(C_t)^2}{\lambda(D_t)}; \\ (3) \quad & \frac{d}{dt}h(D(t)) = \frac{d}{dt} \frac{\sigma(C_t)}{\lambda(D_t)} \leq \frac{-12\pi^2}{\sigma(C_t)\lambda(D_t)} \leq 0. \end{aligned}$$

*Proof.* Using Gauss-Bonnet Theorem, i.e. for simple closed curve  $\int_0^{\sigma(C_t)} \rho ds = 2\pi$ , and (4) we have:

$$\begin{aligned} \frac{d}{dt}\sigma(C_t) &= \frac{d}{dt} \int_0^{2\pi} v(t, u) du = \int_0^{2\pi} -\rho(\rho - 2h)v du = \int_0^{\sigma(C_t)} -\rho(\rho - 2h) ds \\ &= -\int \rho^2 ds + \frac{4\pi\sigma(C_t)}{\lambda(D_t)}. \end{aligned}$$

For the second point, we have

$$\frac{d}{dt}\lambda(D_t) = \int_{C_t} \left\langle \frac{d}{dt}C(t, s), \nu \right\rangle ds = \int_{C_t} -(\rho - 2h) ds = -2\pi + \frac{2\sigma(C_t)^2}{\lambda(D_t)}.$$

Let us write  $\sigma_t = \sigma(C_t)$ ,  $\lambda_t = \lambda(D_t)$  and denote by a dot the derivation with respect to  $t$ ,

$$\begin{aligned} \frac{d}{dt} \frac{\sigma_t}{\lambda_t} &= \frac{1}{\lambda_t} (\dot{\sigma}_t - \frac{\sigma_t \dot{\lambda}_t}{\lambda_t}) = \frac{1}{\lambda_t} \left( -\int \rho^2 ds + \frac{4\pi\sigma_t}{\lambda_t} - \frac{\sigma_t}{\lambda_t} \left( -2\pi + \frac{2\sigma_t^2}{\lambda_t} \right) \right) \\ &\leq \frac{1}{\lambda_t} \left( -\frac{4\pi^2}{\sigma_t} + \frac{4\pi\sigma_t}{\lambda_t} - \frac{\sigma_t}{\lambda_t} \left( -2\pi + \frac{2\sigma_t^2}{\lambda_t} \right) \right) \\ &= \frac{-4\pi^2\lambda_t^2 + 6\pi\sigma_t^2\lambda_t - 2\sigma_t^4}{\lambda_t^3\sigma_t} \\ &= \frac{-2(\sigma_t^2 - 2\pi\lambda_t)(\sigma_t^2 - \pi\lambda_t)}{\sigma_t\lambda_t^3} \\ &\leq \frac{-12\pi^2}{\lambda_t\sigma_t} \leq 0, \end{aligned}$$

where we have used Cauchy-Schwartz inequality and Gauss-Bonnet Theorem in the second line, and the isoperimetric inequality in the last line.

□

**Lemma 2.8.** *If the family of curves  $C(t, \cdot)$ , solution of (2), remain convex for all  $t \in [0, T_c]$ , then the isoperimetric ratio is non-increasing, i.e. for all  $t \in [0, T_c]$ ,*

$$\frac{d}{dt} \frac{\sigma_t^2}{4\pi\lambda_t} \leq 0.$$

*Proof.*

$$\begin{aligned} \frac{d}{dt} \frac{\sigma_t^2}{4\pi\lambda_t} &= \frac{\sigma_t}{4\pi\lambda_t} (2\dot{\sigma}_t - \frac{\sigma_t \dot{\lambda}_t}{\lambda_t}) \\ &= \frac{\sigma_t}{4\pi\lambda_t} \left( -2 \int \rho^2 ds + \frac{8\pi\sigma_t}{\lambda_t} - \frac{\sigma_t}{\lambda_t} (-2\pi + \frac{2\sigma_t^2}{\lambda_t}) \right) \\ &= \frac{\sigma_t}{4\pi\lambda_t} \left( -2 \int \rho^2 ds + \frac{10\pi\sigma_t}{\lambda_t} - \frac{2\sigma_t^3}{\lambda_t^2} \right) \\ &\leq \frac{\sigma_t}{4\pi\lambda_t} \left( -2 \int \rho^2 ds + \frac{2\pi\sigma_t}{\lambda_t} \right) \end{aligned}$$

where we have use isoperimetric inequality in the last line. Let us now recall the convex Gage inequality which is proven in [3], and tells us that for convex  $C^2$  plane curves:

$$\frac{\pi\sigma_t}{\lambda_t} \leq \int \rho^2 ds. \quad (5)$$

Using this inequality in the above computation we get:

$$\frac{d}{dt} \frac{\sigma_t^2}{4\pi\lambda_t} \leq \frac{\sigma_t}{4\pi\lambda_t} \left( -\frac{2\pi\sigma_t}{\lambda_t} + \frac{2\pi\sigma_t}{\lambda_t} \right) = 0$$

□

**Lemma 2.9.** *If the family of curves  $C(t, \cdot)$ , remain simple along the flow (2), then the deficit of isoperimetry is non-increasing, i.e.:*

$$\frac{d}{dt} (\sigma_t^2 - 4\pi\lambda_t) \leq 0.$$

*If moreover the family of curves  $C(t, \cdot)$  remain convex for all  $t \in [0, T_c]$  then for all  $t \in [0, T_c]$  we have:*

$$\begin{aligned} (1) \quad &\frac{d}{dt} (\sigma_t^2 - 4\pi\lambda_t) \leq \frac{-2\pi}{\lambda_t} (\sigma_t^2 - 4\pi\lambda_t), \\ (2) \quad &0 \leq (\sigma_t^2 - 4\pi\lambda_t) \leq (\sigma_0^2 - 4\pi\lambda_0) \left( \frac{\left( -2\pi + \frac{2\sigma_0^2}{\lambda_0} \right) t + \lambda_0}{\lambda_0} \right)^{\frac{-2\pi}{-2\pi + \frac{2\sigma_0^2}{\lambda_0}}}. \end{aligned}$$

*Proof.* By direct computation, and after using Lemma 2.7 and similar computation, we have:

$$\frac{d}{dt} (\sigma_t^2 - 4\pi\lambda_t) = 2\sigma_t \dot{\sigma}_t - 4\pi \dot{\lambda}_t = 2\sigma_t \left( - \int \rho^2 ds + \frac{4\pi\sigma_t}{\lambda_t} \right) - 4\pi \left( -2\pi + \frac{2\sigma_t^2}{\lambda_t} \right)$$

$$\begin{aligned} &\leq 2\sigma_t \left( -\frac{4\pi^2}{\sigma_t} + \frac{4\pi\sigma_t}{\lambda_t} \right) - 4\pi \left( -2\pi + \frac{2\sigma_t^2}{\lambda_t} \right) \\ &\leq 0. \end{aligned}$$

If moreover the family of curves  $C(t, \cdot)$  remain convex, in the second line of the above computation, we can improve the inequality using (5) instead of Gauss-Bonnet Theorem, and we get for all  $t \in [0, T_c[$ :

$$\begin{aligned} \frac{d}{dt} (\sigma_t^2 - 4\pi\lambda_t) &\leq 2\sigma_t \left( -\frac{\pi\sigma_t}{\lambda_t} + \frac{4\pi\sigma_t}{\lambda_t} \right) - 4\pi \left( -2\pi + \frac{2\sigma_t^2}{\lambda_t} \right) \\ &\leq \frac{-2\pi}{\lambda_t} (\sigma_t^2 - 4\pi\lambda_t). \end{aligned}$$

Using Lemmas 2.8 , 2.7 and isoperimetric inequality we deduce that

$$6\pi \leq -2\pi + \frac{2\sigma_t^2}{\lambda_t} \leq \dot{\lambda}_t \leq -2\pi + \frac{2\sigma_t^2}{\lambda_t} \leq -2\pi + \frac{2\sigma_0^2}{\lambda_0},$$

so for all  $t \in [0, T_c[$

$$6\pi t + \lambda_0 \leq \lambda_t \leq \left( -2\pi + \frac{2\sigma_0^2}{\lambda_0} \right) t + \lambda_0.$$

Hence we get:

$$\frac{d}{dt} (\sigma_t^2 - 4\pi\lambda_t) \leq \frac{-2\pi}{\left( -2\pi + \frac{2\sigma_0^2}{\lambda_0} \right) t + \lambda_0} (\sigma_t^2 - 4\pi\lambda_t).$$

After integration we obtain for all  $t \in [0, T_c[$ :

$$0 \leq (\sigma_t^2 - 4\pi\lambda_t) \leq (\sigma_0^2 - 4\pi\lambda_0) \left( \frac{\left( -2\pi + \frac{2\sigma_0^2}{\lambda_0} \right) t + \lambda_0}{\lambda_0} \right)^{\frac{-2\pi}{-2\pi + \frac{2\sigma_0^2}{\lambda_0}}}.$$

□

We get the following corollary, which describes the shape of the curve  $C(t, \cdot)$  as  $t$  goes to infinity.

**Corollary 2.10.** *If the curves  $C(t, \cdot)$  are defined for all times  $t \geq 0$  and remain convex then*

$$\lim_{t \rightarrow \infty} \frac{\sigma_t^2}{\lambda_t} = 4\pi.$$

*After renormalization the curve  $\frac{1}{\sqrt{6t}}[C(t, \cdot) - c_{\text{int}}(t)]$  converge to the circle of center 0 and radius 1 for Hausdorff metric, where  $c_{\text{int}}(t)$  is a center of an inscribed circle of  $C(t, \cdot)$ .*



*Proof.* Using Lemma 2.7 and the isoperimetric inequality, we have  $\dot{\lambda}_t \geq 6\pi$ , so

$$\lambda_t \geq 6\pi t + \lambda_0.$$

Using the above Lemma 2.9 we get

$$0 \leq \frac{\sigma_t^2 - 4\pi\lambda_t}{\lambda_t} \leq \frac{(\sigma_0^2 - 4\pi\lambda_0) \left( \frac{\left( -2\pi + \frac{2\sigma_0^2}{\lambda_0} \right) t + \lambda_0}{\lambda_0} \right)^{\frac{-2\pi}{-2\pi + \frac{2\sigma_0^2}{\lambda_0}}}}{6\pi t + \lambda_0},$$

and the right hand side goes to 0 as  $t$  goes to infinity. For the second point, use again 2.7 and the computation above, to deduce that

$$\dot{\lambda}_t \underset{t \sim \infty}{\sim} 6\pi,$$

so

$$\lambda_t \underset{t \sim \infty}{\sim} 6\pi t.$$

Since  $\lim_{t \rightarrow \infty} \sigma_t^2 - 4\pi\lambda_t = 0$ , and using Bonnesen inequality, i.e.

$$\frac{\pi^2}{\lambda_t} (r_{\text{out}}(t) - r_{\text{int}}(t))^2 \leq \left( \frac{\sigma_t^2}{\lambda_t} - 4\pi \right) \quad (6)$$

where  $r_{\text{out}}(t), r_{\text{int}}(t)$  are respectively the outer and the inner radius of the curve  $C(t, \cdot)$ , we get that

$$(r_{\text{out}}(t) - r_{\text{int}}(t))^2 \leq \frac{\lambda_t}{\pi^2} \left( \frac{\sigma_t^2}{\lambda_t} - 4\pi \right).$$

Let  $c_{\text{int}}(t)$  be a center of an inscribed circle of  $C(t, \cdot)$ , and  $c_{\text{out}}(t)$  be a center of a circumscribed circle of  $C(t, \cdot)$ . Then since  $B(c_{\text{int}}(t), r_{\text{int}}(t)) \subset D(t) \subset B(c_{\text{out}}(t), r_{\text{out}}(t))$  we have by Lemma 2.9, and isoperimetric inequality that there exist two positive constants  $C > 0$  and  $\gamma \in ]0, \frac{1}{6}]$  such that ,

$$|c_{\text{int}}(t) - c_{\text{out}}(t)| \leq r_{\text{out}}(t) - r_{\text{int}}(t) \leq Ct^{-\gamma}$$

and

$$r_{\text{int}}(t) \leq \sqrt{\frac{\lambda_t}{\pi}} \leq r_{\text{out}}(t).$$

For two compact set  $A, B \subset \mathbb{R}^2$  and  $\epsilon \geq 0$  we define

$$A_\epsilon := \{x \in \mathbb{R}^2, d(x, A) \leq \epsilon\}$$

and the Hausdorff distance between  $A$  and  $B$  by

$$d_{\mathcal{H}}(A, B) := \inf\{r > 0, A \subset B_r \text{ and } B \subset A_r\}.$$

Since  $B(c_{\text{out}}(t), r_{\text{out}}(t)) \subset B(c_{\text{int}}(t), r_{\text{int}}(t))_{2(r_{\text{out}}(t) - r_{\text{int}}(t))}$  we easily derive that

$$d_{\mathcal{H}}(B(c_{\text{int}}(t), r_{\text{int}}(t)), B(c_{\text{out}}(t), r_{\text{out}}(t))) \leq 2(r_{\text{out}}(t) - r_{\text{int}}(t)) \leq 2Ct^{-\gamma}$$

and

$$d_{\mathcal{H}}(D(t), B(c_{\text{int}}(t), r_{\text{int}}(t))) \leq 2(r_{\text{out}}(t) - r_{\text{int}}(t)) \leq 2Ct^{-\gamma}.$$

So by Lemma 2.9,

$$\begin{aligned} d_{\mathcal{H}} \left( \frac{D(t) - c_{\text{int}}(t)}{\sqrt{6t}}, B(0, 1) \right) &= \frac{1}{\sqrt{6t}} d_{\mathcal{H}}(D(t), B(c_{\text{int}}(t), \sqrt{6t})) \\ &\leq \frac{1}{\sqrt{6t}} \left( d_{\mathcal{H}}(D(t), B(c_{\text{int}}(t), r_{\text{int}}(t))) + d_{\mathcal{H}}(B(c_{\text{int}}(t), r_{\text{int}}(t)), B(c_{\text{int}}(t), \sqrt{6t})) \right) \\ &\leq \frac{1}{\sqrt{6t}} \left( 2Ct^{-\gamma} + |r_{\text{int}} - \sqrt{6t}| \right) \xrightarrow{t \rightarrow \infty} 0 \end{aligned}$$

Similarly we have that  $\frac{1}{\sqrt{6t}}[C(t, \cdot) - c_{\text{int}}(t)]$  converge to the circle of radius 1 and center 0 for the Hausdorff metric.  $\square$

**2.4. Preserving the convexity, and lower bound of the curvature.** We will now show that if the starting curve is strictly convex and simple then it remains convex and simple during the flow (2) until the lifetime. In a second time we will show that the lifetime is infinite using intensively some ideas developed in [6].

It is possible to parametrize a strictly convex curve in  $\mathbb{R}^2$  using the angle  $\theta$  between the tangent line and the  $x$  axis, i.e.  $T = (\cos(\theta), \sin(\theta))$ . Normally, the angle  $\theta$  depends on  $u$  and  $t$ . Following [5] and [6], after adding to the flow (2) a tangential perturbation, we do not perturb the shape of the curve, and it is possible to find a tangential intensity so that the parameter  $\theta$  does not depend on the time. This change of coordinate will make the equation more simple to understand (since operators  $\partial_{\theta}$  and  $\partial_t$  will commute, contrarily to  $\partial_s$  and  $\partial_t$  as shown in (12)).

Let us quickly recall this fact. We define (since the shape of the curve does not change we will keep the notation, only the parametrisation changes):

$$\partial_t C(t, u) = (-\rho_t(C(t, u)) + 2h(D(t, \cdot)))\nu + \alpha(u, t)T. \quad (7)$$

Differentiating (7) with respect to  $u$  and using  $\partial_u T = -v\rho\nu$  together with  $\partial_u \nu = v\rho T$  we get

$$\partial_t T = \left( -\frac{\partial_u \rho}{v} - \alpha(u, t)\rho \right) \nu$$

and

$$\partial_t v = -v\rho^2 + 2h\nu\rho + \partial_u \alpha.$$

To make  $\theta$  and hence  $T$  independent of  $t$  we take  $\alpha = -\frac{\partial_u \rho}{v\rho}$ . Differentiating  $\partial_t T$  with respect to  $u$ , we get that  $\partial_t(v\rho) = 0$ . Since  $\partial_{\theta} T = -\nu$ , we have by chain rule  $\frac{\partial_u \partial_t T}{\partial \theta} = -\nu$ , hence  $\frac{\partial_u}{\partial \theta} = \frac{1}{v\rho}$  and we deduce that

$$\frac{\partial}{\partial \theta} = \frac{1}{v\rho} \frac{\partial}{\partial u},$$

and so  $d\theta = \rho v du$ .

Since  $\partial_t(v\rho) = 0$  we have:

$$\partial_t \rho = -\frac{(\partial_t v)\rho}{v} = \rho^3 - 2h\rho^2 + \frac{\rho}{v} \partial_u \left( \frac{\partial_u \rho}{v\rho} \right) = \rho^3 - 2h\rho^2 + \frac{\rho}{v} \partial_u (\partial_{\theta} \rho)$$

$$= \rho^2 \partial_\theta (\partial_\theta \rho) + \rho^2 (\rho - 2h)$$

We summarize the result in the next lemma.

**Lemma 2.11.** *When the curvature remains positive, equation (7) for the RMCF yields equation*

$$\partial_t \rho = \rho^2 \partial_\theta^2 \rho + \rho^2 (\rho - 2h) \quad (8)$$

for the curvature.

**Lemma 2.12.** *A strictly convex simple curve  $C(0, \cdot)$ , that evolves according to (7) remains strictly convex and simple during the existence of the flow. Moreover*

$$\rho(\theta, t) \geq \rho_{\inf}(0) e^{-h_0^2 t}.$$

*Proof.* Let  $\mathcal{Q}(\theta, t) = \rho(\theta, t) e^{\mu t}$  for a constant  $\mu$  that will be chosen later, then  $\mathcal{Q}$  will satisfy the following equation:

$$\partial_t \mathcal{Q} = \rho^2 \frac{\partial^2}{\partial \theta^2} \mathcal{Q} + \mathcal{Q} (\rho^2 - 2h\rho + \mu). \quad (9)$$

The reaction term in the above equation is quadratic in  $\rho$ , and the discriminant is  $4(h^2 - \mu)$ . Note that the quantity  $h$  in this equation is the same as in (2), since the geometric quantities are the same for this equation and (7). Also by Lemma 2.7 ((\*) suppose for the moment that the curve remains simple),  $h$  is non-increasing, and

$$4(h^2 - \mu) \leq 4(h_0^2 - \mu).$$

So choosing  $\mu > h_0^2$  such that this discriminant is negative, the coefficient of  $\mathcal{Q}$  remains positive. We will apply the maximum principle for this equation. Let  $\mathcal{Q}_{\inf}(t) := \inf\{\mathcal{Q}(\theta, t), 0 \leq \theta \leq 2\pi\}$ . The proof is by contradiction, suppose that there exist  $0 < \eta < \mathcal{Q}_{\inf}(0)$  and  $t > 0$  such that  $\mathcal{Q}_{\inf}(t) = \eta$ , let  $t_0$  be the first time such that  $\mathcal{Q}_{\inf}(t_0) = \eta$ . This minimum is achieved at some point  $\theta_0$ , and at this point:

$\partial_t \mathcal{Q}(\theta_0, t_0) \leq 0$ ,  $\frac{\partial^2}{\partial \theta^2} \mathcal{Q}(\theta_0, t_0) \geq 0$ , and  $\mathcal{Q}(\theta_0, t_0) = \eta$ . This is in contradiction with Equation (9). Hence for all  $0 < t$ ,  $\mathcal{Q}_{\inf}(t) \geq \mathcal{Q}_{\inf}(0)$  and

$$\rho_{\inf}(t) \geq \rho_{\inf}(0) e^{-\mu t},$$

where  $\rho_{\inf}(t) = \inf\{\rho(\theta, t), 0 \leq \theta \leq 2\pi\}$  so  $\rho(\theta, t) \geq \rho_{\inf}(0) e^{-\mu t}$  for all  $\mu > h_0^2$ . Hence if  $C(0, \cdot)$  is strictly convex then  $C(t, \cdot)$  remains strictly convex at any time  $t$  up to which it is defined.

(\*)Remark that in the above proof we have used Lemma 2.7, all along the duration of the equation (7), without showing that the curve remains simple. But the argument is the same, by contradiction, if the first time  $T_s$  when the curve stops to be simple, occurs strictly before the maximal lifetime of (7), i.e.  $0 < T_s \leq t^* < T_c$  by the same computation as above the curvature  $\rho$  is positive until  $T_s$ ,  $\rho$  and all its derivatives are bounded in  $[0, t^*]$ , so there exist a limiting curve as  $t$  goes to  $T_s$ , that is smooth and has positive curvature (Arzelà-Ascoli theorem). Using Lemma 4.1.1,

and Theorem 4.1.4 in [6] we know that the positive curvature and the conservation of

$$\int_0^{2\pi} \frac{\cos(\theta)}{\rho} d\theta = \int_0^{2\pi} \frac{\sin(\theta)}{\rho} d\theta = 0$$

during the flow characterize the simple close strictly convex curve. So the limiting curve is simple and strictly convex, this contradicts the non simplicity at this time.  $\square$

**Corollary 2.13.** *A strictly convex simple curve  $C(0, \cdot)$ , that evolves according to (7) remains convex and simple during the existence of the flow. Moreover*

$$\rho(\theta, t) \geq \frac{1}{\frac{1}{\rho_{\inf}(0)} + \frac{\sigma_0^2}{6\pi^2\lambda_0} \sqrt{24\pi^2 t + 4\pi\lambda_0}}.$$

*Proof.* Using Lemma 2.12, we have that  $\rho > 0$  so we can define  $W := e^{-\frac{1}{\rho} + \int_0^t 2h_s ds}$ . We compute

$$\begin{aligned} \partial_t W &= (\partial_t e^{-\frac{1}{\rho}}) e^{\int_0^t 2h_s ds} + 2hW \\ &= \left( \frac{\partial_t \rho}{\rho^2} + 2h \right) W \\ &= (\partial_\theta^2 \rho + \rho) W \end{aligned}$$

We will apply the maximum principle for this equation. Define

$$W_{\inf}(t) := \inf\{W(\theta, t), 0 \leq \theta \leq 2\pi\}$$

The proof is by contradiction, suppose that there exists  $0 < \eta < W_{\inf}(0)$  and  $t > 0$  such that  $W_{\inf}(t) = \eta$ . Let  $t_0 > 0$  be the first time such that  $W_{\inf}(t_0) = \eta$ . This minimum is achieved at some point  $\theta_0$ , and at this point (since  $W$  is a non-decreasing function in  $\rho$ ):

$\partial_t W(\theta_0, t_0) \leq 0$ ,  $\frac{\partial^2}{\partial^2 \theta} W(\theta_0, t_0) \geq 0$ , and  $W(\theta_0, t_0) = \eta$ , so  $\frac{\partial^2}{\partial^2 \theta} \rho(\theta_0, t_0) \geq 0$ . This is in contradiction with the equation satisfied by  $W$ . Hence for all  $0 < t$ ,  $W_{\inf}(t) \geq W_{\inf}(0)$ , so

$$e^{-\frac{1}{\rho} + \int_0^t 2h_s ds} \geq e^{-\frac{1}{\rho_{\inf}(0)}}.$$

By Lemma 2.8, we have

$$h(t) := \frac{\sigma_t}{\lambda_t} \leq \frac{\sigma_0^2}{\lambda_0 \sigma_t}.$$

By isoperimetric inequality we have  $\sqrt{4\pi\lambda_t} \leq \sigma_t$  hence by Lemma 2.7 we have  $\sqrt{4\pi(6\pi t + \lambda_0)} \leq \sigma_t$  and

$$h(t) \leq \frac{\frac{\sigma_0^2}{\lambda_0}}{\sqrt{4\pi(6\pi t + \lambda_0)}}.$$

This yield

$$\int_0^t 2h(s) ds \leq \frac{\sigma_0^2}{6\pi^2\lambda_0} [\sqrt{24\pi^2t + 4\pi\lambda_0} - \sqrt{4\pi\lambda_0}],$$

and

$$\rho(t) \geq \frac{1}{\frac{1}{\rho_{\inf}(0)} + \frac{\sigma_0^2}{6\pi^2\lambda_0} \sqrt{24\pi^2t + 4\pi\lambda_0}}.$$

□

**2.5. Upper bound of the curvature and long time existence of the flow for strictly convex simple initial curve.** We will now show that (2) exists for all times if  $C(0, \cdot)$  is a strictly convex simple curve, by first establishing a uniform bound (depending on time horizon) of the maximum of curvature. Similarly to [6] we define the pseudo-median of the curvature:

$$\rho^*(t) = \sup\{\beta, \rho(\theta, t) > \beta \text{ on some interval of length } \pi\}.$$

**Lemma 2.14.** [6] *If a convex closed plane curve encloses an area  $\lambda$  and has length  $\sigma$  then  $\rho^* \leq \frac{\sigma}{\lambda}$ .*

Following the above lemma we have:

**Corollary 2.15.** *If the starting curve  $C(0, \cdot)$  is simple and strictly convex, and the solution to Equation (2) exists for all  $t \in [0, T[$  then for all  $t \in [0, T[$ ,*

$$\rho^*(t) \leq h \leq h_0.$$

*Proof.* We use Lemma 2.12 to get the convexity until  $T$ , Lemma 2.7 for  $h$  non-increasing, and the above Lemma. □

**Lemma 2.16** (Entropic estimate). *If the starting curve  $C(0, \cdot)$  is strictly convex and simple, and the solution of equation (2) is defined for all  $t \in [0, T[$ , then*

$$\text{Ent}(t) := \int_0^{2\pi} \log(\rho(\theta, t)) d\theta \tag{10}$$

*is non-increasing on  $[0, T[$ .*

*Moreover we have the following estimates:*

$$2\pi [\log \rho_{\inf}(0) - h_0^2 t] \leq \text{Ent}(t) \leq \text{Ent}(0) + \frac{\pi \rho_{\inf}^2(0)}{2h_0^2} [e^{-2h_0^2 t} - 1]$$

*and there exists five explicit constants  $C_0, C_1, \tilde{C}_0, \tilde{C}_1, \tilde{C}_2$ , only depending on the geometry of the initial curve, such that*

$$-2\pi \left[ \log(\tilde{C}_0 + \sqrt{\tilde{C}_1 t + \tilde{C}_2}) \right] \leq \text{Ent}(t) \leq \text{Ent}(0) - \frac{2\pi}{C_1} \left[ \log\left(\frac{C_1 t + C_0}{C_0}\right) \right].$$

*Proof.* The proof is an adaptation of the proof in [6]. We will just point the differences. After an integration by part we have:

$$\frac{d}{dt} \int_0^{2\pi} \log(\rho(\theta, t)) d\theta = \int_0^{2\pi} - \left( \frac{\partial}{\partial \theta} \rho \right)^2 + \rho(\rho - 2h) d\theta.$$

Let write the open set  $U = \{\theta, \rho(\theta, t) > \rho^*(t)\}$  as an union of disjoint interval  $I_i$ , by definition of the median the length of all  $I_i$  is smaller than  $\pi$ , and

$$\begin{aligned} \int_{I_i} - \left( \frac{\partial}{\partial \theta} \rho \right)^2 + \rho(\rho - 2h) d\theta &= \int_{I_i} - \left( \frac{\partial}{\partial \theta} (\rho - \rho^*) \right)^2 + \rho^2 d\theta - 2h \int_{I_i} \rho d\theta \\ &\leq \int_{I_i} -(\rho - \rho^*)^2 + \rho^2 d\theta - 2h \int_{I_i} \rho d\theta \\ &= \int_{I_i} 2\rho\rho^* - (\rho^*)^2 d\theta - 2h \int_{I_i} \rho d\theta \\ &\leq (2\rho^* - 2h) \int_{I_i} \rho d\theta - (\rho^*)^2 \int_{I_i} d\theta, \end{aligned}$$

where in the second line we used Wirtinger [6] inequality (recall that on the boundary of  $I_i$ ,  $\rho = \rho^*$ ).

On the complement of  $U$  we have:

$$\begin{aligned} \int_{U^c} - \left( \frac{\partial}{\partial \theta} \rho \right)^2 + \rho(\rho - 2h) d\theta &\leq (\rho^* - 2h) \int_{U^c} \rho d\theta \\ &\leq (2\rho^* - 2h) \int_{U^c} \rho d\theta - \rho^* \int_{U^c} \rho d\theta. \end{aligned}$$

Hence using Lemma 2.14, and  $\rho^*(t) \geq \rho_{\inf}(t)$  we get

$$\begin{aligned} \frac{d}{dt} \int_0^{2\pi} \log(\rho(\theta, t)) d\theta &\leq (2\rho^* - 2h) \int_0^{2\pi} \rho d\theta - \rho^* \int_{U^c} \rho d\theta - (\rho^*)^2 \int_U d\theta \\ &\leq -2\pi \rho_{\inf}^2(t). \end{aligned}$$

So the first part of the lemma is proved. Lemma 2.12 yields after integration:

$$\int_0^{2\pi} \log(\rho(\theta, t)) d\theta \leq \int_0^{2\pi} \log(\rho(\theta, 0)) d\theta + \frac{\pi \rho_{\inf}^2(0)}{h_0^2} \left[ e^{-2h_0^2 t} - 1 \right],$$

and Corollary 2.13 yields the second estimate.

For the lower bound use the bounds of Lemma 2.12 and 2.13.  $\square$

**Proposition 2.17.** *If the starting curve  $C(0, \cdot)$  is strictly convex and simple, and the solution of equation (2) is defined for all  $t \in [0, T]$ , then there exists a constant  $C_0$  that depends only on the initial curve such that:*

$$\int_0^{2\pi} \left( \frac{\partial}{\partial \theta} \rho \right)^2 d\theta \leq \int_0^{2\pi} (\rho)^2 d\theta - 4h_t \int_0^{2\pi} \rho d\theta + C_0.$$

*Proof.* By Lemma 2.12 we know that the curvature  $\rho$  remains positive during the existence of the flow, so we can compute:

$$\begin{aligned}
 & \frac{d}{dt} \int_0^{2\pi} \left( \rho^2 - \left( \frac{\partial}{\partial \theta} \rho \right)^2 - 4h\rho \right) d\theta \\
 &= 2 \int_0^{2\pi} \frac{d\rho}{dt} (\rho + (\partial_\theta^2 \rho) - 2h) d\theta - 4 \frac{dh}{dt} \int_0^{2\pi} \rho d\theta \\
 &= 2 \int_0^{2\pi} \left( \frac{\partial_t \rho}{\rho} \right)^2 d\theta - 4 \frac{dh}{dt} \int_0^{2\pi} \rho d\theta \\
 &\geq -4 \frac{dh}{dt} \int_0^{2\pi} \rho d\theta \\
 &\geq \frac{48\pi^2}{\lambda\sigma} \int_0^{2\pi} \rho d\theta > 0,
 \end{aligned}$$

where we have used an integration by part on the second line, the equation of the curvature at the third line, and Lemma 2.7 at the last line.

Integrating the last inequality we get for all  $t \in [0, T[$ ,

$$\int_0^{2\pi} \left( \rho^2 - \left( \frac{\partial}{\partial \theta} \rho \right)^2 - 4h\rho \right) \Big|_t d\theta \geq \left( \int_0^{2\pi} \left( \rho^2 - \left( \frac{\partial}{\partial \theta} \rho \right)^2 - 4h\rho \right) d\theta \right) \Big|_0 = -C_0$$

We obtained the last inequality using (5) and the upper bound of the volume during the flow (Lemma 2.7).  $\square$

**Proposition 2.18.** *If the starting curve  $C(0, \cdot)$  is strictly convex and simple, and the solution of equation (2) is defined for all  $t \in [0, T[$  then if  $T < \infty$*

$$M = \sup\{\rho(\theta, t), (\theta, t) \in \mathbb{R}/2\pi\mathbb{Z} \times [0, T[ \}$$

*is bounded.*

*Proof.* For  $t < T$  let

$$M_t = \sup\{\rho(\theta, s), (\theta, s) \in \mathbb{R}/2\pi\mathbb{Z} \times [0, t] \}.$$

There exists  $(\theta_1, t_1) \in \mathbb{R}/2\pi\mathbb{Z} \times [0, t]$  such that  $M_t = \rho(\theta_1, t_1)$ . Then for all  $\theta_2 \in \mathbb{R}/2\pi\mathbb{Z}$  we have:

$$\begin{aligned}
 |\rho(\theta_1, t_1) - \rho(\theta_2, t_1)| &= \left| \int_{\theta_2}^{\theta_1} \frac{\partial}{\partial \theta} \rho d\theta \right| \\
 &\leq \sqrt{\int_0^{2\pi} \left( \frac{\partial}{\partial \theta} \rho \right)^2 d\theta} \sqrt{|\theta_1 - \theta_2|} \\
 &\leq \sqrt{2\pi M_t^2 + C_0} \sqrt{|\theta_1 - \theta_2|} \\
 &\leq M_t \sqrt{2\pi + \frac{|C_0|}{M_t^2}} \sqrt{|\theta_1 - \theta_2|},
 \end{aligned}$$

where we have used Proposition 2.17 at the third line, and  $|\theta_1 - \theta_2|$  is the usual distance in  $\mathbb{R}/2\pi\mathbb{Z}$ . We also have,

$$M_t \geq \rho_{sup}(0).$$

So

$$\rho(\theta_1, t_1) - \rho(\theta_2, t_1) \leq C_1 M_t \sqrt{|\theta_1 - \theta_2|},$$

where  $C_1 := \sqrt{2\pi + \frac{|C_0|}{\rho_{sup}^2(0)}}$ .

It follows that for all  $\theta_2$ :

$$\rho(\theta_2, t_1) \geq M_t - C_1 M_t \sqrt{|\theta_1 - \theta_2|}$$

and

$$\begin{aligned} & \int_0^{2\pi} \log(\rho(\theta, t_1)) d\theta \\ &= \int_{|\theta_1 - \theta| \leq (\frac{1}{2C_1})^2} \log(\rho(\theta, t_1)) d\theta + \int_{|\theta_1 - \theta| \geq (\frac{1}{2C_1})^2} \log(\rho(\theta, t_1)) d\theta \\ &\geq \log\left(\frac{M_t}{2}\right) \frac{1}{2C_1^2} + \left(2\pi - \frac{1}{2C_1^2}\right) \log\left(\rho_{inf}(0)e^{-h_0^2 T}\right) \end{aligned}$$

Use Lemma 2.16 we obtain for all  $t \in [0, T[$

$$\log(M_t) \leq C_2(T)$$

where

$$C_2(T) = 2C_1^2 \left( \int_0^{2\pi} \log(\rho(\theta, 0)) d\theta - \left(2\pi - \frac{1}{2C_1^2}\right) (\log(\rho_{inf}(0)) - h_0^2 T) \right)$$

is a function that only depends on the final time and the initial curve. So  $M = \sup \{\rho(\theta, t), (\theta, t) \in \mathbb{R}/2\pi\mathbb{Z} \times [0, T]\}$  is bounded for  $T < \infty$ .  $\square$

**Proposition 2.19.** *If the starting curve  $C(0, \cdot)$  is strictly convex, and the solution of equation (2) is defined for all  $t \in [0, T[$  then if  $T < \infty$  then for all  $n \in \mathbb{N}$*

$$M^{(n)} = \sup \left\{ \frac{\partial^n \rho}{\partial^n \theta}(\theta, t), (\theta, t) \in \mathbb{R}/2\pi\mathbb{Z} \times [0, T[ \right\}$$

is bounded.

*Proof.* Following [6] section 4.4, we will first prove that  $\frac{\partial \rho}{\partial \theta}$  is bounded along the flow. By direct computation,  $\frac{\partial \rho}{\partial \theta}$  satisfies:

$$\frac{\partial}{\partial t} \frac{\partial \rho}{\partial \theta} = \rho^2 \partial_\theta^2 \left( \frac{\partial \rho}{\partial \theta} \right) + 2\rho \left( \frac{\partial \rho}{\partial \theta} \partial_\theta \left( \frac{\partial \rho}{\partial \theta} \right) \right) + [3\rho^2 - 4h\rho] \frac{\partial \rho}{\partial \theta}$$

By Lemma 2.7 and Proposition 2.18,  $[3\rho^2 - 4h\rho]$  is bounded so by the maximum principle  $\frac{\partial \rho}{\partial \theta}$  is bounded for all  $t \in [0, T[$ . (this is easier than in the proof of 2.18).



Since the equation for  $\frac{\partial^2 \rho}{\partial^2 \theta}$  contains a quadratic term it seems not clear to directly use the maximum principle. To control it we will proceed as follows.

With the same computation as in [6] Lemma 4.4.2 we see that modulo a additional term that comes from the non local term  $h$ , which is bounded, we show that  $\int \left( \frac{\partial^2 \rho}{\partial^2 \theta} \right)^4 d\theta$  is bounded on finite intervals of time (i.e. when  $T < \infty$ ). Let us prove this property. To present the difference with the computation in [6], we integrate by part. We get:

$$\begin{aligned} \frac{\partial}{\partial t} \int_0^{2\pi} \left( \frac{\partial^2 \rho}{\partial^2 \theta} \right)^4 d\theta &= 4 \int_0^{2\pi} \partial_\theta^2 (\rho^2 \partial_\theta^2 \rho + \rho^2 (\rho - 2h)) (\partial_\theta^2 \rho)^3 d\theta \\ &= -12 \int_0^{2\pi} (\rho^2 \partial_\theta^3 \rho + 2\rho \partial_\theta \rho \partial_\theta^2 \rho + (3\rho^2 - 4h\rho) \partial_\theta \rho) (\partial_\theta^3 \rho) (\partial_\theta^2 \rho)^2 d\theta. \end{aligned}$$

To simplify notations let us use  $\rho' := \partial_\theta \rho$ , in the above computation:

$$\begin{aligned} \frac{\partial}{\partial t} \int_0^{2\pi} (\rho'')^4 d\theta &= -12 \int_0^{2\pi} \left( \rho^2 (\rho'')^2 (\rho''')^2 + 2\rho \rho' (\rho'')^3 \rho''' \right. \\ &\quad \left. + 3\rho^2 \rho' (\rho'')^2 \rho''' - 4h\rho \rho' (\rho'')^2 \rho''' \right) d\theta. \end{aligned}$$

We use the inequality  $ab \leq \frac{1}{4\epsilon} a^2 + \epsilon b^2$  to bound the three last terms by the first one and some additional terms, after choosing  $\epsilon$  to control the sign of the first term. We obtain that there exist  $C_1, C_2, C_3$  such that:

$$\begin{aligned} \frac{\partial}{\partial t} \int_0^{2\pi} (\rho'')^4 d\theta &\leq C_1 \int_0^{2\pi} (\rho')^2 (\rho'')^4 d\theta + C_2 \int_0^{2\pi} (\rho)^2 (\rho')^2 (\rho'')^4 d\theta \\ &\quad + C_3 \int_0^{2\pi} (\rho')^2 (\rho'')^2 d\theta \end{aligned}$$

where the constant  $C_3$  depends on  $h(0)$  which is the maximum of  $h$  by Lemma 2.7. Since  $\rho$  is bounded by proposition 2.18 and  $\rho'$  is bounded we deduce from the above inequality and Cauchy-Schwarz inequality that  $\int_0^{2\pi} (\rho'')^4 d\theta$  remains bounded on  $[0, T[$ .

By the same kind of computation, we will show that  $\int \left( \frac{\partial^3 \rho}{\partial^3 \theta} \right)^2 d\theta$  remains bounded for  $t \in [0, T[$  (when  $T < \infty$ ). After a integration by part, we have:

$$\begin{aligned} \frac{d}{dt} \int (\rho''')^2 d\theta &= 2 \int (\rho''') (\rho^2 \rho'' + \rho^2 (\rho - 2h))''' d\theta \\ &= -2 \int (\rho''''') (\rho^2 \rho'' + \rho^2 (\rho - 2h))'' d\theta \\ &= -2 \int (\rho''''') \left( \rho^2 \rho'''' + 2\rho \rho' \rho''' + 2\rho (\rho'')^2 + 2(\rho')^2 \rho'' + 3\rho^2 \rho'' \right. \\ &\quad \left. + 6\rho (\rho')^2 - 4h(\rho')^2 - 4h\rho (\rho'') \right) d\theta. \end{aligned}$$

Using again the inequality  $ab \leq \frac{1}{4\epsilon}a^2 + \epsilon\beta^2$  for a well choosed  $\epsilon$  to bound the seven last terms by the first one and some additional terms, we get that there exist  $C_1, C_2, C_3, \dots, C_6, C_7$  (that all depend on  $\epsilon$ , and  $C_6, C_7$  depend also on  $h_0$ , the upper bound of  $h$  by Lemma 2.7) such that:

$$\begin{aligned} & \frac{d}{dt} \int (\rho''')^2 d\theta \\ & \leq C_1 \int (\rho\rho''')^2 d\theta + C_2 \int (\rho'')^4 d\theta + C_3 \int \left( \frac{(\rho')^2 \rho''}{\rho} \right)^2 d\theta \\ & + C_4 \int (\rho\rho'')^2 d\theta + C_5 \int (\rho')^4 d\theta + C_6 \int \left( \frac{(\rho')^2}{\rho} \right)^2 d\theta + C_7 \int (\rho'')^2 d\theta. \end{aligned}$$

Since on finite intervals of time,  $\rho$  is bounded by Lemma 2.18 and by the computation above  $\rho'$  and  $\int (\rho'')^4 d\theta$  are bounded, and the lower bound  $\rho \geq \rho_{\inf}(0)e^{-h_0^2 t}$  (Lemma 2.12), using Cauchy-Swartz inequality we get for other constants (that depend on the time horizon  $T$ ):

$$\frac{d}{dt} \int (\rho''')^2 d\theta \leq C_0 \int (\rho''')^2 d\theta + C_2.$$

Hence  $\int \left( \frac{\partial^3 \rho}{\partial \theta^3} \right)^2 d\theta$  remains bounded for  $t \in [0, T[$ , and so  $\frac{\partial^2 \rho}{\partial \theta^2}$  is bounded, using fundamental calculus theorem, or Sobolev inequality in  $\mathbb{S}^1$ .

For all  $n \geq 3$  the equation for  $\frac{\partial^n \rho}{\partial \theta^n} =: \rho^{(n)}$  writes

$$\begin{aligned} \frac{d}{dt} \rho^{(n)} &= \rho^2 \rho^{(n+2)} + 2n\rho\rho' \rho^{(n+1)} + [2\rho\rho'' + 3\rho^2 - 4h\rho \\ & + (n)(n-1)(\rho' + \rho\rho'')] \rho^{(n)} + P(h, \rho, \rho', \dots, \rho^{(n-1)}) \end{aligned}$$

where  $P(h, \rho, \rho', \dots, \rho^{(n-1)})$  is a polynomial in  $(h, \rho, \rho', \dots, \rho^{(n-1)})$ . Since  $\rho, \rho', \rho''$  are bounded by the computation above and  $h$  is bounded by Lemma 2.7, we can apply the maximum principle to get an exponential bound for  $\rho'''$ , so  $\rho'''$  is bounded (when  $T$  is finite). Using the above equation for  $\rho^{(n)}$ , we get by induction and maximum principle that for all  $n$ ,  $\rho^{(n)}$  is uniformly bounded on  $[0, T[$  when  $T < \infty$ .  $\square$

### 2.6. Proof of Theorem 2.3.

We prove the long time existence of the flow. Assume that the starting curve  $C(0, \cdot)$  is simple and strictly convex, and the flow exists for all  $t \in [0, T[$ . Then by Lemma 2.12 we know that the solution of (7) remains convex and simple during the flow. Since the solution of (7) has similar shape as the solution of (2) (just the parametrisation changes) we know that the solution of (2) remains convex and simple, so the quantity  $h$  remains defined for all  $t \in [0, T[$ . Using Lemma 2.7 we

get that the quantity  $h$  remains bounded as soon as the flow exists. By Propositions 2.18 and 2.19 we know that  $\rho$  and all spacial derivatives of  $\rho$  are bounded in  $[0, T]$ , if  $T < \infty$ , hence the same for time derivative of  $\rho$ . So by Arzelà-Ascoli Theorem,  $\rho$  converges to a  $C^\infty$  function  $\rho(T, \cdot)$  as  $t \rightarrow T$ . Using equation (2) there exists a limiting curve  $C(T, \cdot)$  and this limiting curve is associated to  $\rho(T, \cdot)$  (in the sense of Lemma 4.1.1 in [6]), so  $C(T, \cdot)$  is strictly convex and simple. By the small time existence if  $T < \infty$ , we can extend the time interval on which the solution is defined using the solution that starts at  $C(T, \cdot)$ . This proves that the solution of (2) starting with at a simple strictly convex curve exists for all time. By Lemma 2.8, the isoperimetric ratio is non-increasing. It is in fact decreasing until the curve becomes a circle (take strict inequality in the isoperimetric inequality in the proof of Lemma 2.8), but if it would becomes a circle in finite time we could reverse the flow and get that the starting curve is a circle. So the isoperimetric ratio is decreasing unless if the starting curve is circular. Using Lemma 2.9 we get that the deficit of isoperimetry converges to 0 polynomially, and Corollary 2.10 shows that the family of curves becomes more and more circular, and the isoperimetric ratio decreases to  $4\pi$ . Also this corollary yields, with Bonnesen inequality, the convergence after normalization to a circle, i.e.  $\frac{1}{\sqrt{6t}}[C(t, \cdot) - c_{\text{int}}(t)]$  converges to circle of center 0 and radius 1 for Hausdorff metric, where  $c_{\text{int}}(t)$  is a center of a inscribed circle of  $C(t, \cdot)$ . □

**Remark 2.20.** *If the starting curve is convex and not simple, as in Figure 2(a), the flow is not well defined. And if the starting curve is simple but non convex, the existence in long time is not automatic as in Figure 2 b).*

### 3. THE STOCHASTIC RENORMALIZED MEAN CURVATURE FLOW (SRMCF) IN $\mathbb{R}^2$

**3.1. Equations of geometric quantities along the stochastic flow of curves in  $\mathbb{R}^2$ .**  
Let  $u \in \mathbb{T} \mapsto C_0(u)$  be a simple  $C^{2+\alpha}$  closed curve in  $\mathbb{R}^2$ .

**Definition 3.1.** *Let  $C(t, u) : \mathbb{R}^+ \times \mathbb{T} \rightarrow \mathbb{R}^2$  be a family of simple closed curves. We say that  $C(t, \cdot)$  evolves under the renormalized stochastic mean curvature flow, shortly (SRMCF), if it satisfies the following equation*

$$\begin{cases} d_t C(t, u) &= ([-\rho_t(C(t, u)) + 2h_t]dt + \sqrt{2}dB_t) \nu_t(C(t, u)) \\ C(0, u) &= C_0(u) \end{cases} \quad (11)$$

where  $h_t$ ,  $\nu_t$  and  $\rho_t$  standing respectively for  $h(D(t))$ ,  $\nu_{C(t)}$  and  $\rho_{C(t)}$ .

When convenient and when there is no possible confusion, the index  $t \geq 0$  will even be omitted. It is an important goal of this paper to show that the above equation admits a solution in the whole temporal interval  $\mathbb{R}_+$  under some assumptions. Until it is proven, when we consider a time  $t \geq 0$ , it will be implicitly assumed that  $t$  is before the (random) time up to which a solution of (11) is constructed.

**Remark 3.2.** For the short time existence of (11), i.e. the existence of a stopping time  $\tau > 0$  such that equation (11) is defined for all  $t \in [0, \tau[$ , we refer to Theorem 40 in [2], where the authors use Doss-Sussman method and the theory of quasi-linear equations, as well as inverse function theorem. In the sequel, any such positive stopping time  $\tau$  will be called a **lifetime** for (11).

Concerning the regularity, for any  $0 < \alpha < 1$ , the solution of (11) is  $C^{\alpha/2, \infty}$  if the starting curve  $C(0, \cdot)$  is smooth,. In fact it is enough to consider that the starting curve  $C(0, \cdot)$  are  $C^{\alpha+n}$ , for  $n \geq 2$  and  $0 < \alpha < 1$ , to get the  $C^{\alpha/2, \alpha+n}$  regularity of the solution of (11) (cf. Chapter 8 of Lunardi [9] and Chapter 4 in [2]). So, to justify the existence of all the derivatives one may need, it is sufficient to take  $C_0$  regular enough, but we will not insist on the regularity of  $C_0$  in the rest of the paper.

To a solution  $(C(t))_{t \geq 0}$  of (11), as in the deterministic situation, associate

$$\forall t \geq 0, \forall u \in \mathbb{T}, \quad v(t, u) := |\partial_u C(t, u)|$$

and the arc-length parametrization  $\partial_s := \frac{1}{v} \partial_u$  (equivalently  $ds = v du$ ).

For any  $t \geq 0$  and  $u \in \mathbb{T}$ ,  $T_t(u)$  will stand for the unit tangent vector of the curve  $C(t)$  at  $u$  (i.e. in  $\mathbb{R}^2$  we have  $\mathcal{R}(T) = \nu$ , where  $\mathcal{R}$  is the rotation of angle  $-\frac{\pi}{2}$ ).

The evolution of these objects is dictated by the following result, for  $C_0$  regular enough (say  $C^{4+\alpha}$  for the last equation).

**Lemma 3.3.** Let  $(C(t))_{t \geq 0}$  be a solution of (11). We have the following equations in the  $(t, s)$ -domain of validity:

$$\begin{cases} d_t v_t &= v_t \rho_t ((-\rho_t + 2h_t)dt + \sqrt{2}dB_t) \\ [d_t, \partial_s] &= \rho_t ((3\rho_t - 2h_t)dt - \sqrt{2}dB_t) \partial_s - \sqrt{2}\rho_t dB_t \partial_s d_t \\ d_t T_t &= -\frac{1}{v_t} (\partial_u \rho_t) \nu_t dt \\ d_t \rho_t(s) &= (\partial_s^2 \rho_t) dt + \rho_t^2 ((3\rho_t - 2h_t)dt - \sqrt{2}dB_t) \end{cases} \quad (12)$$

*Proof.* Since  $C(t, u)$  satisfies

$$d_t C(t, u) = (-\rho_t(C(t, u)) + 2h(D(t))) \nu_{C(t, u)} dt + \sqrt{2} \nu_{C(t, u)} dB_t$$

we have, after differentiation, cf. remark 3.2, and shortening the notation:

$$d_t \partial_u C = (-\partial_u \rho_t) \nu_t dt + \left( (-\rho_t + 2h_t)dt + \sqrt{2}dB_t \right) \partial_u \nu_t,$$

Also we have

$$\partial_u \nu_t = v \partial_s \nu_t = v \rho_t T_t,$$

so that

$$d_t \partial_u C = (-\partial_u \rho_t) \nu_t dt + v_t \rho_t \left( (-\rho_t + 2h_t)dt + \sqrt{2}dB_t \right) T_t.$$

Hence we have the following equation:

$$\begin{aligned}
 d_t(v_t)^2 &= d_t|\partial_u C(t, u)|^2 \\
 &= 2\langle d_t\partial_u C(t, u), \partial_u C(t, u) \rangle + \langle d_t\partial_u C(t, u), d_t\partial_u C(t, u) \rangle \\
 &= 2v_t^2\rho_t \left( (-\rho_t + 2h_t)dt + \sqrt{2}dB_t \right) + 2v_t^2\rho_t^2 dt \\
 &= 2v_t^2\rho_t \left( (2h_t)dt + \sqrt{2}dB_t \right).
 \end{aligned}$$

Also

$$dv_t^2 = 2v_t dv_t + dv_t dv_t,$$

where the semi-martingale bracket notation  $\langle dv_t, dv_t \rangle$  has been simplified into  $dv_t dv_t$ .

Hence

$$2v_t dv_t + dv_t dv_t = 2v_t^2\rho_t \left( 2h_t dt + \sqrt{2}dB_t \right),$$

so the Doob-Meyer decomposition of  $v_t$  is  $dv_t = \sqrt{2}v_t\rho_t dB_t + dA_t$  where  $A_t$  is a process with finite variation. After identification we find:

$$dA_t = v_t\rho_t(-\rho_t + 2h)dt$$

and so

$$d_tv_t = v_t\rho_t \left( (-\rho_t + 2h)dt + \sqrt{2}dB_t \right).$$

For the second equation let us observe that for a vector-valued process  $X_t$ :

$$\begin{aligned}
 d_t\partial_s X_t &= d_t \left( \frac{1}{v_t} \partial_u X_t \right) \\
 &= d_t \left( \frac{1}{v_t} \right) \partial_u X_t + \frac{1}{v_t} \partial_u d_t X_t + d_t \left( \frac{1}{v_t} \right) d_t(\partial_u X_t) \\
 &= \frac{\rho_t}{v_t} \left( (3\rho_t - 2h)dt - \sqrt{2}dB_t \right) \partial_u X_t + \partial_s d_t X_t - \sqrt{2} \frac{\rho_t}{v_t} dB_t \partial_u d_t X_t \\
 &= \rho_t \left( (3\rho_t - 2h)dt - \sqrt{2}dB_t \right) \partial_s X_t + \partial_s d_t X_t - \sqrt{2}\rho_t dB_t \partial_s d_t X_t.
 \end{aligned}$$

In other words, we have:

$$[d_t, \partial_s] = \rho_t \left( (3\rho_t - 2h)dt - \sqrt{2}dB_t \right) \partial_s - \sqrt{2}\rho_t dB_t \partial_s d_t$$

For the third point, let us compute:

$$\begin{aligned}
 d_t T_t &= d_t \left( \frac{1}{v_t} \partial_u C_t \right) = d_t \left( \frac{1}{v_t} \right) \partial_u C_t + \frac{1}{v_t} d_t \partial_u C_t + d_t \left( \frac{1}{v_t} \right) d_t \partial_u C_t \\
 &= -\frac{\rho_t}{v_t} \left( (-3\rho_t + 2h)dt + \sqrt{2}dB_t \right) v_t T_t \\
 &\quad + \frac{1}{v_t} \left( (-\partial_u \rho_t) v_t dt + v_t \rho_t \left( (-\rho_t + 2h_t)dt + \sqrt{2}dB_t \right) T_t \right) \\
 &\quad - 2\rho_t^2 T_t dt \\
 &= -\frac{1}{v_t} (\partial_u \rho_t) v_t dt
 \end{aligned}$$

that is  $d_t \partial_s C(t, s) = -(\partial_s \rho_t) \nu_t dt$ . This equation is equivalent to

$$d_t \nu_t = d_t \mathcal{R}(T_t) = (\partial_s \rho_t) T_t dt.$$

So the processes  $T_t$  and  $\nu_t$  have finite variation.

For the last point, the mean curvature  $\rho_t$  satisfies:

$$\begin{aligned} d_t \rho_t &= -d_t \langle \partial_s T_t, \nu_t \rangle \\ &= -\langle d_t \partial_s T_t, \nu_t \rangle - \langle \partial_s T_t, d_t \nu_t \rangle \\ &= -\langle d_t \partial_s T_t, \nu_t \rangle. \end{aligned}$$

In the last line we used that  $\partial_s T_t$  is in the normal direction. Let us compute, using the commutation formula in the first term in the above bracket and the fact that  $\partial_s C_t$  has finite variation:

$$\begin{aligned} d_t \partial_s T_t &= d_t (\partial_s \partial_s C_t) \\ &= \rho_t \left( (3\rho_t - 2h) dt - \sqrt{2} dB_t \right) \partial_s \partial_s C_t + \partial_s d_t \partial_s C_t - \sqrt{2} \rho_t dB_t \partial_s d_t \partial_s C_t \\ &= \rho_t \left( (3\rho_t - 2h) dt - \sqrt{2} dB_t \right) \partial_s \partial_s C_t + \partial_s d_t \partial_s C_t \\ &= \rho_t \left( (3\rho_t - 2h) dt - \sqrt{2} dB_t \right) \partial_s \partial_s C_t + \partial_s (-\partial_s \rho_t) \nu_t dt \\ &= -\rho_t^2 \left( (3\rho_t - 2h) dt - \sqrt{2} dB_t \right) \nu_t + (-\partial_s^2 \rho_t) \nu_t - \rho_t (\partial_s \rho_t) T_t dt. \end{aligned}$$

Hence

$$\begin{aligned} d_t \rho_t &= -\langle -\rho_t^2 \left( (3\rho_t - 2h) dt - \sqrt{2} dB_t \right) \nu_t + (-\partial_s^2 \rho_t) \nu_t - \rho_t (\partial_s \rho_t) T_t dt, \nu_t \rangle \\ &= \partial_s^2 \rho_t dt + \rho_t^2 \left( (3\rho_t - 2h) dt - \sqrt{2} dB_t \right). \end{aligned} \tag{13}$$

□

**Remark 3.4.** We want to stress that (13) is a SPDE with mobile barrier (recall the parameter  $s$  lives in a time dependent interval  $[0, \sigma_t]$ ).

**Remark 3.5.** Note also that after integration, we recover the equation satisfied by  $\sigma_t$  and  $\lambda_t$  obtained by  $\mathcal{L}$  diffusion technique (resp. Proposition 57 and equation (106) in [2]) in the case of a simple curve:

$$\begin{aligned} d_t \sigma_t &= d_t \int_0^{2\pi} v_t(u) du \\ &= \int_0^{2\pi} v_t \rho_t \left( (-\rho_t + 2h) dt + \sqrt{2} dB_t \right) du \\ &= \left( -\int \rho_t^2 ds + 2h_t \int \rho_t ds \right) dt + \left( \int \rho_t ds \right) \sqrt{2} dB_t \\ &= -\left( \int \rho_t^2 ds \right) dt + 4\pi h_t dt + 2\sqrt{2}\pi dB_t, \end{aligned}$$

Using, on one hand Stokes theorem, namely  $\int_D \operatorname{div}(b) d\lambda = \int_{\partial D} \langle \nu, b \rangle ds$ , here in dimension 2 and with  $b$  the identity vector field, and on the other hand the computation done in the proof of Lemma 3.3, we get:

$$\begin{aligned}
 d_t \lambda_t &= d_t \frac{1}{2} \int_0^{2\pi} \langle C(t, u), \mathcal{R}(\partial_u C(t, u)) \rangle du \\
 &= \frac{1}{2} \left( \int_0^{2\pi} v_t \left( (-\rho_t + 2h) dt + \sqrt{2} dB_t \right) du + \int_0^{2\pi} \langle C(t, u), (\partial_u \rho_t(u)) T dt \rangle du \right. \\
 &\quad \left. + \int_0^{2\pi} v_t \rho_t \left( (-\rho_t + 2h) dt + \sqrt{2} dB_t \right) \langle C(t, u), \nu \rangle du + 2 \int_0^{2\pi} v_t \rho_t dt du \right).
 \end{aligned}$$

For the second term in the right hand side we integrate by part and we use  $\partial_u T = -v_t \rho_t \nu_t$  to get:

$$\int_0^{2\pi} \langle C(t, u), (\partial_u \rho_t(u)) T dt \rangle du = - \int_0^{2\pi} v_t \rho_t dt du - \rho_t^2 v_t \langle C(t, u), \nu_t dt \rangle du.$$

And then

$$\begin{aligned}
 d_t \lambda_t &= \frac{1}{2} \left( \int_0^{2\pi} v_t (2h dt + \sqrt{2} dB_t) du \right. \\
 &\quad \left. + \int_0^{2\pi} v_t \rho_t (2h dt + \sqrt{2} dB_t) \langle C(t, u), \nu \rangle du \right).
 \end{aligned}$$

In the last term of the right hand side we can integrate by part again to get

$$\begin{aligned}
 \int_0^{2\pi} v_t \rho_t \langle C(t, u), \nu \rangle du &= - \int_0^{2\pi} \langle C(t, u), \partial_u T \rangle du \\
 &= \int_0^{2\pi} (-\partial_u \langle C(t, u), T \rangle + v_t) du = \int_0^{2\pi} v_t du.
 \end{aligned}$$

Hence:

$$\begin{aligned}
 d_t \lambda_t &= \int_0^{2\pi} v_t (2h dt + \sqrt{2} dB_t) du \\
 &= \frac{2\sigma_t^2}{\lambda_t} dt + \sqrt{2} \sigma_t dB_t.
 \end{aligned}$$

Let us recall that for a simple strictly convex closed curve in  $\mathbb{R}^2$  it is possible to parametrize the curve using the angle  $\theta$  between the tangent line and the  $x$  axis, i.e.  $T = (\cos(\theta), \sin(\theta))$ . Usually, the angle  $\theta$  depends on  $u$  and  $t$ . Following [5] and [6], after adding to the stochastic flow (11) a tangential perturbation with finite variation, we do not perturb the shape of the curve, and as detailed below, it is possible to find a tangential intensity so that the parameter  $\theta$  does not depend on the time. This change of coordinate will make the equation easier to understand (since operators  $\partial_\theta$  and  $\partial_t$  will commute, contrary to  $\partial_s$  and  $\partial_t$  as shown in Lemma 3.3).

Let us quickly show this fact. Consider the following tangential perturbation of equation (11):

$$\begin{cases} d_t \tilde{C}(t, u) &= \left( -\tilde{\rho}_t(u) + 2h(\tilde{D}_t) \right) dt + \sqrt{2} dB_t \tilde{\nu}_t(u) + (\alpha_t(u) dt) \tilde{T}_t(u) \\ \tilde{C}(0, u) &= C_0(u) \end{cases} \quad (14)$$

Since only the parametrisation changes, the lifetime of (14) is the same as the one of (11), so for simplicity we will drop the tilde in the above equation, and sometimes the parameter  $u$ . With similar computation as we have done before in the proof of Lemma 3.3 we get:

**Lemma 3.6.** *Letting  $C_t(u)$  be a solution of (14), we have:*

$$\begin{aligned} d_t \partial_u C_t &= ((-\partial_u \rho_t - \rho_t v_t \alpha_t) dt) \nu_t \\ (1) \quad &+ \left( v_t \rho_t [(-\rho_t + 2h) dt + \sqrt{2} dB_t] + \partial_u \alpha_t dt \right) T_t. \\ (2) \quad dv_t &= v_t \rho_t [(-\rho_t + 2h) dt + \sqrt{2} dB_t] + \partial_u \alpha_t dt. \\ (3) \quad d_t T_t &= -\frac{1}{v_t} (\partial_u \rho_t + \rho_t v_t \alpha_t) dt \nu_t. \end{aligned}$$

*Proof.* For the first point, we differentiate term by term and we use that:

$$\partial_u \nu_t = v_t \rho_t T_t$$

and so

$$\partial_u T_t = -v_t \rho_t \nu_t.$$

For the second point:

$$\begin{aligned} d_t (v_t)^2 &= d_t |\partial_u C(t, u)|^2 = 2 \langle d_t \partial_u C(t, u), \partial_u C(t, u) \rangle + \langle d_t \partial_u C(t, u), d_t \partial_u C(t, u) \rangle \\ &= 2v_t \left( v_t \rho_t \left( (-\rho_t + 2h_t) dt + \sqrt{2} dB_t \right) + \partial_u \alpha_t dt \right) + 2v_t^2 \rho_t^2 dt. \end{aligned}$$

Then we write  $dv_t^2 = 2v_t dv_t + dv_t dv_t$ , and we identify the martingale part and the finite variation part of  $v_t$  to get the conclusion.

For the last point we compute:

$$\begin{aligned} d_t T_t &= d_t \left( \frac{1}{v_t} \partial_u C_t \right) = d_t \left( \frac{1}{v_t} \right) \partial_u C_t + \frac{1}{v_t} d_t \partial_u C_t + d_t \left( \frac{1}{v_t} \right) d_t \partial_u C_t \\ &= \left( -\frac{\rho_t}{v_t} \left( (-3\rho_t + 2h) dt + \sqrt{2} dB_t \right) - \frac{\partial_u \alpha_t}{v_t^2} dt \right) v_t T_t \\ &+ \frac{1}{v_t} \left( (-\partial_u \rho_t - \rho_t v_t \alpha_t) \nu_t dt + \left( v_t \rho_t [(-\rho_t + 2h_t) dt + \sqrt{2} dB_t] + \partial_u \alpha_t dt \right) T_t \right) \\ &- 2\rho_t^2 T_t dt \\ &= -\frac{1}{v_t} (\partial_u \rho_t + \rho_t v_t \alpha_t) dt \nu_t. \end{aligned}$$

□

In the above lemma, if the curvature is positive and if we take

$$\alpha_t = \frac{-\partial_u \rho_t}{v_t \rho_t}$$

we get that  $T_t$  and  $\nu_t$  become constant in time hence the angle  $\theta$  becomes constant in time, as desired.

The following lemma will describe the evolution of the curvature in this system of coordinates.



**Lemma 3.7.** *If  $C_0$  is  $C^{\alpha+4}$  and strictly convex, the solution to*

$$d_t \rho_t(\theta) = \rho_t^2(\theta) (\partial_\theta^2 \rho_t(\theta)) dt + \rho_t^2(\theta) \left( (3\rho_t(\theta) - 2h) dt - \sqrt{2} dB_t \right),$$

*is well-defined for all  $0 \leq t < \tau_0 \wedge \tau$  where,  $\tau_0 = \inf\{t \geq 0, \text{s.t. } \exists u \in [0, 1], \rho_t(u) = 0\}$  and  $\tau$  is the lifetime of (14). Also by definition of  $\tau$  all the quantities  $h, \rho, \partial_\theta \rho, \partial_\theta^2 \rho$  are bounded until  $\tau$ .*

*Proof.* By the above choice of  $\alpha$  we have:

$$\begin{aligned} 0 &= \partial_u d_t T_t = d_t(\partial_u T_t) \\ &= d_t(-\rho_t \nu_t \nu_t). \end{aligned}$$

Since  $\nu_t$  is constant in time we get:

$$d_t(v_t \rho_t) = 0,$$

and so

$$0 = d_t(v_t \rho_t) = d_t(v_t) \rho_t + v_t d_t \rho_t + dv_t d \rho_t.$$

We get that  $\rho_t$  satisfies the following stochastic differential equation:

$$\begin{aligned} v_t d \rho_t &= -\rho_t d_t(v_t) - dv_t d \rho_t \\ &= -\rho_t \left( v_t \rho_t [(-\rho_t + 2h) dt + \sqrt{2} dB_t] + \partial_u \alpha_t dt \right) - dv_t d \rho_t. \end{aligned}$$

After identification, the martingale part of  $d \rho_t$  is  $-\sqrt{2} \rho_t^2 dB_t$ , hence by this choice of  $\alpha$

$$\begin{aligned} d \rho_t &= -\rho_t^2 \left( (-\rho_t + 2h) dt + \sqrt{2} dB_t \right) - \frac{\rho_t}{v_t} \partial_u \alpha_t dt + 2 \rho_t^3 dt \\ &= -\rho_t^2 \left( (-3\rho_t + 2h) dt + \sqrt{2} dB_t \right) - \frac{\rho_t}{v_t} \partial_u \alpha_t dt \\ &= -\frac{\rho_t}{v_t} \partial_u \left( \frac{-\partial_u \rho_t}{v_t \rho_t} \right) dt - \rho_t^2 \left( (-3\rho_t + 2h) dt + \sqrt{2} dB_t \right). \end{aligned}$$

Recall that  $T = (\cos(\theta), \sin(\theta))$  and so  $\partial_\theta T = -\nu$ . So by the chain rule we have:

$$-\nu = \frac{\partial T}{\partial \theta} = \frac{\partial u}{\partial \theta} \frac{\partial T}{\partial u} = \frac{\partial u}{\partial \theta} (-v \rho) \nu.$$

Hence

$$\frac{\partial u}{\partial \theta} = \frac{1}{v \rho} \quad \text{and} \quad \partial_\theta = \frac{1}{v \rho} \partial_u. \quad (15)$$

The previous evolution equation of  $\rho_t$  becomes

$$\begin{aligned} d \rho_t &= \frac{\rho_t}{v_t} \partial_u \partial_\theta \rho_t dt - \rho_t^2 \left( (-3\rho_t + 2h) dt + \sqrt{2} dB_t \right) \\ &= \rho_t^2 \partial_\theta^2 \rho_t dt + \rho_t^2 \left( (3\rho_t - 2h) dt - \sqrt{2} dB_t \right). \end{aligned}$$

□

**Theorem 3.8.** *Let  $(C_0(\theta))_{\theta \in \mathbb{R}/(2\pi\mathbb{Z})}$  be a strictly convex simple closed curve,*

parametrized by the angle  $\theta$  (in particular  $\rho_0 > 0$ ).

Let  $\rho_t(\theta)$  be a solution of the following elliptic partial stochastic differential equation:

$$\begin{cases} d_t \rho_t(\theta) &= \rho_t^2(\theta) (\partial_\theta^2 \rho_t) dt + \rho_t^2(\theta) \left( (3\rho_t(\theta) - 2h) dt - \sqrt{2} dB_t \right) \\ \rho_0(\theta) &= \rho_0(\theta). \end{cases} \quad (16)$$

with lifetime  $\tau_2$ , namely the solution has to be regular up to order 2 at least for all times smaller than  $\tau_2$ .

Then  $\tau_0 \wedge \tau \leq \tau_2$ , and for all  $t < \tau_2$ ,  $\rho_t(\theta) > 0$  for all  $\theta \in \mathbb{R}/(2\pi\mathbb{Z})$  and (16) provides the solution of (11) by :

$$C(t, \theta) := \tilde{C}(t, \theta) + \int_0^t (-\partial_\theta \rho_u(0), \rho_u(0) - 2h_u) du - (0, \sqrt{2} B_t)$$

where

$$\tilde{C}(t, \theta) = \left( \int_0^\theta \frac{\cos(\theta_1)}{\rho_t(\theta_1)} d\theta_1, \int_0^\theta \frac{\sin(\theta_1)}{\rho_t(\theta_1)} d\theta_1 \right).$$

Thus by uniqueness of the solution of (11), (16) admits a unique regular solution too, and  $\tau_2 \leq \tau$ .

*Proof.* By lemma 3.7, (16) admit a solution and  $\tau_0 \wedge \tau \leq \tau_2$ . Note that the quantity  $h_t$  could be expressed in terms of  $\sigma_t$  and  $\lambda_t$  and these quantities also depend on the integral of  $\rho$  as seen in Remark 3.10 below and so  $h$  is bounded until  $\tau_2$ .

From (16), we get for all  $t < \tau_2$ ,

$$\rho_t(\theta) = \rho_0(\theta) \exp \int_0^t -\sqrt{2} \rho_s(\theta) dB_s + (\rho_s(\theta) \partial_\theta^2 \rho_s(\theta) + 2\rho_s(\theta)(\rho_s(\theta) - h_s)) ds$$

which is positive.

Recall Lemma 4.1.1 in [6], or see the beginning of Section 5, that says a  $2\pi$  periodic positive function  $\rho$  represents the curvature of a simple closed strictly convex plane curve if and only if

$$\int_0^{2\pi} \frac{\cos(\theta)}{\rho(\theta)} d\theta = \int_0^{2\pi} \frac{\sin(\theta)}{\rho(\theta)} d\theta = 0. \quad (17)$$

Here this equation is satisfied by  $\rho_0(\theta)$ . So we have to check that this relation is conserved over time for  $\rho_t(\theta)$  solution of (16). We will only verify this fact for the first coordinate, the computation will be the same for the second one. Using Itô calculus we get for  $0 \leq t < \tau_2$ :

$$\begin{aligned} d_t \frac{1}{\rho_t} &= -\frac{1}{\rho_t^2} d\rho_t + \frac{1}{\rho_t^3} d\rho_t d\rho_t \\ &= -(\partial_\theta^2 \rho_t(\theta)) dt - \left( (3\rho_t(\theta) - 2h) dt - \sqrt{2} dB_t \right) + 2\rho_t dt \\ &= -(\partial_\theta^2 \rho_t(\theta)) dt - \left( (\rho_t(\theta) - 2h) dt - \sqrt{2} dB_t \right). \end{aligned} \quad (18)$$

And so after integration by part we get for  $0 \leq t < \tau_2$ :

$$\begin{aligned}
d_t \int_0^{2\pi} \frac{\cos(\theta)}{\rho_t(\theta)} d\theta &= \int_0^{2\pi} d_t \frac{\cos(\theta)}{\rho_t(\theta)} d\theta \\
&= \int_0^{2\pi} \cos(\theta) \left( -(\partial_\theta^2 \rho_t(\theta)) dt - \left( (\rho_t(\theta) - 2h) dt - \sqrt{2} dB_t \right) \right) d\theta \\
&= - \left( \int_0^{2\pi} \cos(\theta) (\partial_\theta^2 \rho_t(\theta) + \rho_t(\theta)) d\theta \right) dt \\
&= 0.
\end{aligned}$$

We get that, for all  $0 \leq t < \tau_2$ ,  $\rho_t$  is the curvature of a simple closed strictly convex plane curve. Let us write the curve as:

$$\tilde{C}(t, \theta) = \left( \int_0^\theta \frac{\cos(\theta_1)}{\rho_t(\theta_1)} d\theta_1, \int_0^\theta \frac{\sin(\theta_1)}{\rho_t(\theta_1)} d\theta_1 \right).$$

We only have to check that  $(C(t, \theta))_\theta$  solves Equation (11) up to some tangential component.

We have:

$$\begin{aligned}
d_t C(t, \theta) &= d_t \left( \int_0^\theta \frac{\cos(\theta_1)}{\rho_t(\theta_1)} d\theta_1, \int_0^\theta \frac{\sin(\theta_1)}{\rho_t(\theta_1)} d\theta_1 \right) \\
&\quad + (-\partial_\theta \rho_t(0) dt, (\rho_t(0) - 2h_t) dt - \sqrt{2} dB_t) \\
&= \left( \int_0^\theta \cos(\theta_1) \left( -(\partial_{\theta_1}^2 \rho_t(\theta_1)) dt - \left( (\rho_t(\theta_1) - 2h) dt - \sqrt{2} dB_t \right) \right) d\theta_1, \right. \\
&\quad \left. \int_0^\theta \sin(\theta_1) \left( -(\partial_{\theta_1}^2 \rho_t(\theta_1)) dt - \left( (\rho_t(\theta_1) - 2h) dt - \sqrt{2} dB_t \right) \right) d\theta_1 \right) \\
&\quad + (-\partial_\theta \rho_t(0) dt, (\rho_t(0) - 2h_t) dt - \sqrt{2} dB_t).
\end{aligned}$$

After two integrations by parts, we have for the first term in the right hand side:

$$\begin{aligned}
&\int_0^\theta \cos(\theta) \left( -(\partial_{\theta_1}^2 \rho_t(\theta_1)) dt - \left( (\rho_t(\theta_1) - 2h) dt - \sqrt{2} dB_t \right) \right) d\theta_1 \\
&= - \left\{ [\cos(\theta_1) \partial_{\theta_1} \rho_t]_0^\theta dt + [\sin(\theta_1) \rho_t]_0^\theta dt + [\sin(\theta)] \left( -2h dt - \sqrt{2} dB_t \right) \right\} \\
&= -\cos(\theta) \partial_\theta \rho_t(\theta) dt + \partial_\theta \rho_t(0) dt - \sin(\theta) \left( (\rho_t - 2h) dt - \sqrt{2} dB_t \right).
\end{aligned}$$

For the second term, we have:

$$\begin{aligned}
& \int_0^\theta \sin(\theta) \left( -(\partial_{\theta_1}^2 \rho_t(\theta_1)) dt - ((\rho_t(\theta_1) - 2h) dt - \sqrt{2} dB_t) \right) d\theta_1 \\
&= - \left\{ [\sin(\theta_1) \partial_{\theta_1} \rho_t]_0^\theta dt - [\cos(\theta_1) \rho_t]_0^\theta dt - [\cos(\theta)]_0^\theta \left( -2h dt - \sqrt{2} dB_t \right) \right\} \\
&= - \sin(\theta) \partial_\theta \rho_t(\theta) dt + \cos(\theta) \left( (\rho_t - 2h) dt - \sqrt{2} dB_t \right) \\
&\quad - \left( (\rho_t(0) - 2h) dt - \sqrt{2} dB_t \right).
\end{aligned}$$

Hence:

$$d_t C(t, \theta) = ((-\rho_t + 2h) dt + \sqrt{2} dB_t) \nu - (\partial_\theta \rho_t dt) T.$$

This is (14) and so up a parametrization, this is a solution of (11). Since a solution of (16) produces a solution of (11), by uniqueness of solution of (11), we get the uniqueness of the solution of (16), and  $\tau_2 \leq \tau$ .  $\square$

We will show that Equation (11) preserves the positivity of the mean curvature.

**Lemma 3.9.** *Consider the solution of (11), if  $C_0$  is  $C^{\alpha+4}$  and  $\rho_0 > 0$  then  $\rho_t > 0$  for all  $t < \tau$ , where  $\tau$  is any lifetime of (11), moreover  $\tau = \tau_2$ .*

*Proof.* Suppose that  $\tau_0 < \tau$ , so  $h_t, \rho_t(\theta), \partial_\theta \rho_t(\theta), \partial_\theta^2 \rho_t(\theta)$  are bounded for all  $t \leq \tau_0$ , and

$$\rho_{\tau_0}(\theta) = \rho_0(\theta) \exp^{\int_0^{\tau_0} -\sqrt{2}\rho_s(\theta) dB_s + (\rho_s(\theta) \partial_\theta^2 \rho_s(\theta) + 2\rho_s(\theta)(\rho_s(\theta) - h_s)) ds},$$

and we get a contradiction. By Theorem 3.8, we get  $\tau = \tau_2$ .  $\square$

**Remark 3.10.** *Let us compute the equation satisfied by  $h$  when we know the equation of  $\rho$ . Resorting to (18) and recalling from (15) that here  $1/(v\rho) = \partial u / \partial \theta = 1$ , we get by Stokes Theorem:*

$$\begin{aligned}
d\sigma_t &= d \int_0^{2\pi} |\partial_\theta C(t, \theta)| d\theta \\
&= d \int_0^{2\pi} \frac{1}{\rho_t(\theta)} d\theta \\
&= \int_0^{2\pi} \left( -\partial_\theta^2 \rho_t(\theta) dt - ((\rho_t(\theta) - 2h) dt - \sqrt{2} dB_t) \right) d\theta \\
&= \left( - \int_0^{2\pi} \rho_t(\theta) d\theta \right) dt + 4\pi h_t dt + 2\sqrt{2}\pi dB_t \\
&= \left( - \int_0^{\sigma_t} \rho_t^2(s) ds \right) dt + 4\pi h_t dt + 2\sqrt{2}\pi dB_t.
\end{aligned}$$

Also by similar computation as above, we have:

$$\begin{aligned}
 d\lambda_t &= d\frac{1}{2} \int_0^{2\pi} \langle C(t, \theta), \nu_t(\theta) \rangle \frac{d\theta}{\rho_t(\theta)} \\
 &= \frac{1}{2} \left\{ \int_0^{2\pi} \left( \langle dC(t, \theta), \nu_t(\theta) \rangle \frac{1}{\rho_t(\theta)} + \langle C(t, \theta), \nu_t(\theta) \rangle d\left(\frac{1}{\rho_t(\theta)}\right) + \right. \right. \\
 &\quad \left. \left. + \langle dC(t, \theta), \nu_t(\theta) \rangle d\left(\frac{1}{\rho_t(\theta)}\right) \right) d\theta \right\} \\
 &= \frac{1}{2} \left\{ \int_0^{2\pi} \left( \left( (-\rho_t(\theta) + 2h)dt + \sqrt{2}dB_t \right) \frac{1}{\rho_t(\theta)} \right. \right. \\
 &\quad \left. \left. + \langle C(t, \theta), \nu_t(\theta) \rangle \left( -\partial_\theta^2 \rho_t(\theta)dt - ((\rho_t(\theta) - 2h)dt - \sqrt{2}dB_t) \right) + 2dt \right) d\theta \right\}
 \end{aligned}$$

After integrating by part two times and using  $\partial_\theta \nu = T$ , we get:

$$\begin{aligned}
 \int_0^{2\pi} -\langle C(t, \theta), \nu_t(\theta) \rangle \partial_\theta^2 \rho_t(\theta) d\theta &= \int_0^{2\pi} \partial_\theta (\langle C(t, \theta), \nu_t(\theta) \rangle) \partial_\theta \rho_t(\theta) d\theta \\
 &= \int_0^{2\pi} \partial_\theta \rho_t(\theta) \langle C(t, \theta), T_t(\theta) \rangle d\theta \\
 &= - \int_0^{2\pi} \rho_t(\theta) \left( \frac{1}{\rho_t(\theta)} - \langle C(t, \theta), \nu_t(\theta) \rangle \right) d\theta.
 \end{aligned}$$

Taking into account that  $\partial_\theta T = -\nu$ , we have

$$\begin{aligned}
 \int_0^{2\pi} \langle C(t, \theta), \nu_t(\theta) \rangle d\theta &= - \int_0^{2\pi} \langle C(t, \theta), \partial_\theta T_t(\theta) \rangle d\theta \\
 &= - \int_0^{2\pi} \partial_\theta (\langle C(t, \theta), T_t(\theta) \rangle) - \frac{1}{\rho_t(\theta)} d\theta = \int_0^{2\pi} \frac{1}{\rho_t(\theta)} d\theta.
 \end{aligned}$$

Putting the two computations above in the evolution equation of  $\lambda_t$  we get:

$$\begin{aligned}
 d\lambda_t &= \frac{1}{2} \left\{ \int_0^{2\pi} \left( \left( 2hdt + \sqrt{2}dB_t \right) \frac{1}{\rho_t(\theta)} \right. \right. \\
 &\quad \left. \left. - \langle C(t, \theta), \nu_t(\theta) \rangle \left( -2hdt - \sqrt{2}dB_t \right) \right) d\theta \right\} \\
 &= \int_0^{2\pi} \frac{1}{\rho_t(\theta)} d\theta \left( 2hdt + \sqrt{2}dB_t \right) = d\lambda_t = \frac{2\sigma_t^2}{\lambda_t} dt + \sqrt{2}\sigma_t dB_t.
 \end{aligned}$$

So we have to interpret (16) as a system where we have:

$$\begin{cases} d\sigma_t &= \left( -\int_0^{2\pi} \rho_t(\theta) d\theta \right) dt + 4\pi \frac{\sigma_t}{\lambda_t} dt + 2\sqrt{2\pi} dB_t \\ d\lambda_t &= \frac{2\sigma_t^2}{\lambda_t} dt + \sqrt{2}\sigma_t dB_t \\ h_t &= \frac{\sigma_t}{\lambda_t} \end{cases} \quad (19)$$

Using the above theorem and Lemma 3.7, we get the following corollary:

**Corollary 3.11.** *If the starting curve is simple closed and strictly convex, there is a one to one correspondence between the solutions of (11), (16) and (14).*

*Proof.* Use Theorem 3.8, Lemma 3.7 and 3.9. □

Consider the following stochastic mean curvature flow,

$$\begin{cases} d_t C(t, u) &= (-\rho_t(C(t, u))) \nu_{C(t, u)} dt + \sqrt{2} \nu_{C(t, u)} dB_t \\ C(0, u) &= C_0(u) \end{cases} \quad (20)$$

**Corollary 3.12.** *Consider the solution of (20), if  $C_0$  is  $C^{\alpha+4}$  and  $\rho_0 > 0$ , then  $\rho_t > 0$  for all  $t < \tau$ , where  $\tau$  is any lifetime of (20).*

*Proof.* With similar computation as in the above lemma, and since  $\rho_0 > 0$ , we have:

$$\begin{cases} d_t \rho_t(\theta) &= \rho_t^2(\theta) (\partial_\theta^2 \rho_t(\theta)) dt + \rho_t^2(\theta) ((3\rho_t(\theta)) dt - \sqrt{2} dB_t), \\ \rho_0 &= \rho_0 \end{cases} \quad (21)$$

and the proof is similar to the proof of 3.7, 3.8, 3.9. □

**Corollary 3.13.** *If the starting curve is simple closed and strictly convex, there is a one to one correspondence between the solutions of (20) and (21).*

*Proof.* The proof is similar to that of Theorem 3.8, essentially remove all the  $h_t$ . □

## 4. LONG TIME EXISTENCE

### 4.1. Evolution of geometric quantities along the stochastic flow (11).

**Proposition 4.1.** *Let  $C_0$  be a strictly convex curve. Let  $C_t$  be the solution of (11),  $\lambda_t$  the volume of  $D_t$  and  $\sigma_t$  the perimeter of  $C_t$ . Then we have the followings equations for  $t < \tau$  (where  $\tau$  is any lifetime of (11)):*

- i)  $d_t(\sigma_t^2 - 4\pi\lambda_t) \leq -2\pi \left( \frac{\sigma_t^2 - 4\pi\lambda_t}{\lambda_t} \right) dt,$
- ii)  $d \frac{1}{\rho_t(\theta)} = -\partial_\theta^2 \rho_t(\theta) dt - (\rho_t(\theta) - 2h) dt + \sqrt{2} dB_t,$
- iii)  $d \int_0^{2\pi} \frac{1}{\rho_t^2} d\theta = -2 \int_0^{2\pi} (\partial_\theta \log(\rho_t))^2 d\theta dt + 2d\lambda_t,$

iv)

$$\begin{aligned} d \int_0^{2\pi} \log(\rho_t(\theta)) d\theta &= - \int_0^{2\pi} (\partial_\theta \rho_t)^2 d\theta dt + 2 \int_0^{2\pi} \left( \rho_t(\theta) - \frac{h}{2} \right)^2 d\theta dt \\ &\quad - \pi h^2 dt - \sqrt{2} \int_0^{2\pi} \rho_t(\theta) d\theta dB_t. \end{aligned}$$

*Proof.* For equation i): using equation (19) and Itô formula we have

$$\begin{aligned} d(\sigma_t^2 - 4\pi\lambda_t) &= 2\sigma_t d\sigma_t + d\sigma_t d\sigma_t - 4\pi d\lambda_t \\ &= 2\sigma_t \left( - \int_0^{2\pi} \rho_t(\theta) d\theta dt + 4\pi \frac{\sigma_t}{\lambda_t} dt + 2\sqrt{2}\pi dB_t \right) + 8\pi^2 dt \\ &\quad - \frac{8\pi\sigma_t^2}{\lambda_t} dt - 4\pi\sqrt{2}\sigma_t dB_t \\ &= 2\sigma_t \left( - \int_0^{2\pi} \rho_t(\theta) d\theta \right) dt + 8\pi^2 dt \\ &\leq \left( -2\pi \frac{\sigma_t^2}{\lambda_t} + 8\pi^2 \right) dt \\ &= -2\pi \left( \frac{\sigma_t^2 - 4\pi\lambda_t}{\lambda_t} \right) dt \\ &\leq 0 \end{aligned}$$

where we use the preservation of the convexity along the flow (Lemma 3.9) and Gage inequality for convex curve [3]:

$$\pi h(D) = \pi \frac{\sigma(C)}{\lambda(D)} \leq \int_C \rho^2(s) ds = \int_0^{2\pi} \rho(\theta) d\theta. \quad (22)$$

Also in the last inequality we use the isoperimetric estimate. So the isoperimetric deficit  $\sigma_t^2 - 4\pi\lambda_t$  is non-increasing along the flow. One of the geometric meaning of the isoperimetric deficit is the following Bonnesen inequality [4]:

$$\pi^2 (r_{\text{out}} - r_{\text{int}})^2 \leq \sigma^2(\partial D) - 4\pi\lambda(D)$$

where  $r_{\text{int}}, r_{\text{out}}$  are respectively the inradius and the circumradius of  $D$ .

For equation ii): it is done in the proof of Theorem 3.8.

For equation iii): using Itô formula in the point (ii) we get

$$\begin{aligned} d_t \frac{1}{\rho_t^2} &= \frac{2}{\rho_t} \left( -(\partial_\theta^2 \rho_t(\theta)) dt - ((\rho_t(\theta) - 2h) dt + \sqrt{2} dB_t) \right) + 2dt \\ &= -\frac{2}{\rho_t} \partial_\theta^2 \rho_t(\theta) dt + \frac{4}{\rho_t} h dt + \frac{2\sqrt{2}}{\rho_t} dB_t. \end{aligned}$$

Integrating the above equality we get (since  $\int_0^{2\pi} \frac{1}{\rho} d\theta = \sigma_t$ )

$$\begin{aligned} d \int_0^{2\pi} \frac{1}{\rho_t^2} d\theta &= -2 \int_0^{2\pi} \left( \frac{(\partial_\theta \rho_t)}{\rho_t} \right)^2 d\theta dt + 4 \frac{\sigma_t^2}{\lambda_t} dt + 2\sqrt{2}\sigma_t dB_t \\ &= -2 \int_0^{2\pi} (\partial_\theta \log(\rho_t))^2 d\theta dt + 2d\lambda_t. \end{aligned}$$

For equation iv) we use (16) and Itô formula:

$$\begin{aligned} d \log(\rho_t(\theta)) &= \frac{1}{\rho_t(\theta)} d\rho_t(\theta) - \frac{1}{2\rho_t^2(\theta)} d\rho_t(\theta) d\rho_t(\theta) \\ &= \rho_t(\theta) (\partial_\theta^2 \rho_t) dt + \rho_t(\theta) \left( (3\rho_t(\theta) - 2h) dt - \sqrt{2} dB_t \right) - \rho_t^2(\theta) dt \\ &= \rho_t(\theta) (\partial_\theta^2 \rho_t) dt + 2\rho_t(\theta) (\rho_t(\theta) - h) dt - \sqrt{2} \rho_t(\theta) dB_t. \end{aligned}$$

Integrating the above equation, we get:

$$\begin{aligned} d \int_0^{2\pi} \log(\rho_t(\theta)) d\theta &= - \int_0^{2\pi} (\partial_\theta \rho_t)^2 d\theta dt + 2 \int_0^{2\pi} \rho_t(\theta) (\rho_t(\theta) - h) d\theta dt \\ &\quad - \sqrt{2} \int_0^{2\pi} \rho_t(\theta) d\theta dB_t. \\ &= - \int_0^{2\pi} (\partial_\theta \rho_t)^2 d\theta dt + 2 \int_0^{2\pi} \left( \rho_t(\theta) - \frac{h}{2} \right)^2 d\theta dt \\ &\quad - \pi h^2 dt - \sqrt{2} \int_0^{2\pi} \rho_t(\theta) d\theta dB_t. \end{aligned}$$

□

**Remark 4.2.** Note that  $\int_0^{2\pi} \frac{1}{\rho_t^2} d\theta - 2\lambda_t \geq \frac{1}{2\pi} \sigma_t^2 - 2\lambda_t \geq 0$  where we use isoperimetric estimate for the last point. Hence

$$0 \leq \int_0^{2\pi} \frac{1}{\rho_t^2} d\theta - 2\lambda_t = -2 \int_0^t \int_0^{2\pi} (\partial_\theta \log(\rho_s))^2 d\theta ds + a_0$$

where  $a_0 = \int_0^{2\pi} \frac{1}{\rho_0^2} d\theta - 2\lambda_0 \geq 0$ . So if moreover  $C_0$  is a curve in the set  $\mathcal{S}_n$  of  $n$ -symmetric convex curves with star shaped skeleton for some  $n \geq 2$  (see section 5 for the definition) using Proposition 5.5,  $C_t \in \mathcal{S}_n$  and  $\theta \mapsto \rho_t(\theta)$  is non-decreasing and the above equation gives:

$$0 < 2\lambda_t \leq \frac{1}{\rho_t(0)} \sigma_t \leq \frac{2\pi}{\rho_t^2(0)}$$

so  $0 < \rho_t(0) \leq \sqrt{\frac{\pi}{\lambda_t}}$  and  $0 < \rho_t(0) \leq \frac{h_t}{2}$ . On the other hand we have

$$0 < \int_0^{2\pi} \frac{1}{\rho_t^2} d\theta \leq a_0 + 2\lambda_t,$$



and if  $C_0 \in \mathcal{S}_n$  then

$$0 < \frac{2\pi}{\rho_t^2(\pi/2)} \leq \frac{\sigma_t}{\rho_t(\pi/2)} \leq a_0 + 2\lambda_t$$

so  $\sqrt{\frac{2\pi}{a_0+2\lambda_t}} \leq \rho_t(\pi/2)$  and note also by the Gage inequality (22) we have  $\frac{h_t}{2} \leq \rho_t(\pi/2)$ .

**Lemma 4.3.**  $h_t$  is a positive super martingale, and so it is almost surely bounded for all  $t < \tau$ .

*Proof.* Using equation (19) and Itô formula we have

$$\begin{aligned} dh_t &= d\left(\frac{\sigma_t}{\lambda_t}\right) \\ &= \frac{1}{\lambda_t} d\sigma_t + \sigma_t d\left(\frac{1}{\lambda_t}\right) + d\sigma_t d\left(\frac{1}{\lambda_t}\right) \\ &= \frac{1}{\lambda_t} \left( \left( -\int_0^{2\pi} \rho_t(\theta) d\theta \right) dt + 4\pi \frac{\sigma_t}{\lambda_t} dt + 2\sqrt{2}\pi dB_t \right) - \frac{\sqrt{2}\sigma_t^2}{\lambda_t^2} dB_t - \frac{4\pi\sigma_t}{\lambda_t^2} dt \\ &= -\frac{1}{\lambda_t} \left( \int_0^{2\pi} \rho_t(\theta) d\theta \right) dt + \sqrt{2} \left( \frac{2\pi\lambda_t - \sigma_t^2}{\lambda_t^2} \right) dB_t \\ &\leq -\frac{\pi h_t}{\lambda_t} dt + \sqrt{2} \left( \frac{2\pi\lambda_t - \sigma_t^2}{\lambda_t^2} \right) dB_t \end{aligned}$$

□

In the sequel we will encounter random constants, they will be denoted under the form  $c(\omega)$ , where  $\omega$  stands for the randomness associated to underlying Brownian motions. This is a generic notation and the exact value of  $c(\omega)$  may change from line to line.

**Proposition 4.4.** Let  $C_0$  be a strictly convex curve. Let  $C_t$  be the solution of (11), and  $\tau$  be any lifetime of (11). Then there exists a positive random variable  $c(\omega) < \infty$  such that for all  $t < \tau(\omega)$ ,  $h_t(\omega) \leq c(\omega)$  and

$$\frac{1}{\frac{1}{\inf \rho_0} + \sqrt{2} \sup_{[0,t]} B_s + 2c(\omega)t} \leq \inf_{\theta} \rho_t(\theta).$$

*Proof.* Let  $J_t = \frac{1}{\rho_t(\theta)} - \sqrt{2}B_t - 2 \int_0^t h(s) ds - \frac{1}{\inf \rho_0}$ . By Lemma 3.9 this quantity is well defined, and by Proposition 4.1 we have

$$\begin{aligned} dJ_t(\theta) &= -\partial_{\theta}^2 \rho_t(\theta) dt - \rho_t(\theta) dt \\ &= \left( \rho_t^2(\theta) \partial_{\theta}^2 \left( \frac{1}{\rho_t(\theta)} \right) - 2 \frac{(\partial_{\theta} \rho_t(\theta))^2}{\rho_t(\theta)} - \rho_t(\theta) \right) dt \\ &\leq \left( \frac{1}{J_t(\theta) + \sqrt{2}B_t + 2 \int_0^t h(s) ds + \frac{1}{\inf \rho_0}} \right)^2 \partial_{\theta}^2 J_t dt. \end{aligned}$$

Using the maximum principle, we will show that  $J_t \leq 0$  for all  $t \in [0, \tau[$ . Suppose that there exists  $t_0 \in [0, \tau[$  and  $\theta_0$  such that  $J_{t_0}(\theta_0) = \alpha > 0$ . Let  $W_t := e^{-t} J_t$ , then  $W_{t_0}(\theta_0) = e^{-t_0} \alpha > 0$  and  $\sup_{\theta} W_{t_0} \geq e^{-t_0} \alpha > 0$ . Consider the time  $t_* = \inf\{t \in [0, t_0], s.t. \sup_{\theta} W_t = W_{t_0}(\theta_0)\}$ , and let  $\theta_*$  such that  $W_{t_*}(\theta_*) = \sup_{\theta} W_{t_*}$ . We have  $t_* > 0$  and

$$\partial_t W_t \leq \left( \frac{1}{e^t W_t(\theta) + \sqrt{2} B_t + 2 \int_0^t h(s) ds + \frac{1}{\inf \rho_0}} \right)^2 \partial_{\theta}^2 W_t - W_t.$$

Note that since  $0 \leq \partial_t W_t(\theta_*)|_{t_*}, \partial_{\theta}^2 W_{t_*}(\theta)|_{\theta_*} \leq 0$  and  $W_{t_*}(\theta_*) = e^{-t_*} \alpha > 0$  we get a contradiction. Hence for all  $t \in [0, \tau[$  we have

$$\frac{1}{\rho_t(\theta)} \leq \frac{1}{\inf \rho_0} + \sqrt{2} B_t + 2 \int_0^t h(s) ds.$$

Since  $h_t$  is a positive super martingale by Lemma 4.3, it is almost surely bounded in  $[0, \tau[$ , so there exists a positive random variable  $c(\omega) < \infty$  such that  $h_t(\omega) \leq c(\omega)$  and

$$\frac{1}{\frac{1}{\inf \rho_0} + \sqrt{2} \sup_{[0,t]} B_s + 2c(\omega)t} \leq \inf_{\theta} \rho_t(\theta).$$

□

**4.2. When there is a sufficient number of symmetries.** The goal of this section is to find a necessary condition on the strictly convex domain to guarantee the existence of the solution of (11) for all times. We will see that the entropy will be a supermartingale if the initial domain has enough symmetries. This condition seems to be a technical hypothesis and we think that it is not necessary, and perhaps there exists an entropy which is more adapted to our situation. From Lemma 3.7, we deduce the evolution of the entropy (defined in (10), it also coincides with the relative entropy of the curvature density with respect to the arc length Lebesgue measure, up to normalizations in terms of the length of the curve):

$$\begin{aligned} d\text{Ent}_t &= d \int_0^{2\pi} \log(\rho_t(\theta)) d\theta \\ &= - \int_0^{2\pi} (\partial_{\theta} \rho_t)^2 d\theta dt + 2 \int_0^{2\pi} \rho_t(\theta) (\rho_t(\theta) - h) d\theta dt \\ &\quad - \sqrt{2} \int_0^{2\pi} \rho_t(\theta) d\theta dB_t. \end{aligned}$$

**Proposition 4.5.** *If the boundary of the domain is strictly convex (recall Definition 1.1) then we have the following estimate*

$$2\rho_{\inf} \leq h \leq 2\rho_{\sup}$$

*Proof.* Let  $p$  be the support function, namely  $p(s) = \langle x(s), \nu(s) \rangle$ . Green Theorem asserts  $\lambda(D) = \frac{1}{2} \int_{\gamma} p(s) ds$  and we have  $\sigma(\partial D) = \int_{\gamma} p(s) \rho(s) ds$ . Indeed, we compute that

$$\begin{aligned} \int_{\gamma} p(s) \rho(s) ds &= \int_{\gamma} \langle x(s), \rho(s) \nu(s) \rangle ds \\ &= - \int_{\gamma} \langle x(s), x(s)'' \rangle ds \end{aligned}$$

and it remains to integrate by part to recognize  $\sigma(\partial D)$ .

Remark also that we can suppose that the origin is contained in the domain (else translate and all the quantities are invariant under translation). By convexity of the domain we have that  $p(\theta) > 0$ . Recalling that  $d\theta = \rho ds$ , we have  $\sigma(\partial D) = \int_{\mathbb{T}} p(\theta) d\theta$ , so that

$$\lambda(D) = \frac{1}{2} \int_{\mathbb{T}} \frac{p(\theta)}{\rho(\theta)} d\theta \leq \frac{1}{2\rho_{\inf}} \sigma(\partial D).$$

Hence  $2\rho_{\inf} \leq h(D)$ . The other inequality is a direct consequence of Gage inequality.  $\square$

**Proposition 4.6.** *If  $f(0) = f(L) = 0$  then*

$$\int_0^L f^2 d\theta \leq \left(\frac{L}{\pi}\right)^2 \int_0^L f'^2 d\theta$$

*Proof.* This is the Wirtinger inequality which can be proved by Fourier series.  $\square$

**Definition 4.7.** *We will say that a domain  $D$  have  $n$  axes of symmetries, if up to a translation there exists a linear straight line  $\Delta$  such that  $D$  is symmetric with respect to  $\Delta, R_{\pi/n}(\Delta), \dots, R_{(n-1)\pi/n}(\Delta)$ , where  $R_{\theta}$  is a rotation of angle  $\theta$ .*

**Proposition 4.8.** *If the boundary of the initial domain is strictly convex (recall Definition 1.1), and the domain has  $n$  axes of symmetries, with  $n \geq 3$  then the entropy is a super-martingale.*

*Proof.* Using proposition 4.5 and the symmetries there exists  $\theta_k \in [\frac{k\pi}{n}, \frac{(k+1)\pi}{n}]$  for  $k \in \llbracket 0, 2n-1 \rrbracket$  such that  $\rho(\theta_k) = \frac{h}{2}$ . Note that we can further impose that  $|\theta_k - \theta_{k-1}| \leq \frac{2\pi}{n}$  for  $k \in \llbracket 0, 2n \rrbracket$  (with  $\theta_{2n} = \theta_0 + 2\pi$ ).

So using Proposition 4.6, we get

$$\int_{\theta_k}^{\theta_{k+1}} \left(\rho(\theta) - \frac{h}{2}\right)^2 d\theta \leq \frac{4}{n^2} \int_{\theta_k}^{\theta_{k+1}} \rho'(\theta)^2 d\theta.$$

Hence

$$- \int_{\mathbb{T}} \rho'(\theta)^2 d\theta \leq -\frac{n^2}{4} \int_{\mathbb{T}} \left(\rho(\theta) - \frac{h}{2}\right)^2 d\theta,$$

and, if  $n \geq 3$  we have

$$\begin{aligned} d\text{Ent}_t &\leq -\frac{n^2}{4} \int_{\mathbb{T}} \left( \rho(\theta) - \frac{h}{2} \right)^2 d\theta + 2 \int_0^{2\pi} \left( \rho_t(\theta) - \frac{h}{2} \right)^2 d\theta dt - \pi h^2 dt \\ &\quad - \sqrt{2} \int_0^{2\pi} \rho_t(\theta) d\theta dB_t. \\ &\leq -\pi h^2 dt - \sqrt{2} \int_0^{2\pi} \rho_t(\theta) d\theta dB_t \end{aligned}$$

□

**Remark 4.9.** *The Green-Osher's inequality, see Theorem 0.2 of [8], shows*

$$\text{Ent}_t = \int_{\mathbb{T}} \ln(\rho_t) d\theta \geq \pi \ln \left( \frac{\pi}{\lambda_t} \right).$$

Since  $\frac{1}{\lambda_t}$  is a positive martingale, the r.h.s. is a super-martingale (at least on its domain of definition). Of course, this is not sufficient to insure that  $(\text{Ent}_t)_t$  itself is a super-martingale.

In the sequel we will use comparison of processes up to a continuous martingale term: when  $(X_t)_{t \in [0, \tau]}$  and  $(Y_t)_{t \in [0, \tau]}$  are two predictable processes with respect to the same underlying filtration and are defined on the same time-interval  $[0, \tau)$  (where  $\tau$  is a stopping time), we write

$$\forall t \in [0, \tau) \quad X_t \stackrel{(m)}{\leq} Y_t$$

to mean there exists a continuous martingale  $(M_t)_{t \in [0, \tau)}$  such that

$$\forall t \in [0, \tau) \quad X_t \leq Y_t + M_t$$

In the next four results,  $\tau$  will stand the maximal time up to which the equation of Lemma 3.7 admits a solution.

**Proposition 4.10.** *If  $D_0$  is strictly convex then:*

$$d \int (\partial_\theta \rho_t)^2 d\theta \stackrel{(m)}{\leq} \left( \left( \frac{13}{3} \right)^2 + 16 \right) \int \rho^4 d\theta dt$$

*Proof.* From Lemma 3.7, we deduce that on  $[0, \tau)$ , via integrations by parts,

$$\begin{aligned} &d \int (\partial \rho)^2 \\ &= 2 \int \partial \rho d\partial \rho + \int d\partial \rho d\partial \rho \\ &= 2 \int \partial \rho \partial d\rho + \int \partial d\rho \partial d\rho \\ &= -2 \int \partial^2 \rho d\rho + 2 \int (\partial \rho^2)^2 dt \end{aligned}$$

$$\begin{aligned}
 &= 2 \int \rho^2 \partial^2 \rho [(2h - 3\rho - \partial^2 \rho) dt + \sqrt{2} dB_t] + 8 \left( \int (\rho^2 \partial \rho) \partial \rho \right) dt \\
 &= 2 \left( \int \rho^2 \partial^2 \rho (2h - 3\rho - \partial^2 \rho) \right) dt - \frac{8}{3} \left( \int \rho^3 \partial^2 \rho \right) dt \\
 &\quad + 2\sqrt{2} \left( \int \rho^2 \partial^2 \rho \right) dB_t \\
 &\stackrel{(m)}{=} 2 \left( \int \rho^2 \left[ -(\partial^2 \rho)^2 + \left( 2h - \frac{13}{3} \rho \right) \partial^2 \rho \right] \right) dt \\
 &= 2 \left( \int \rho^2 \left[ \left( h - \frac{13}{6} \rho \right)^2 - \left( \partial^2 \rho + \frac{13}{6} \rho - h \right)^2 \right] \right) dt \\
 &\leq 2 \left( \int \rho^2 \left( h - \frac{13}{6} \rho \right)^2 \right) dt \\
 &\leq 4 \left( \int \left( \frac{13}{6} \right)^2 \rho^4 + h^2 \rho^2 \right) dt
 \end{aligned}$$

where  $\partial$  stands for the differentiation with respect to the underlying parameter  $\theta$  (which commutes with respect to the “stochastic differentiation with respect to time”  $d$ ).

Taking into account Gage’s inequality, we get

$$\begin{aligned}
 h^2 \int \rho^2 &\leq \frac{1}{\pi^2} \left( \int \rho \right)^2 \int \rho^2 \\
 &\leq 4 \int \rho^4
 \end{aligned}$$

and finally the desired bound.  $\square$

This observation leads us to investigate the evolution of  $\int \rho^4$  itself:

**Proposition 4.11.** *If  $D_0$  is strictly convex and has  $n$  axes of symmetries, with  $n \geq 7$  then*

$$d \int (\rho_t)^4 d\theta \stackrel{(m)}{\leq} c(\omega) dt$$

where  $c(\omega)$  is a finite random constant (independent of time), as mentioned before Proposition 4.4.

*Proof.* We compute

$$\begin{aligned}
 d \int \rho^4 &= 4 \int \rho^3 d\rho + 6 \int \rho^2 d\rho d\rho \\
 &\stackrel{(m)}{=} 4 \left( \int 6\rho^6 + \rho^5 \partial^2 \rho - 2h\rho^5 \right) dt \\
 &= 4 \left( \int 6\rho^6 - 5\rho^4 (\partial \rho)^2 - 2h\rho^5 \right) dt
 \end{aligned}$$

$$= 4 \left( \int 6\rho^6 - \frac{5}{9} (\partial\rho^3)^2 - 2h\rho^5 \right) dt$$

To deal with the middle term, let us resort to Wirtinger inequality, assuming  $n \geq 7$  axes of symmetry for  $D_0$ . Since the evolution equation is invariant by these symmetries, for any time  $t \in [0, \tau)$ , we still have that  $D_t$  has  $n$  axes of symmetry. We deduce that

$$\begin{aligned} \int (\partial\rho^3)^2 &= \int (\partial(\rho^3 - \rho_{\text{inf}}^3))^2 \\ &\geq \frac{49}{4} \int (\rho^3 - \rho_{\text{inf}}^3)^2 \end{aligned}$$

so that, taking into account Proposition 4.5,

$$\begin{aligned} d \int \rho^4 &\stackrel{(m)}{\leq} 4 \left( \int -\frac{29}{36}\rho^6 + \frac{245}{18}\rho^3\rho_{\text{inf}}^3 - \frac{245}{36}\rho_{\text{inf}}^6 - 2\rho^5h \right) dt \\ &\leq 2 \left( \int -\frac{29}{18}\rho^6 + \frac{245}{9}\rho^3\rho_{\text{inf}}^3 - \frac{389}{18}\rho_{\text{inf}}^6 \right) dt \\ &\leq c(\omega) dt \end{aligned}$$

To get the desired result, recall that  $\rho_{\text{inf}} \leq h/2$  and that  $h$  is a positive supermartingale and is thus a.s. bounded on its domain of definition.  $\square$

**Proposition 4.12.** *If  $D_0$  is strictly convex and has  $n$  axes of symmetries with  $n \geq 7$ , then there exists a finite random variable  $c(\omega)$  such that on the event  $\tau < \infty$ :*

$$\forall t \in [0, \tau), \quad \int (\partial_\theta \rho_t)^2 d\theta \leq c(\omega) \quad (23)$$

*Proof.* Let us show that there exists a finite random variable  $c_1(\omega)$  such that on the event  $\tau < \infty$ :

$$\forall t \in [0, \tau), \quad \int (\rho_t)^4 d\theta \leq c_1(\omega) \quad (24)$$

According to the previous proposition, there exist a finite random constant  $c(\omega) \geq 0$  and a continuous martingale  $(M_t)_{t \in [0, \tau)}$  such that

$$\forall t \in [0, \tau), \quad \int (\rho_t)^4 d\theta \leq c(\omega)t + M_t$$

Up to enriching the underlying probability space, we can find a Brownian motion  $(W_t)_{t \geq 0}$  such that

$$\forall t \in [0, \tau), \quad \int (\rho_t)^4 d\theta \leq c(\omega)t + W_{\langle M \rangle_t} \quad (25)$$

Thus on  $\{\tau < +\infty\}$ , we will deduce (24) as soon as we show

$$\lim_{t \rightarrow \tau^-} \langle M \rangle_t < +\infty$$

Note that if we had

$$\lim_{t \rightarrow \tau^-} \langle M \rangle_t = +\infty$$

we would get from (25) that

$$\inf_{t \in [0, \tau)} \int (\rho_t)^4 d\theta = -\infty$$

which is a contradiction. Hence there exists  $c_1(\omega)$  such that (24) is satisfied on the event  $\tau < \infty$ . According to Proposition 4.10 there exist a finite constant  $c_2(\omega) \geq 0$  and a continuous martingale  $(\tilde{M}_t)_{t \in [0, \tau)}$  such that on  $\{\tau < +\infty\}$

$$\forall t \in [0, \tau), \quad \int (\partial_\theta \rho_t)^2 d\theta \leq c_2(\omega)t + \tilde{M}_t$$

We deduce (23) by the same argument used to get (24).  $\square$

**Proposition 4.13.** *If  $D_0$  is strictly convex and has  $n$  axes of symmetries, with  $n \geq 7$  then there exists a random variable  $c(\omega)$  such that on the event  $\tau < \infty$ :*

$$\rho_t \leq c(\omega) < \infty \quad \forall t \in [0, \tau[.$$

*Proof.* On the event  $\tau < \infty$ , according to Propositions 4.12 and 4.8, there exists a random constant  $c(\omega) < \infty$  such that for all  $t < \tau$  we have:

$$\text{Ent}_t \leq c(\omega)$$

$$\int (\partial_\theta \rho_t)^2 \leq c(\omega).$$

Let  $r_t := \sup\{\rho_s(\theta), (\theta, s) \in [0, 2\pi] \times [0, t]\}$  for  $t < \tau$ . Then there exists  $(\theta_1, t_1) \in [0, 2\pi] \times [0, t]$  such that  $\rho_{t_1}(\theta_1) = r_t$ . For all  $\theta_2 \in [0, 2\pi]$ , we have

$$\begin{aligned} |\rho_{t_1}(\theta_1) - \rho_{t_1}(\theta_2)| &= \left| \int_{\theta_1}^{\theta_2} \partial_\theta \rho_{t_1}(\theta) d\theta \right| \\ &\leq \sqrt{|\theta_1 - \theta_2|} \sqrt{c(\omega)}, \end{aligned}$$

so

$$r_t - \sqrt{|\theta_1 - \theta_2|} \sqrt{c(\omega)} \leq \rho_{t_1}(\theta_2).$$

Then using Proposition 4.4 we get

$$\begin{aligned} \text{Ent}_{t_1} &\geq \int_{|\theta - \theta_1| \leq \frac{r_t^2}{4c(\omega)} \wedge \frac{\pi}{2}} \log(\rho_{t_1}(\theta)) d\theta + \int_{|\theta - \theta_1| \geq \frac{r_t^2}{4c(\omega)} \wedge \frac{\pi}{2}} \log(\rho_{t_1}(\theta)) d\theta \\ &\geq 2 \log\left(\frac{r_t}{2}\right) \left(\frac{r_t^2}{4c(\omega)} \wedge \frac{\pi}{2}\right) \\ &\quad + \left(2\pi - 2\left(\frac{r_t^2}{4c(\omega)} \wedge \frac{\pi}{2}\right)\right) \log\left(\frac{1}{\frac{1}{\inf \rho_0} + \sqrt{2} \sup_{[0, t_1]} B_s + 2c(\omega)t_1}\right) \\ &\geq 2 \log\left(\frac{r_t}{2}\right) \left(\frac{r_t^2}{4c(\omega)} \wedge \frac{\pi}{2}\right) \\ &\quad + \left(2\pi - 2\left(\frac{r_t^2}{4c(\omega)} \wedge \frac{\pi}{2}\right)\right) \log\left(\frac{1}{\frac{1}{\inf \rho_0} + \sqrt{2} \sup_{[0, \tau]} B_s + 2c(\omega)\tau}\right). \end{aligned}$$

On the event  $\tau < \infty$  the last term of the above equation is almost surely bounded, since the entropy is bounded from above on  $[0, \tau[$ . We get that  $\rho_t$  has to be a.s. uniformly bounded on  $t \in [0, \tau)$ .  $\square$

We will need the following lemma which is a small refinement of Lemma 4.1.1 from [6].

**Lemma 4.14.** *Let a  $2\pi$  periodic positive function  $\rho \in C^\alpha(\mathbb{T})$ , with  $\alpha \in (0, 1)$ , be such that*

$$\int_0^{2\pi} \frac{\cos(\theta)}{\rho(\theta)} d\theta = \int_0^{2\pi} \frac{\sin(\theta)}{\rho(\theta)} d\theta = 0. \quad (26)$$

*Consider the curve  $X : \theta \mapsto (\int_0^\theta \frac{\cos(u)}{\rho(u)} du, \int_0^\theta \frac{\sin(u)}{\rho(u)} du)$ , as before parametrized by the angle  $\theta \in \mathbb{T}$  of its tangent with respect to the horizontal axis, and whose curvature function is  $\rho$ . When  $X$  is parametrized by its arc-length, it becomes  $C^{2+\alpha}$ .*

*Proof.* Note that under the parametrization of  $X$  by  $\theta$  the curve seems to be only of order  $C^{1+\alpha}$ . Let us check it is in fact  $C^{2+\alpha}$  under the arc-length parametrization. Denoting  $s$  the arc length parametrization of  $X$ , we have  $\partial_s = \rho \partial_\theta$  and  $s(\theta) = \int_0^\theta \frac{1}{\rho(u)} du$ ,  $\partial_s \theta(s) = \rho(\theta(s))$ ,  $T(s) = (\cos(\theta(s)), \sin(\theta(s)))$  (as it should be, by definition of the parametrization by  $\theta$ ). From  $\partial_s \theta(s) = \rho(\theta(s))$ , we see that  $s \mapsto \theta(s)$  is  $C^{1+\alpha}$ . Furthermore, in the parameter  $s$ , the curve  $\tilde{X}(s) := X(\theta(s))$  satisfies  $\partial_s \tilde{X} = (\cos(\theta(s)), \sin(\theta(s)))$ , so we get that  $\tilde{X}$  is  $C^{2+\alpha}$ .  $\square$

**Theorem 4.15.** *If  $D_0$  is strictly convex and has  $n$  axes of symmetries, and  $n \geq 7$  then a.s.  $\tau = \infty$ , where  $\tau$  is the maximal lifetime of (11).*

*Proof.* Suppose that  $\mathbb{P}(\tau < \infty) > 0$ . Let  $C_t(\theta)$  be the solution of (11) namely

$$\begin{cases} d_t C(t, \theta) &= ([-\rho_t(C(t, \theta)) + 2h_t] dt + \sqrt{2} dB_t) \nu_t(C(t, \theta)) \\ C(0, \theta) &= C_0(\theta) \end{cases}$$

On the event  $\{\tau < \infty\}$ , using Lemma 4.3 and 4.13 we have for all  $t < \tau$ ,  $h_t \leq c(\omega) < \infty$  and  $\rho_t(\theta) \leq c(\omega) < \infty$ . Since  $\|\nu_t(C(t, \theta))\| = 1$  we have for  $s, t < \tau$  such that  $|t - s|$  is small:

$$|C(s, \theta) - C(t, \theta)| \leq c_1(\omega) |t - s|^{\frac{1}{2} - \epsilon},$$

where the random variable  $c_1$  depends on  $c$ . Hence there exists  $C_\tau : \mathbb{T} \mapsto \mathbb{R}^2$  such that  $C_t$  converges uniformly to  $C_\tau$ . On the other hand, using Proposition 4.12 we get by Hölder inequality that for all  $t < \tau$

$$|\rho_t(\theta) - \rho_t(\beta)| \leq \left| \int_\beta^\theta \partial \rho_t(\gamma) d\gamma \right| \leq c(\omega) \sqrt{|\theta - \beta|}.$$

Hence  $\rho$  is equi-continuous. So using again Proposition 4.13 and Ascoli Theorem we get that there exists a sequence  $(t_n)_n$  converging to  $\tau$  and a  $C^{\frac{1}{2}}$  function  $\rho_\tau$  such that  $\rho_{t_n}$  converges uniformly to  $\rho_\tau$ .



We want to show that  $C_\tau$ , is in fact  $C^{2+\frac{1}{2}}$ .

By Theorem 3.8 we have the following representation of the solution of (11):

$$C(t, \theta) := \tilde{C}(t, \theta) + \int_0^t (-\partial_\theta \rho_u(0), \rho_u(0) - 2h_u) du - (0, \sqrt{2}B_t)$$

where

$$\tilde{C}(t, \theta) = \left( \int_0^\theta \frac{\cos(\theta_1)}{\rho_t(\theta_1)} d\theta_1, \int_0^\theta \frac{\sin(\theta_1)}{\rho_t(\theta_1)} d\theta_1 \right).$$

Since  $C(t_n, 0) \rightarrow C_\tau(0)$ , there exists  $A \in \mathbb{R}^2$  such that

$$\int_0^{t_n} (-\partial_\theta \rho_u(0), \rho_u(0) - 2h_x) du - (0, \sqrt{2}B_{t_n}) \rightarrow A.$$

Also since  $\rho_{t_n}$  converges uniformly to  $\rho_\tau$  and by Proposition 4.4,  $\rho_\tau > 0$ , we have that  $\tilde{C}(t_n, \cdot)$  converges uniformly to  $\left( \int_0^\cdot \frac{\cos(\theta_1)}{\rho_\tau(\theta_1)} d\theta_1, \int_0^\cdot \frac{\sin(\theta_1)}{\rho_\tau(\theta_1)} d\theta_1 \right)$ . Hence

$$C(t_n, \cdot) \rightarrow \left( \int_0^\cdot \frac{\cos(\theta_1)}{\rho_\tau(\theta_1)} d\theta_1, \int_0^\cdot \frac{\sin(\theta_1)}{\rho_\tau(\theta_1)} d\theta_1 \right) + A = C_\tau(\cdot)$$

By Lemma 4.14 we get that the curve  $C_\tau$  is  $C^{2+\frac{1}{2}}$ . Using Theorem 22 in [2], and the Markov property we can extend the solution after the time  $\tau$  by a solution starting at the curve  $C_\tau$ , which is in contradiction with the maximality of  $\tau$ .  $\square$

We have the following corollary of Theorem 61 in [2].

**Corollary 4.16.** *Let  $D_0$  be strictly convex and have  $n$  axes of symmetries, with  $n \geq 7$ , and consider  $(D_t)_{t \geq 0}$  the solution of (11). We have a.s. in the Hausdorff metric,*

$$\lim_{t \rightarrow +\infty} \frac{D_t}{\sqrt{\lambda(D_t)}} = B(0, 1/\sqrt{\pi})$$

where  $B(0, 1/\sqrt{\pi})$  is the Euclidean ball centered at 0 of radius  $1/\sqrt{\pi}$ .

## 5. SYMMETRIC CONVEX IN $\mathbb{R}^2$ WITH STAR SHAPED SKELETONS

Let  $\mathcal{C}$  be the set of smooth closed simple and strictly convex curves embedded in  $\mathbb{R}^2$ .

Fix  $n \geq 2$ . Let  $\mathcal{T}_n$  be the set of closed curves symmetric with respect to the vertical axis, denoted  $\Delta$ , and invariant by the rotation  $R_{2\pi/n}$  of angle  $2\pi/n$  (and thus invariant by the group  $G_n$  generated by these two isometries).

Let us describe the set  $\mathcal{C}$  in terms of its curvature. Let  $C_0 \in \mathcal{C}$ , and let  $C : \mathbb{T} \rightarrow \mathbb{R}^2$  be the parametrization of  $C_0$  such that  $\theta$  is the angle between the tangent line and the  $x$  axis at the point  $C(\theta)$  i.e a tangent vector is  $(\cos(\theta), \sin(\theta)) \in T_{C(\theta)}C$ . Note that this parametrization is possible since  $\partial_s \theta = \rho(\theta) > 0$  where  $s$  is the arc-length parametrization (due to Frénet equation). From now on, we will take this parametrization for curves in  $\mathcal{C}$ .

Recall from Lemma 4.1.1 of Gage and Hamilton [6] that a  $2\pi$  periodic positive function  $\rho$  represents the curvature of a simple closed strictly convex plane curve if and only if  $I_{c,\rho}(2\pi) = I_{s,\rho}(2\pi) = 0$ , where

$$I_{c,\rho}(\beta) := \int_0^\beta \frac{\cos(\theta)}{\rho(\theta)} d\theta, \quad I_{s,\rho}(\beta) := \int_0^\beta \frac{\sin(\theta)}{\rho(\theta)} d\theta \quad \beta \in \mathbb{T}.$$

More precisely, we have

$$\mathcal{C} \simeq \{\rho \in C^1(\mathbb{T}, ]0, \infty[) : I_{c,\rho}(2\pi) = I_{s,\rho}(2\pi) = 0\} \times \mathbb{R}^2$$

through the reciprocal bijections given by

$$\begin{aligned} \{\theta \mapsto C(\theta)\} &\longmapsto (\{\theta \mapsto \rho(\theta)\}, C(0)) \\ \{\theta \mapsto (I_{c,\rho}(\theta), I_{s,\rho}(\theta)) + X\} &\longleftarrow (\{\theta \mapsto \rho(\theta)\}, X) \end{aligned}$$

Let us describe the set  $\mathcal{C} \cap \mathcal{T}_n$  in terms of its curvature. Let  $C \in \mathcal{C} \cap \mathcal{T}_n$ . For any  $\theta \in \mathbb{T}$ , denote  $S_\theta$  the symmetry with respect to  $R_\theta(\Delta)$ . Using the symmetry  $S_0$  we have  $C(-\theta) = S_0(C(\theta))$  implying that  $C(0) = S_0(C(0))$  and

$$C(0) = (0, -b) \quad \text{for some } b \geq 0. \quad (27)$$

Using the symmetry  $S_{\pi/n} = R_{2\pi/n}S_0$  (thus belonging to  $G_n$ ) we have:  $C(2\pi/n - \theta) = S_{\pi/n}(C(\theta))$ , yielding for  $\theta = \pi/n$ :

$$C\left(\frac{\pi}{n}\right) = R_{\pi/n}((0, -a)) \quad \text{for some } a \geq 0. \quad (28)$$

The two numbers  $b, a$  are positive since  $(0, 0) \in \text{int}(C)$  by convexity. Also  $C$  is completely defined by its restriction to  $[0, \frac{\pi}{n}]$ . Using the invariance by  $G_n$  we have the following property of the associated curvature

$$\rho\left(\theta + \frac{\pi}{n}\right) = \rho\left(\frac{\pi}{n} - \theta\right), \theta \in \left[0, \frac{\pi}{n}\right], \quad \text{and } \rho \text{ is } \frac{2\pi}{n}\text{-periodic.} \quad (29)$$

So  $\partial_\theta \rho\left(\frac{k\pi}{n}\right) = 0$  for all  $k \in \{0, \dots, 2n-1\}$ .

A fundamental object for the study of elements of  $\mathcal{C} \cap \mathcal{T}_n$  will be the projection to some well chosen lines. Let  $C \in \mathcal{C} \cap \mathcal{T}_n$ . For  $\theta \in (0, \frac{\pi}{n}]$ , let  $(0, \Pi(\theta))$  be the intersection of the line  $D_\theta$  orthogonal to  $C$  at the point  $C(\theta)$  and the vertical axis  $\Delta$ . Define

$$\mathcal{S}_n := \left\{ C \in \mathcal{C} \cap \mathcal{T}_n, \Pi \text{ is increasing on } \left[0, \frac{\pi}{n}\right] \right\}. \quad (30)$$

Define also

$$\mathcal{S}_n^\downarrow := \left\{ C \in \mathcal{C} \cap \mathcal{T}_n, \rho \text{ is decreasing on } \left[0, \frac{\pi}{n}\right] \right\}. \quad (31)$$

Notice that for  $C \in \mathcal{S}_n$  or  $C \in \mathcal{S}_n^\downarrow$ , since  $C \in \mathcal{C} \cap \mathcal{T}_n$ , it is characterized by its values for  $\theta \in [0, \frac{\pi}{n}]$ .

**Proposition 5.1.** *Let  $C \in \mathcal{S}_n$ . Then  $\Pi(\pi/n) = 0$ , and  $\Pi$  has a limit  $-y_0 < 0$  as  $\theta \searrow 0$ , so it extends to a  $C^1$  nonpositive non-increasing function on  $[0, \pi/n]$ .*

*Proof.* Since the outward normal at  $C(\theta)$  is  $\nu(\theta) := (\sin(\theta), -\cos(\theta))$  we have for all  $\theta \in (0, \frac{\pi}{n})$

$$\Pi(\theta) = -b + \int_0^\theta \frac{\sin(\beta)}{\rho(\beta)} d\beta + \cot(\theta) \int_0^\theta \frac{\cos(\beta)}{\rho(\beta)} d\beta = -b + \int_0^\theta \frac{\cos(\theta - \beta)}{\rho(\beta) \sin(\theta)} d\beta$$

with  $b$  defined in (27), so

$$\lim_{\theta \searrow 0} \Pi(\theta) = -b + \frac{1}{\rho(0)} =: -y_0. \quad (32)$$

On the other hand, by symmetry of  $C$ , the point  $(0, \Pi(\pi/n))$  also belongs to  $R_{2\pi/n}(\Delta)$ , so  $\Pi(\pi/n) = 0$ . As a consequence, since we have assumed that  $\Pi$  is non-decreasing, we have  $y_0 > 0$  and  $\Pi$  is negative on  $[0, \pi/n)$ .

From now on we let  $\Pi(0) := -y_0$ .

Using an integration by part we have for  $\theta \in (0, \pi/n)$

$$\begin{aligned} \Pi'(\theta) &= \frac{1}{\rho(\theta) \sin(\theta)} - \frac{1}{\sin^2(\theta)} \int_0^\theta \frac{\cos(\beta)}{\rho(\beta)} d\beta \\ &= \frac{1}{\rho(\theta) \sin(\theta)} + \frac{1}{\sin^2(\theta)} \left( \left[ -\frac{\sin(\beta)}{\rho(\beta)} \right]_0^\theta - \int_0^\theta \frac{\rho'(\beta) \sin(\beta)}{\rho^2(\beta)} d\beta \right) \\ &= \frac{-1}{\sin^2(\theta)} \int_0^\theta \frac{\rho'(\beta) \sin(\beta)}{\rho^2(\beta)} d\beta. \end{aligned} \quad (33)$$

Note that

$$\Pi \in C^1 \left( \left(0, \frac{\pi}{n}\right) \right) \cap C^0 \left( \left[0, \frac{\pi}{n}\right] \right).$$

Taking into account that  $\lim_{\theta \searrow 0} \Pi'(\theta) = 0$ , due to  $\rho'(0) = 0$ , we end up with  $\Pi \in C^1([0, \frac{\pi}{n}])$ .  $\square$

The following result is a direct consequence of Equation (33):

**Proposition 5.2.** *We have  $\mathcal{S}_n^\downarrow \subset \mathcal{S}_n$ .*

Consider the mapping  $r$  defined by

$$\forall \theta \in [0, \pi/n], \quad r(\theta) := \|C(\theta) - (0, \Pi(\theta))\|$$

Since the curve does not cross the vertical axis before  $\frac{\pi}{n}$ ,  $r \in C^1((0, \frac{\pi}{n}) \cap C^0([0, \frac{\pi}{n}])$ . Hence we have the following parametrization of the curve  $C$ , for  $\theta \in (0, \frac{\pi}{n}]$

$$C(\theta) = (0, \Pi(\theta)) + r(\theta)(\sin(\theta), -\cos(\theta)) \quad (34)$$

**Lemma 5.3.** *The map  $\theta \mapsto (\Pi(\theta), r(\theta))$  extends to a  $C^1$  map defined on  $[0, \pi/n]$ , and satisfying*

$$(\Pi(0), r(0)) = \left( -b + \frac{1}{\rho(0)}, \frac{1}{\rho(0)} \right).$$

*Proof.* We are only left to prove the assertion for the map  $r$ . Putting the two parametrizations together, since  $\langle C'(\theta), N(\theta) \rangle = 0$  and  $\langle C'(\theta), T(\theta) \rangle = \frac{1}{\rho(\theta)}$ , from (15), we deduce from (34) that for any  $\theta \in (0, \pi/n]$ ,

$$\begin{aligned} \lim_{\theta \searrow 0} r(\theta) &= \frac{1}{\rho(0)}, \\ \begin{cases} -\Pi'(\theta) \cos(\theta) + r'(\theta) &= 0 \\ \Pi'(\theta) \sin(\theta) + r(\theta) &= \frac{1}{\rho(\theta)}, \end{cases} \end{aligned} \quad (35)$$

i.e

$$\begin{cases} \begin{pmatrix} \Pi(\theta) \\ r(\theta) \end{pmatrix}' + \begin{pmatrix} 0 & \frac{1}{\sin(\theta)} \\ 1 & \cot \theta \end{pmatrix} \begin{pmatrix} 0 \\ r(\theta) \end{pmatrix} = \begin{pmatrix} 0 & \frac{1}{\sin(\theta)} \\ 1 & \cot \theta \end{pmatrix} \begin{pmatrix} 0 \\ \frac{1}{\rho(\theta)} \end{pmatrix} \\ \begin{pmatrix} \Pi(\frac{\pi}{n}) \\ r(\frac{\pi}{n}) \end{pmatrix} = \begin{pmatrix} 0 \\ a \end{pmatrix} \end{cases} \quad (36)$$

where  $a$  is defined in (28). Using the first equation of (35), we get that  $\lim_{\theta \searrow 0} r'(\theta) = 0$ , so  $r \in C^1([0, \frac{\pi}{n}])$ .  $\square$

**Proposition 5.4.** *Let  $C$  be a curve in  $\mathcal{C} \cap \mathcal{T}_n$ .*

- (1) *If  $\Pi$  is non-decreasing on  $[0, \pi/n]$  (i.e. if  $C \in \mathcal{S}_n$ ), then the skeleton of  $C$  is  $G_n(\{0\} \times [-y_0, 0])$ .*
- (2) *If the skeleton of  $C$  is  $G_n(\{0\} \times [-y, 0])$  then  $\Pi$  is non-decreasing.*

*Proof.* (1) First assume that  $C \in \mathcal{S}_n$ . Denote by  $S$  the skeleton of  $C$ .

a) First we prove that  $G_n(\{0\} \times [-y_0, 0]) \subset S$ . For this it is sufficient to prove that for all  $\theta \in [0, \pi/n]$ , the point  $(0, \Pi(\theta))$  belongs to  $S$ .

We only need to prove it for  $\theta \in (0, \pi/n)$  since the skeleton is closed. For the same reason we can also assume that  $\Pi'(\theta) > 0$ . So let  $\theta \in (0, \pi/n)$  with  $\Pi'(\theta) > 0$ . The closed disk  $\bar{B}((0, \Pi(\theta)), r(\theta))$  centered at  $(0, \Pi(\theta))$  and with radius  $r(\theta)$  meets  $C$  at least at the two points  $C(\theta)$  and  $C(-\theta)$ . To prove that  $(0, \Pi(\theta)) \in S$  we need to prove that it is inside  $\bar{D}$ . This will be done in two steps.

- We prove that the set

$$\left\{ (0, \Pi(\theta)) + r(\cos \varphi, \sin \varphi) \mid 0 \leq r \leq r(\theta), -\frac{\pi}{2} - \frac{\pi}{n} \leq \varphi \leq -\frac{\pi}{2} + \frac{\pi}{n} \right\}$$


is included in  $\bar{D}$ .

The proof is by contradiction, assume there exists  $\theta' \in [0, \theta)$  such that  $\|C(\theta') - (0, \Pi(\theta))\| = r(\theta)$ . Consider the closed disk  $\mathcal{O}$  centred at  $(0, \Pi(\theta))$  of radius  $r(\theta)$ , passing through  $C(\theta)$  and  $C(\theta')$ , see Figure 1.

On one hand, by (35) we have  $r(\theta) < \frac{1}{\rho(\theta)}$ , so for  $\alpha < \theta$  and  $\alpha$  close to  $\theta$ , the points  $C(\alpha)$  are outside the disk  $\mathcal{O}$ . On the other hand, since  $\Pi(\alpha) < \Pi(\theta)$ , there exists  $q_\alpha \in D_\alpha \cap ((0, \Pi(\theta)), C(\theta'))$ . As we can see in the proof of the point b) below,  $C(\alpha)$  is the nearest point of  $q_\alpha$  in  $C$ . We have  $\|q_\alpha - C(\alpha)\| \leq \|q_\alpha - C(\theta')\|$ . Hence

$$\begin{aligned} \|C(\alpha) - (0, \Pi(\theta))\| &< \|q_\alpha - C(\alpha)\| + \|q_\alpha - (0, \Pi(\theta))\| \\ &\leq \|q_\alpha - C(\theta')\| + \|q_\alpha - (0, \Pi(\theta))\| = r(\theta) \end{aligned}$$

and we get a contradiction.



A similar contradiction is obtained if we assume there exists  $\theta' \in (\theta, \pi/n]$ , with  $\|C(\theta') - (0, \Pi(\theta))\| = r(\theta)$ .

We get the wanted inclusion.

- We easily check that the convex hull  $H(\theta)$  of the  $n$  pieces of disks

$$G_n \left( \left\{ (0, \Pi(\theta)) + r(\cos \varphi, \sin \varphi) \mid 0 \leq r \leq r(\theta), -\frac{\pi}{2} - \frac{\pi}{n} \leq \varphi \leq -\frac{\pi}{2} + \frac{\pi}{n} \right\} \right)$$

contains  $\bar{B}((0, \Pi(\theta)), r(\theta))$  (check for instance that the curvature of its boundary is everywhere smaller than  $1/r(\theta)$ ). But  $H(\theta) \subset \bar{D}$  since  $\bar{D}$  is left invariant by  $G_n$  and convex. As a conclusion,  $\bar{B}((0, \Pi(\theta)), r(\theta)) \subset \bar{D}$ , so  $(0, \Pi(\theta)) \in S$ .

b) Finally we prove that  $S \subset G_n(\{0\} \times [-y_0, 0])$ . For  $\theta \in [0, \pi/n]$  and  $r \in (0, r(\theta))$ , consider the point  $P = (0, \Pi(\theta)) + r\nu(\theta)$ . We have to prove that it does not belong to  $S$ . Consider  $\theta' \in [0, 2\pi)$  such that  $C(\theta')$  minimizes the distance between  $P$  and  $C$ . First note that we must have  $\theta' \in [0, \pi/n]$ , otherwise the minimizing segment would cross an axis of symmetry, allowing to construct a shorter segment from  $P$  to  $C$ . Next let us show that necessarily  $\theta' = \theta$ . Indeed, otherwise, the lines  $D_\theta$  and  $D_{\theta'}$  would then intersect at  $P$ . Assume for instance that  $\theta < \theta'$ , then we would get that  $\Pi(\theta) > \Pi(\theta')$ , which is forbidden. Finally, since  $d(P, C(\theta)) < r(\theta) \leq 1/\rho(\theta)$ , the distance to  $C$  is not singular at  $P$  and  $P$  cannot belong to  $S$ . Using all symmetries, this proves that the complementary of  $G_n(\{0\} \times [-y_0, 0])$  in  $\bar{D}$  does not meet the cutlocus  $S$  of distance to  $C$ .

(2) Assume that the skeleton of  $C$  is  $G_n(\{0\} \times [-y_0, 0])$ . Then for all  $\theta \in (0, \pi/n)$ , we have  $B((0, \Pi(\theta)), r(\theta)) \subset D$ . This implies that  $r(\theta) \leq 1/\rho(\theta)$ . Then by (35) we get

$$\Pi'(\theta) = \frac{\frac{1}{\rho(\theta)} - r(\theta)}{\sin(\theta)} \geq 0,$$

so  $\Pi$  is non-decreasing. □

**Proposition 5.5.** *The set of curve  $\mathcal{S}_n^\downarrow$  is stable under the stochastic mean curvature flow namely (20). It is also stable under the usual deterministic mean curvature flow.*

*Proof.* Let  $C_0$  be a curve in  $\mathcal{S}_n^\downarrow$ , and  $\rho_0$  the associated curvature function, by hypothesis  $\partial\rho_0(\theta) \leq 0$  for  $\theta \in [0, \frac{\pi}{n}]$ . Let  $C_t$  be the solution of the stochastic mean curvature started at  $C_0$ , namely the solution of:

$$\begin{cases} d_t C(t, u) &= (-\rho_t(C(t, u)))\nu_{C(t, u)} dt + \sqrt{2}\nu_{C(t, u)} dB_t \\ C(0, u) &= C_0(u). \end{cases}$$

Using the parametrization by the angle  $\theta$  of the tangent vector and the horizontal axis as above we have, denoting  $\rho(t, \theta) := \rho_t(\theta)$ ,

$$\begin{cases} d_t \rho(t, \theta) = \rho^2(t, \theta)(\partial_\theta^2 \rho(t, \theta))dt + \rho^2(t, \theta)(3\rho(t, \theta)dt - \sqrt{2}dB_t), \\ \rho(0, \cdot) = \rho_0. \end{cases}$$

(see (16) with  $h$  replaced by 0).

Using Lemme 3.9 we get that  $\rho_t > 0$  for  $t < \tau$  where  $\tau$  is any lifetime of the stochastic mean curvature flow. Using Itô formula, we have for  $0 \leq t < \tau$

$$d \frac{1}{\rho_t(\theta)} = (-\partial_\theta^2 \rho_t(\theta) - \rho_t(\theta))dt + \sqrt{2}dB_t, \quad (37)$$

Computations similar to those of the proof of Theorem 3.8 show that  $I_{c, \rho_t}(2\pi) = I_{s, \rho_t}(2\pi) = 0$ . Recall  $S_0$  is the reflection with respect to the vertical axis. Using the uniqueness of the stochastic mean curvature flow, we have

$$S_0(C_t(C_0)) = C_t(S_0(C_0)) = C_t(C_0).$$

Doing the same thing with the rotation  $R_{2\pi/n}$ , it follows that  $\rho_t$  satisfies Equation (29). To get the result we only have to show that  $\partial_\theta \rho_t(\theta) \leq 0$  for  $0 \leq t < \tau$  and  $\theta \in (0, \frac{\pi}{n})$ .

Differentiating (37) in  $\theta$  we get:

$$d \left( \frac{\partial_\theta \rho_t(\theta)}{\rho_t^2(\theta)} \right) = \partial_\theta^2 (\partial_\theta \rho_t(\theta))dt + \partial_\theta \rho_t(\theta)dt.$$

Let  $\psi_t(\theta) = \frac{\partial_\theta \rho_t(\theta)}{\rho_t^2(\theta)}$ , then  $\psi$  satisfies the following partial differential equation with stochastic coefficient and with lifetime  $\tau$  (see Lemma 3.9):

$$\begin{aligned} \partial_t \psi_t(\theta) &= \partial_\theta^2 (\psi_t(\theta) \rho_t^2(\theta)) + \psi_t(\theta) \rho_t^2(\theta) \\ &= \rho_t^2(\theta) \partial_\theta^2 \psi_t(\theta) + 4\rho_t(\theta) (\partial_\theta \psi_t(\theta)) (\partial_\theta \rho_t(\theta)) + \psi_t(\theta) (\partial_\theta^2 \rho_t^2(\theta) + \rho_t^2(\theta)) \end{aligned}$$

with initial condition  $\psi_0(\theta) = \frac{\partial_\theta \rho_0(\theta)}{\rho_0^2(\theta)}$ . By hypothesis  $\psi_0(\theta) \leq 0$ . Note also by the conservation of the symmetry that we have the boundary conditions  $\psi_t(0) = \psi_t(\frac{\pi}{n}) = 0$ . To show that  $\partial_\theta \rho_t(\theta) \leq 0$  for all  $t < \tau$  we will argue by contradiction. Suppose that there exists  $t^* < \tau$  and  $\theta \in [0, \frac{\pi}{n}]$  such that  $\partial_\theta \rho_{t^*}(\theta) > 0$  so  $\psi_{t^*}(\theta) > 0$ . Let

$$\mu = -2 \left( \|\partial_\theta^2 \rho^2(\cdot)\|_{[0, t^*] \times [0, \frac{\pi}{2}]} + \|\rho^2(\cdot)\|_{[0, t^*] \times [0, \frac{\pi}{2}]} \right) > -\infty,$$

and  $W_t(\theta) = e^{\mu t} \psi_t(\theta)$ , which satisfies the following equation:

$$\begin{aligned} & \partial_t W_t(\theta) \\ &= \rho_t^2(\theta) \partial_\theta^2 W_t(\theta) + 4\rho_t(\theta) (\partial_\theta \rho_t(\theta)) (\partial_\theta W_t(\theta)) + W_t(\theta) (\partial_\theta^2 \rho_t^2(\theta) + \rho_t^2(\theta) + \mu). \end{aligned} \quad (38)$$

Define  $\alpha := \sup_{\theta \in [0, \frac{\pi}{n}]} W_{t^*}(\theta) > 0$ ,

$$t_0 := \inf \left\{ t \leq t^*, \text{ s.t. } \sup_{\theta \in [0, \frac{\pi}{n}]} W_t(\theta) = \alpha \right\}$$

and let  $\theta^*$  be such that  $W_{t_0}(\theta^*) = \alpha$ . From boundary conditions we have  $\theta^* \in ]0, \frac{\pi}{n}[$ . At  $(t_0, \theta^*)$  we have

$$\partial_t W_t(\theta^*)|_{t_0} \geq 0, \quad \partial_\theta^2 W_{t_0}(\theta)|_{\theta^*} \leq 0, \quad \partial_\theta W_{t_0}(\theta)|_{\theta^*} = 0.$$

Using equation (38) we get the contradiction, since

$$0 \leq \partial_t W_t(\theta^*)|_{t_0} \leq \alpha \frac{\mu}{2} < 0.$$

With a similar proof, we get the second part, namely the conservation of the class  $\mathcal{S}_n^\downarrow$  under the usual deterministic mean curvature flow.  $\square$

**Corollary 5.6.** *The class of domain  $\mathcal{S}_n^\downarrow$  is also stable under the normalized stochastic mean curvature flow (11).*

*Proof.* Since the solutions of (11) are obtained by a change of probability from the solutions of the stochastic mean curvature flow, the state space does not change, and the result follows from Proposition 5.5.  $\square$

## 6. A NEW ISOPERIMETRIC ESTIMATE

Let us end our consideration of  $\mathcal{S}_n$  by observing that its elements are quite round when  $n^2$  is much larger than the length of their skeleton:

**Proposition 6.1.** *For any curve  $C$  in the set  $\mathcal{S}_n$  defined in (30) (and in particular with skeleton  $G_n(\{0\} \times [-L(C)/n, 0])$ ) we have*

$$\begin{aligned} \pi^2 (r_{\text{out}} - r_{\text{int}})^2 &\leq \sigma^2(C) - 4\pi \text{vol}(D) \\ &\leq \frac{2\pi^2}{n^2} L(C)^2 \left( 1 - \frac{\sin\left(\frac{2\pi}{n}\right)}{\frac{2\pi}{n}} \right) \\ &\leq \frac{4\pi^4}{3n^4} L(C)^2, \end{aligned}$$

where  $L(C)$  is the length of the skeleton of  $C$ .

*Proof.* The lower bound on  $\sigma^2(C) - 4\pi \text{vol}(D)$  is just Bonnesen inequality (6). For the upper bound, let  $\rho$  be the curvature function associated to  $C$ , and  $p(\theta) = \langle C(\theta), \nu(\theta) \rangle$  the support function. Using computation in (35) we have

$$p(\theta) = -\Pi(\theta) \cos(\theta) + r(\theta),$$

$$p'(\theta) = \Pi(\theta) \sin(\theta)$$

$$p''(\theta) + p(\theta) = \frac{1}{\rho(\theta)}.$$

By symmetry of  $C$  we have the following Fourier series of  $p$ :

$$p(\theta) = a_0 + \sum_{k \geq 1} a_k \cos(kn\theta).$$

Also  $\text{vol}(D) = \frac{1}{2} \int_0^{2\pi} p(\theta)(p(\theta) + p''(\theta))d\theta = \pi a_0^2 + \frac{\pi}{2} \sum_{k \geq 2} a_k^2(1 - n^2 k^2)$  and  $a_0 = \frac{1}{2\pi} \int p(\theta)d\theta = \frac{1}{2\pi} \sigma(C)$ . Hence

$$\begin{aligned} \sigma^2(C) - 4\pi \text{vol}(D) &= 2\pi^2 \sum_{k \geq 1} a_k^2(n^2 k^2 - 1) \\ &\leq 2\pi \int_0^{2\pi} p'(\theta)^2 d\theta \\ &= 4n\pi \int_0^{\pi/n} \Pi^2(\theta) \sin^2(\theta) d\theta \\ &\leq 4n\pi \left(\frac{L(C)}{n}\right)^2 \int_0^{\pi/n} \sin^2(\theta) d\theta \\ &= \frac{2\pi^2}{n^2} L(C)^2 \left(1 - \frac{\sin\left(\frac{2\pi}{n}\right)}{\frac{2\pi}{n}}\right) \\ &\leq \frac{4\pi^4}{3n^4} L(C)^2 \end{aligned}$$

since  $1 - \sin(x)/x \leq x^2/6$  for any  $x \in \mathbb{R}$  (with the usual convention  $\sin(0)/0 = 1$ ).  $\square$




FIGURE 2.



## REFERENCES

- [1] Marc Arnaudon, Koléhè Coulibaly-Pasquier and Laurent Miclo. Construction of set-valued dual processes on manifolds. Preprint on the open repository *Hal*, December 2020.
- [2] Koléhè Coulibaly-Pasquier and Laurent Miclo. On the evolution by duality of domains on manifolds. *Mém. Soc. Math. Fr., Nouv. Sér.*, 171:1–110, 2021.
- [3] Michael E. Gage. An isoperimetric inequality with applications to curve shortening. *Duke Mathematical Journal*, volume 50, pages 1225–1229, 1983.
- [4] Michael E. Gage. Curve shortening makes convex curves circular. *Invent. Math.*, volume 76, 1984.
- [5] Michael E. Gage. On an area-preserving evolution equation for plane curves. *Nonlinear problems in geometry* (Mobile, Ala., 1985), volume 51, pages 51–62, 1986.
- [6] Michael E. Gage and Richard S. Hamilton. The heat equation shrinking convex plane curves. *Journal of Differential Geometry*, volume 23, pages 69–96, 1986.
- [7] Matthew A. Grayson. The heat equation shrinks embedded plane curves to round points. *J. Differential Geom.* 26 (1987), pp. 285–314
- [8] Mark Green and Stanley Osher. Steiner polynomials, Wulff flows, and some new isoperimetric inequalities for convex plane curves. *Asian J. Math.*, 3(3):659–676, 1999.
- [9] Alessandra Lunardi. Analytic semigroups and optimal regularity in parabolic problems. Modern Birkhäuser Classics. Birkhäuser/Springer Basel AG, Basel, 1995.

INSTITUT DE MATHÉMATIQUES DE BORDEAUX, UMR 5251,  
 UNIV. BORDEAUX, CNRS, BORDEAUX INP  
*Email address:* marc.arnaudon@math.u-bordeaux.fr

INSTITUT ÉLIE CARTAN DE LORRAINE, UMR 7502,  
 UNIVERSITÉ DE LORRAINE AND CNRS  
*Email address:* kolehe.coulibaly@univ-lorraine.fr

TOULOUSE SCHOOL OF ECONOMICS, UMR 5314,  
 CNRS AND UNIVERSITÉ DE TOULOUSE  
*Email address:* laurent.miclo@math.cnrs.fr