

Computation of L_{\oplus} for several cubic Pisot numbers

Bernat Julien

Institut de Mathématiques de Luminy
 UMR 6206
 Campus de Luminy, Case 907
 13288 MARSEILLE Cedex 9
 FRANCE
 email: bernat@iml.univ-mrs.fr

In this article, we are dealing with β -numeration, which is a generalization of numeration in a non-integer base. We consider the class of simple Parry numbers such that $d_{\beta}(1) = 0.k_1^{d-1}k_d$ with $d \in \mathbb{N}$, $d \geq 2$ and $k_1 \geq k_d \geq 1$. We prove that these elements define Rauzy fractals that are stable under a central symmetry. We use this result to compute, for several cases of cubic Pisot units, the maximal length among the lengths of the finite β -fractional parts of sums of two β -integers, denoted by L_{\oplus} . In particular, we prove that $L_{\oplus} = 5$ in the Tribonacci case.

Keywords: Tribonacci number, β -numeration, β -integers, Rauzy fractal.

Contents

1	Definitions and notation	4
1.1	Substitutions	4
1.2	Introduction to β -numeration	5
1.3	Notion of β -substitution	6
1.4	Rauzy fractals	6
2	Rauzy fractals and symmetric sets	7
2.1	Set of β -integers and admissibility	7
2.2	The symmetric property for confluent Parry numbers	8
3	Computation of L_{\oplus} in several cases of cubic Pisot numbers	10
3.1	Fractional parts of sums or differences of β -integers	10
3.2	Algorithmic computation of L_{\oplus} in several cubic Pisot cases	13
3.3	The particular case of Tribonacci	14

Introduction

This article deals with β -numeration, which is a generalization of numeration in a non-integer base. The β -numeration is born in the late 50's [35, 31]. It has known a major realization with the arrival of quasicrystals during the 80's [38], since the β -numeration allows a modelling of quasicrystals [13, 10]. The number systems defined by the β -numeration are closely related to canonical number systems [6, 36], number systems generated by iterated function systems [39] or by substitutive systems of Pisot type [15]. A common feature between these fields is the property of self-similarity, which yields results in number theory [7], geometry [24], topology [8, 14], dynamical systems [40, 33, 34] and theoretical computer science [9]. Nowadays, the interactions between these domains still set up the core of the research on β -numeration. We focus with this article on applications in computer arithmetics.

Let $\beta > 1$ be an algebraic integer, with $\beta \notin \mathbb{N}$. In the same way as in the case of integral bases, it is possible to expand $x \in [0, 1]$ as $x = \sum_{k \in \mathbb{N}^*} \varepsilon_k \beta^{-k}$, where the sequence $(\varepsilon_k)_{k \in \mathbb{N}^*}$, called *expansion of x in base β* , is defined on the alphabet $\mathcal{A}_\beta = \{0, \dots, [\beta]\}$. Among the expansions of 1 in base β , the greatest for the lexicographic order is called *β -expansion of 1*, and denoted by $d_\beta(1)$. This expansion is constructed by the *greedy algorithm*, that is, $d_\beta(x) = 0.\varepsilon_1 \dots \varepsilon_n \dots$, with for all $k \in \mathbb{N}^*$, $\varepsilon_k = [\beta T_\beta^{k-1}(x)]$, where $T_\beta : [0, 1] \rightarrow [0, 1]$, $x \mapsto \{\beta x\}$. The notion of β -expansion was historically introduced by Rényi [35]. Parry produced in [31] many interesting results concerning the β -numeration, creating and studying among others a dynamical system associated to this numeration.

Let us observe that a natural expansion of 1 in base β is 1. However, defining $d_\beta(1)$ as the greatest sequence for the lexicographic order among expansions of 1 in base β provides useful algebraic informations on β . This is why $d_\beta(1)$ is not defined as 1. Another interesting expansion of 1 in base β is the greatest sequence for the lexicographical order among those which do not end by 0's. This expansion, denoted by $d_\beta^*(1)$, is an improper expansion of 1 in base β . At the opposite of the case of numerations in integer bases, $d_\beta^*(1)$ is worthwhile since it enables to define the Parry condition, defined below, which plays a key role in the construction and the study of the associated arithmetical language \mathcal{L}_β . Notably, Parry showed that for any $x \in [0, 1[$, $d_\beta(x)$ is the only expansion of x in base β which satisfies the following condition called the *Parry condition* [31]:

$$\text{for all } n \in \mathbb{N}, S^n((\varepsilon_k)_{k \in \mathbb{N}^*}) <_{lex} d_\beta^*(1),$$

where S is the shift map: $\mathcal{A}_\beta^{\mathbb{N}^*} \rightarrow \mathcal{A}_\beta^{\mathbb{N}^*}$ defined by $S((\varepsilon_k)_{k \in \mathbb{N}^*}) = (\varepsilon_{k+1})_{k \in \mathbb{N}^*}$. The factors of a sequence in $\mathcal{A}_\beta^{\mathbb{N}^*}$ that fulfills the Parry condition are said to be *admissible*, and they form the arithmetical language \mathcal{L}_β .

In the same way as for elements of $[0, 1]$, any positive real number $x > 1$ can be uniquely expanded as $x = \sum_{i=0}^n \varepsilon_{-i} \beta^i + \sum_{i \in \mathbb{N}^*} \varepsilon_i \beta^{-i}$, where $(\varepsilon_i)_{i \geq -n}$ satisfies the Parry condition. The part with non-negative powers of β is then called the *β -integer part of x* , denoted by $[x]_\beta$; the part with negative powers of β is called the *β -fractional part of x* , denoted by $\{x\}_\beta = x - [x]_\beta$. This allows a natural generalization for the definition of integers in base β .

Definition 0.1. *The set of β -integers, denoted by \mathbb{Z}_β , is the set of real numbers x for which there exists $n \in \mathbb{N}$ such that $x = \pm \sum_{k=0}^n v_k \beta^k$, where $v_n \dots v_0$ is an admissible word.*

The following set, $\text{Fin}(\beta)$, is introduced in [20]. It allows to generalize the framework of numeration to the case of a non-integral base, since the elements of $\text{Fin}(\beta)$ play the role of decimal numbers in base β .

Definition 0.2. *The set of real numbers having a finite β -fractional part is*

$$\text{Fin}(\beta) = \bigcup_{k \in \mathbb{N}} \beta^{-k} \mathbb{Z}_\beta.$$

Remark 1. *For a given algebraic number $\beta > 1$, the relation $\text{Fin}(\beta) = \mathbb{Z}[\beta^{-1}]$ is known as the finiteness property, denoted by (\mathcal{F}) . Whereas not yet fully characterized, the class of numbers satisfying the finiteness property has been extensively studied, and relations with ergodic and combinatorial properties of the associated dynamical system have been highlighted. Moreover, the finiteness property provides topologic and geometric properties for the geometrical realization of the associated dynamical system known as the Rauzy fractal, defined in Section 1.4. For instance, due to Akiyama [2], the finiteness property implies that 0 is an inner point of the Rauzy fractal \mathcal{T} under the additional condition that β is a unit. Also, Theorem 3 of the same article provides a characterization of numbers such that \mathcal{T} is arcwise connected. One may refer to [1, 4, 20] for additional informations.*

The sets \mathbb{Z}_β and $\text{Fin}(\beta)$ are not stable under usual operations like addition and multiplication. For example, when β is the golden ratio $\frac{1+\sqrt{5}}{2}$, which is called the *Fibonacci case*, one checks that $\beta^{-1} + \beta^{-2} = 1$. Then $d_\beta(1) = 0.11$, and the set of admissible words consists of words defined on the alphabet $\{0, 1\}$ which do not contain two consecutive occurrences of 1. The Fibonacci case is the most studied one, since $\frac{1+\sqrt{5}}{2}$ is the only Pisot number of degree two which belongs to $]1, 2[$. This means that it defines from an

algebraic point of view the simplest case of numeration in a non-integer base. See for instance [26] for several properties of this numeration. It is clear that $1 \in \mathbb{Z}_{\beta}$, since $1 \in \mathcal{A}_{\beta}$ holds for any $\beta > 1$. Since $1 + 1 = 2 = \beta + \beta^{-2}$, one has $d_{\beta}(2) = 10.01$, hence \mathbb{Z}_{β} is not stable under addition.

Likewise, let β be the positive root of the polynomial $X^4 - 2X^3 - X - 1$. Then $d_{\beta}(1) = 0.2011$, with $1, 2 \in \mathbb{Z}_{\beta}$ and $d_{\beta}(3) = 10.111(00012)^{\infty}$, hence the sum of two β -integers can have an infinite β -expansion.

In order to perform arithmetics on β -integers, say, for instance, to compute the addition of two β -integers, one must be able to renormalize expansions in base β of real numbers obtained after adding β -integers. The renormalization step is performed by transducers, see for instance [18]. Another method consists in applying the greedy algorithm to the real number obtained after adding or multiplying β -integers, which produces its β -expansion. However, if the β -expansion of the sum of two β -integers is neither finite nor ultimately periodic, the required β -expansion will not be produced after a finite number of steps.

The following notations, L_{\oplus} and L_{\odot} , are introduced in [22]. They represent the maximal possible finite length of the β -fractional parts which may appear when one adds or multiplies two β -integers. The computation of these values gives an indication on the difficulty of performing arithmetics on \mathbb{Z}_{β} .

Definition 0.3. *The bound L_{\oplus} is defined as $\min\{n \in \mathbb{N} \mid \forall x, y \in \mathbb{Z}_{\beta}, x + y \in \text{Fin}(\beta) \Rightarrow \beta^n(x + y) \in \mathbb{Z}_{\beta}\}$ when this set is not empty, $+\infty$ otherwise.*

The bound L_{\odot} is defined as $\min\{n \in \mathbb{N} \mid \forall x, y \in \mathbb{Z}_{\beta}, xy \in \text{Fin}(\beta) \Rightarrow \beta^n xy \in \mathbb{Z}_{\beta}\}$ when this set is not empty, $+\infty$ otherwise.

Let us explain now why we are interested in the case where L_{\oplus} and L_{\odot} are finite. Indeed, if the sum or the product of two β -integer belongs to $\text{Fin}(\beta)$, then the length of the β -fractional part of this sum or product is bounded by a constant which only depends on β . In this case, one can decide whether a given improper expansion can be renormalized into a finite or an ultimately periodic expansion, in the sense that, if during the renormalization process one gets a β -fractional part of length greater than L_{\oplus} , then the improper expansion corresponds to a real number that does not belong to $\text{Fin}(\beta)$. Conversely, if the set of lengths of sums of two β -integers is unbounded, then performing arithmetics in \mathbb{Z}_{β} will be very difficult if not impossible, since one cannot compute in a finite time any operation on β -integers.

It has been proven that L_{\oplus} and L_{\odot} are finite when β is a Pisot number, respectively in [20] and [22]. The computation of these values is however not so easy, especially for L_{\odot} . The case of quadratic Pisot numbers has been studied in [13] when β is a unit, and in [23] otherwise. However, when β is of higher degree, it is a difficult problem to compute the exact value of L_{\oplus} or L_{\odot} , and even to compute upper and lower bounds for these two constants. Several examples are studied in [9], where a method is described in order to compute upper bounds for L_{\oplus} and L_{\odot} for Pisot numbers satisfying additional algebraic properties. We prove in a forthcoming article [11] that the bounds L_{\oplus} and L_{\odot} are finite for the more general case of Perron numbers.

Let us detail now connections between arithmetics in base β and the study of some mathematical structures, namely, model sets and Meyer sets, which allow a modelling of quasicrystals. For any $\beta > 1$, \mathbb{Z}_{β} is a discrete subset of \mathbb{R} . The question of finding an algebraic characterization of numbers β such that \mathbb{Z}_{β} is uniformly discrete is still open; however it is known that \mathbb{Z}_{β} is uniformly discrete when β is a Parry number. Since \mathbb{Z}_{β} is relatively dense in \mathbb{R} for all $\beta > 1$, \mathbb{Z}_{β} is a Delone set [25] as soon as \mathbb{Z}_{β} is uniformly discrete. If moreover there exists F , a finite set, such that $\mathbb{Z}_{\beta} + \mathbb{Z}_{\beta} \subset \mathbb{Z}_{\beta} + F$, \mathbb{Z}_{β} is said to be a Meyer set [29]. Such sets satisfy a strong property of regularity. A particular subset of Meyer sets is the class of model sets, which are produced by a cut-and-project scheme. Roughly speaking, these sets are obtained as the intersection of the lattice \mathbb{Z}^d with a window, that is, a compact set of a space called internal space, that is pushed up along a half-line of another space called physical space. In the particular case of β -numeration, the window is the Rauzy fractal \mathcal{T} defined in Section 1.4. See also [28, 30] for additional details.

When \mathbb{Z}_{β} is a Meyer set, a method which enables the computation of a minimal finite set F satisfying $\mathbb{Z}_{\beta} + \mathbb{Z}_{\beta} \subset \mathbb{Z}_{\beta} + F$ is defined in [5] and studied in [21], where the problem of recognizing the language which consists of sums of β -integers is studied as well. This question is related to the determination of L_{\oplus} , since one can choose the sets of β -fractional parts of length less than L_{\oplus} for F when the finiteness property (\mathcal{F}) holds. Note however that the set of β -fractional parts of length less than L_{\oplus} is not optimal, since they may exist another set F_0 which contains less elements such that $\mathbb{Z}_{\beta} + \mathbb{Z}_{\beta} \subset \mathbb{Z}_{\beta} + F_0$. Let us

remind that \mathbb{Z}_β is a Meyer set when β is a Pisot number [13], and that \mathbb{Z}_β is a Meyer set implies that β is either a Pisot or a Salem number [29].

The aim of this paper is, first to study a geometrical property for a particular subclass of Parry numbers, then to apply these results in order to improve the computation of an upper bound for L_\oplus . This allows us to determine the exact value of L_\oplus for several cases of cubic Pisot numbers. This article is structured in the following way. Section 1 gathers definitions and notations that are introduced in the framework of β -numeration. Section 2 is devoted to the class of β which satisfy the following relation:

$$\text{there exist } d \geq 2, k_1, k_d \in \mathbb{N}, k_1 \geq k_d \text{ such that } d_\beta(1) = 0.k_1^{d-1}k_d. \quad (1)$$

Such numbers define confluent numeration systems, introduced and studied by Frougny [17]. This is why we will refer to *confluent Parry numbers* for any Parry number satisfying (1) from now on. We find the following geometric interpretation for confluent Parry numbers.

Proposition 2.5. Let β be a confluent Parry number. There exists $\vec{C} \in \mathbb{R}^{d-1}$ such that the Rauzy fractal \mathcal{T} is stable under the central symmetry $S_{\vec{C}} : \mathbb{R}^{d-1} \rightarrow \mathbb{R}^{d-1}, \vec{X} \mapsto 2\vec{C} - \vec{X}$.

For a given Pisot number β , we introduce and study a symmetric centered set \mathcal{T}' , which contains the Rauzy fractal \mathcal{T} . The equality $\mathcal{T}' = \mathcal{T}$ holds exactly when β is a confluent Parry number. The study of \mathcal{T}' in Section 3.1 provides inequalities satisfied by any Galois conjugate of β (Propositions 3.4, 3.5). This allows us to improve a method detailed in [9], which enables the computation of upper and lower bounds for L_\oplus under an algebraic condition on β . In particular, we determine the exact value of L_\oplus for several cases of cubic Pisot units in Section 3.2. Finally, we focus in Section 3.3 on the *Tribonacci* case, that is, when β is the positive root of the polynomial $X^3 - X^2 - X - 1$. We improve by the following theorem the result $5 \leq L_\oplus \leq 6$ [9].

Theorem 3.9. When $d_\beta(1) = 0.111$, we have $L_\oplus = 5$.

1 Definitions and notation

We refer mainly for the notation introduced here to [32] and [26].

1.1 Substitutions

Let \mathcal{A} be a finite set that is called *alphabet*. Endowed with the concatenation, \mathcal{A} generates a monoid called *set of finite words*, denoted by \mathcal{A}^* . Let ε denote the empty word. When \mathcal{A} is a totally ordered set, the sets $\mathcal{A}^{\mathbb{Z}}$, $\mathcal{A}^{\mathbb{N}}$ and \mathcal{A}^* are endowed with the lexicographic order $<_{lex}$.

Let $v = v_1 \dots v_n \in \mathcal{A}^*$. The number of letters of v , called *length* of v , is denoted by $|v|$; the number of occurrences of $a \in \mathcal{A}$ in v is denoted by $|v|_a$. A word w is called *factor* of v if there exists $p, s \in \mathcal{A}^*$ such that $v = pws$. The *mirror image* of $v = v_1 \dots v_n$ is defined as $\tilde{v} = v_n \dots v_1$, and the *mirror image map* as the map $\mathcal{A}^* \rightarrow \mathcal{A}^*, u \mapsto \tilde{u}$. A word $v \in \mathcal{A}^*$ is said to be a *palindrome* when $\tilde{v} = v$. When $v \in \mathcal{A}^{\mathbb{N}}$, we set $|v| = +\infty$; a word w is said to be a factor of v if there exists v' , a prefix of v , such that w is a factor of v' .

Let $\mathcal{A} = \{a_1, \dots, a_d\}$. Let $(\vec{e}_j)_{j \in \llbracket 1, \dots, d \rrbracket}$ be the canonical basis of \mathbb{Z}^d . The morphism $f : \mathcal{A}^* \rightarrow \mathbb{Z}^d$ that fulfills $f(a_j) = \vec{e}_j$ for all $j \in \llbracket 1, \dots, d \rrbracket$ is called the *abelianization homomorphism*.

Any map $\sigma : \mathcal{A} \rightarrow \mathcal{A}^*$, extended to a morphism on \mathcal{A}^* by concatenation, is called *substitution*. The substitution σ is said *non-erasing* if for every $a \in \mathcal{A}$, $\sigma(a) \neq \varepsilon$. Let σ be a substitution defined on the alphabet \mathcal{A} . The *incidence matrix* of σ is defined as the square matrix $M_\sigma \in \mathcal{M}_d(\mathbb{N})$ such that, for every $(i, j) \in \llbracket 1, \dots, d \rrbracket^2$, $M_\sigma[i, j] = |\sigma(a_j)|_{a_i}$. The substitution is *primitive* if there exists $n \in \mathbb{N}$ such that for every $(i, j) \in \llbracket 1, \dots, d \rrbracket^2$, $M_\sigma^n[i, j] \geq 1$. The substitution is said to be *unimodular* if $|\det M_\sigma| = 1$. The substitution is said to be *of Pisot type* if the eigenvalues of the incidence matrix M_σ satisfy the following: there exists a dominant eigenvalue $\beta > 1$ such that for every other eigenvalue α , one has $0 < |\alpha| < 1$. Note that the characteristic polynomial of M_σ is irreducible when σ is of Pisot type. Let us remind that a Pisot number is an algebraic integer whose any Galois conjugate $\alpha \neq \beta$ satisfies $|\alpha| < 1$, and that the dominant eigenvalue of M_σ is a Pisot number when σ is of Pisot type.

Let σ be a substitution. If there exists $k \in \mathbb{N}^*$ such that $\sigma^k(\omega) = \omega$, we say that ω is a *periodic point* of σ . In particular, when $\sigma(\omega) = \omega$, ω is said to be a *fixed point* of σ . When σ is primitive, we denote by \mathcal{L}_σ the language generated by the substitution, that is, $\mathcal{L}_\sigma = \bigcup_{k \in \mathbb{N}} \{\text{factors of } \sigma^k(a)\}$, where $a \in \mathcal{A}$. Note that, since we assume that σ is primitive, the definition of \mathcal{L}_σ does not depend on the choice of a .

1.2 Introduction to β -numeration

Let $\beta > 1$ be an algebraic integer. The set $\mathbb{N} \cap [0, \beta[$ is called β -*alphabet* and denoted by \mathcal{A}_β . Let $x \in [0, 1]$. Any sequence $(\varepsilon_k)_{k \in \mathbb{N}^*} \in \mathcal{A}_\beta^{\mathbb{N}^*}$ satisfying $x = \sum_{k \in \mathbb{N}^*} \varepsilon_k \beta^{-k}$ is called *expansion of x in base β* . The notion of expansion of x in base β is extended on \mathbb{R}_+ by considering sequences of $\mathcal{A}_\beta^{\mathbb{Z}}$. Among the expansions of x in base β , the greatest for the lexicographical order is called β -*expansion* of x . This expansion, denoted by $d_\beta(x) = 0.\varepsilon_1 \dots$, is constructed by the *greedy algorithm*, that is, $\varepsilon_k = \lfloor \beta T_\beta^{k-1}(x) \rfloor$ for all $k \in \mathbb{N}^*$, where $T_\beta : [0, 1[\rightarrow [0, 1[$, $y \mapsto \{\beta y\}$. Let d_β denote the map: $[0, 1] \rightarrow \mathcal{A}_\beta^{\mathbb{N}^*}$, $x \mapsto (\varepsilon_i)_{i \in \mathbb{N}^*}$. We recall that the map d_β is increasing if $\mathcal{A}_\beta^{\mathbb{N}^*}$ is endowed with the lexicographic order.

Parry showed in [31] that, for any $x \in [0, 1]$, $d_\beta(x)$ is the unique expansion of x in base β which satisfy the following relation called *Parry condition*:

$$\text{for all } n \in \mathbb{N}, S^n(v) <_{\text{lex}} d_\beta^*(1). \quad (2)$$

Any word or sequence $v \in \mathcal{A}_\beta^* \cup \mathcal{A}_\beta^{\mathbb{N}}$ which satisfy (2) is said to be *admissible*. Let \mathcal{L}_β denote the set of admissible words. The set of non-negative β -integers, denoted by \mathbb{Z}_β^+ , is defined as the set of non-negative real numbers x satisfying $x = \sum_{k=0}^n v_k \beta^k$, where $v_n \dots v_0 \in \mathcal{L}_\beta$. The set of β -integers is defined as $\mathbb{Z}_\beta = \pm \mathbb{Z}_\beta^+$. Since \mathbb{Z}_β^+ is a discrete set, one may define for $x \in \mathbb{R}_+$ the β -*integer part* of x as $[x]_\beta = \max_{p \in \mathbb{Z}_\beta^+} \{p \leq x\}$, and the β -*fractional part* of x as $\{x\}_\beta = x - [x]_\beta$. For $x > 1$, we set $d_\beta(x) = v.w$, where v and w respectively are the unique expansions of $[x]_\beta$ and $\{x\}_\beta$ in base β which satisfy the Parry condition.

For $x \in \mathbb{R}_+$, let $l_i(x)$ denote the length of the β -integer part of x , that is, $l_i(x) = N + 1$, where $N \in \mathbb{N}$ satisfies $\beta^N \leq x < \beta^{N+1}$ when $x \geq 1$, 0 otherwise. If there exists $M \in \mathbb{N}$ such that $\varepsilon_M \neq 0$ and $\varepsilon_n = 0$ for all $n > M$, we say that the β -expansion of x is finite. In this case, the ending consecutive occurrences of the letter 0 are omitted; $d_\beta(x) = \varepsilon_N \dots \varepsilon_0.\varepsilon_{-1} \dots \varepsilon_{-M}$, and we set $l_f(x) = M$, which denotes the length of the β -fractional part of x . The set of non-negative real numbers having a finite β -expansion is denoted by $\text{Fin}(\beta)^+$, and we set $\text{Fin}(\beta) = \pm \text{Fin}(\beta)^+$.

When $d_\beta(1)$ is either finite or ultimately periodic, β is said to be a *Parry number*. More precisely, β is said to be a *sofic Parry number* when $d_\beta(1)$ is ultimately periodic, and a *simple Parry number* when $d_\beta(1)$ is finite. The notion of Parry number was historically introduced by Parry [31] as β -*number*. Bertrand [12] and Schmidt [37] have independently proven that Pisot numbers are Parry numbers. When β is a simple Parry number, we denote by m the length of $d_\beta(1)$. In the sofic Parry case, we respectively denote by n and p the lengths of the preperiod and of the period of $d_\beta(1)$, and we set $m = n + p$ for convenience. More details about Parry numbers can be found in [26].

Remark 2. *One easily checks that $m \geq d$, where d is the algebraic degree of β . There exist Parry numbers such that $m > d$; for instance, the smallest Parry number is the positive root of the polynomial $X^3 - X - 1$, for which one has $d_\beta(1) = 0.10001$.*

Let β be an algebraic integer such that there exists a Galois conjugate α of β which satisfies $|\alpha| < 1$. Let $z \in \mathbb{C}$. We call α -*expansion* of z any admissible sequence $(\varepsilon_k)_{k \in \mathbb{Z}}$ such that $z = \sum_{k \in \mathbb{Z}} \varepsilon_k \alpha^{-k}$.

Remark 3. *There exist complex numbers having multiple α -expansions. For instance, consider the Fibonacci case, for which the Galois conjugate of β is $\alpha = \frac{1-\sqrt{5}}{2}$. Let the admissible sequences $(\varepsilon_k)_{k \in \mathbb{Z}}$ and $(\delta_k)_{k \in \mathbb{Z}}$ be defined by $\varepsilon_j = 1$ when $j \in 2\mathbb{N} + 1$, $\varepsilon_j = 0$ otherwise; $\delta_j = 1$ when $j \in \{-1\} \cup 2\mathbb{N}^*$, $\delta_j = 0$ otherwise. Then $-1 = \sum_{k \in \mathbb{Z}} \varepsilon_k \beta^{-k} = \sum_{k \in \mathbb{Z}} \delta_k \beta^{-k}$. Hence the real number -1 admits both $(\varepsilon_k)_{k \in \mathbb{Z}}$ and $(\delta_k)_{k \in \mathbb{Z}}$ as α -expansions.*

1.3 Notion of β -substitution

For any Parry number β , it is possible to define a substitution which is associated to β in a canonical way. Let β be a simple (resp. sofic) Parry number, with $d_\beta(1) = 0.k_1 \dots k_m$ (resp. $d_\beta(1) = 0.k_1 \dots k_n(k_{n+1} \dots k_{n+p})^\infty$). We remind that the elements of $(k_j)_{j \in \llbracket 1, \dots, m \rrbracket}$ are non-negative integers, and that for all $n \in \mathbb{N}^*$, $S^n(d_\beta(1)) <_{lex} d_\beta(1)$. The substitution σ_β is defined on $\{a_1, \dots, a_m\}$ by:

1. for all $j \in \llbracket 1 \dots m-1 \rrbracket$, $\sigma_\beta(a_j) = a_1^{k_j} a_{j+1}$,
2. $\sigma_\beta(a_m) = a_1^{k_m}$ in the simple Parry case, $\sigma_\beta(a_m) = a_1^{k_m} a_{n+1}$ otherwise.

The substitution associated in this way to a simple (sofic) Parry number β is called a *simple (sofic) β -substitution*, and is denoted by σ_β . Note that a β -substitution is primitive, admits a unique fixed point ω_β and is defined on an alphabet containing at least two letters. In particular, the set of the factors of ω_β defines a language denoted by \mathcal{L}_σ . See [16, 41], where these substitutions are introduced, for more details.

When β is a simple (sofic) Parry number, the incidence matrix M_σ of σ is

$$\begin{pmatrix} k_1 & \dots & \dots & \dots & \dots & k_m \\ 1 & 0 & \dots & \dots & \dots & 0 \\ 0 & \ddots & \ddots & & & \vdots \\ \vdots & \ddots & \ddots & \ddots & & 0 \\ \vdots & & \ddots & \ddots & \ddots & \vdots \\ 0 & \dots & \dots & 0 & 1 & 0 \end{pmatrix} \text{ (simple case) or } \begin{pmatrix} k_1 & \dots & \dots & \dots & \dots & k_{n+p} \\ 1 & 0 & \dots & \dots & \dots & 0 \\ 0 & \ddots & \ddots & & & \vdots \\ \vdots & \ddots & \ddots & \ddots & & 1 \\ \vdots & & \ddots & \ddots & \ddots & \vdots \\ 0 & \dots & \dots & 0 & 1 & 0 \end{pmatrix} \text{ (sofic case).}$$

Remark 4. Let us recall that, if σ_β is of Pisot type, then β is a Pisot number, but the converse is false. For example, the positive root β of the polynomial $X^3 - X - 1$ is a Pisot number, whereas σ_β is not of Pisot type, since the characteristic polynomial of the incidence matrix is $X^5 - X^4 - 1 = (X^3 - X - 1)(X^2 - X + 1)$, which admits two roots of modulus 1.

1.4 Rauzy fractals

Rauzy fractals were first introduced by Rauzy in [33, 34], then by Thurston in [41]. They provide a geometrical representation of the dynamical system defined by the associated β -substitution. In the framework of numeration, they are called *central tiles* by Akiyama [2, 3, 4].

Let β be a Pisot number of degree d , and $\{\alpha_j\}_{j \in \llbracket 1, \dots, r+s \rrbracket}$ be the set of the Galois conjugates which differ from β and which have a non-negative imaginary part. Let $\mu_\beta(X) = X^d - \sum_{j=1}^d b_j X^{d-j}$ denote the minimal polynomial of β on \mathbb{Z} , which admits $r+1$ real roots and $2s$ complex roots, hence $d-1 = r+2s$. For convenience, we set $J = \llbracket 1, \dots, r+s \rrbracket$.

Let M_β be the companion matrix of μ_β , defined as follows:

$$M_\beta = \begin{pmatrix} b_1 & \dots & \dots & \dots & b_d \\ 1 & 0 & \dots & \dots & 0 \\ 0 & \ddots & \ddots & & \vdots \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & \dots & 0 & 1 & 0 \end{pmatrix}.$$

For all $j \in J$, let \mathcal{H}_j denote the eigenspace associated to the eigenvalue α_j , and let \mathcal{D} denote the eigenspace associated to the dominant eigenvalue β . Note that $\mathcal{H}_j \simeq \mathbb{R}$ when $\alpha_j \in \mathbb{R}$, $\mathcal{H}_j \simeq \mathbb{C}$ otherwise. It is possible to describe \mathbb{R}^d as $\mathcal{D} \oplus \mathcal{H}$, where \mathcal{H} denotes the direct sum of the stable sub-spaces $\{\mathcal{H}_j\}_{j \in J}$. Moreover, there exists $\vec{w} \in \mathcal{D}$ and a base of $\mathcal{H} = \bigoplus_{j \in J} \mathcal{H}_j$ constituted by $r+2s$ complex eigenvectors $(\vec{v}_j)_{j \in \llbracket 1, \dots, r+2s \rrbracket}$, which can be chosen such that the following equalities hold:

$$\vec{e}_1 = \sum_{j \in J} \vec{v}_j + \vec{w},$$

$$\forall j \in \llbracket 1, \dots, s \rrbracket, \forall i \in \llbracket 1, \dots, d \rrbracket, v_{r+j}[i] = \overline{v_{r+s+j}[i]},$$

where $v_k[i]$ denotes the i -th coordinate of $v_k \in \mathbb{R}^r \times \mathbb{C}^{d-r}$ and $z \mapsto \bar{z}$ the standard conjugacy on \mathbb{C} . These relations allow us to define $\overrightarrow{p_{\mathcal{D}}}$ and, for all $j \in J$, $\overrightarrow{p_{\mathcal{H}_j}}$, which are the projections on each associated subspace along the direct sum of the others. We then set $p_{\mathcal{D}}$ and, for all $j \in J$, $p_{\mathcal{H}_j}$ as the coordinate maps in the base $(\overrightarrow{w}, \overrightarrow{v}_1, \dots, \overrightarrow{v_{r+2s}})$ of the associated projection maps, that is, one gets for all $\overrightarrow{X} \in \mathbb{R}^d$:

$$\overrightarrow{p_{\mathcal{H}_j}}(\overrightarrow{X}) = p_{\mathcal{H}_j}(\overrightarrow{X}) \overrightarrow{v}_j, \text{ for all } j \in \llbracket 1, \dots, r \rrbracket,$$

$$\overrightarrow{p_{\mathcal{H}_j}}(\overrightarrow{X}) = p_{\mathcal{H}_j}(\overrightarrow{X}) \operatorname{Re}(\overrightarrow{v}_j), \text{ for all } j \in \llbracket r+1, \dots, r+s \rrbracket.$$

For all $j \in J$, we define τ_j as the field morphism: $\mathbb{Q}(\beta) \rightarrow \mathbb{Q}(\alpha_j)$, $\beta \mapsto \alpha_j$. We denote by τ the map:

$$\tau : \mathbb{Q}(\beta) \rightarrow \mathbb{R}^r \times \mathbb{C}^s \simeq \mathbb{R}^{d-1}, x \mapsto (\tau_1(x), \dots, \tau_{r+s}(x)).$$

Definition 1.1. The set $\{\tau(x), x \in \mathbb{Z}_{\beta}^+\}$ is called Rauzy fractal or central tile, denoted by \mathcal{T} .

Note that a Rauzy fractal is a compact set by definition, since it is a bounded set.




Fig. 1: Rauzy fractal \mathcal{T} defined by $d_{\beta}(1) = 0.111$

Example 1. Figure 1 shows the Rauzy fractal associated to the Tribonacci case, defined by $d_{\beta}(1) = 0.111$.

2 Rauzy fractals and symmetric sets

In this section, we introduce a geometric representation set \mathcal{T}' for the numeration defined by β , a given algebraic number. The set \mathcal{T}' , which admits a symmetric property that we explain later, contains the Rauzy fractal \mathcal{T} . Then, following [9], the study of \mathcal{T}' provides inequalities satisfied by the images under τ_j of the finite β -fractional parts of sums of two β -integers. This allows us to define an algorithm which improves the determination of an upper bound and of a lower bound for L_{\oplus} .

2.1 Set of β -integers and admissibility

The set of non-negative β -integers is defined as

$$\mathbb{Z}_{\beta}^+ = \bigcup_{n \in \mathbb{N}} \left\{ \sum_{k=0}^n v_k \beta^k, v_n \dots v_0 \in \mathcal{L}_{\beta} \right\}.$$

When one performs arithmetics on β -integers, the computation of the admissibility condition is not so easy, and the renormalization process significantly slows down any algorithmic process. If we remove the admissibility condition, we obtain a set that contains \mathbb{Z}_{β}^+ , which is

$$\Sigma_{\beta}^+ = \bigcup_{n \in \mathbb{N}} \left\{ \sum_{k=0}^n v_k \beta^k, v_n \dots v_0 \in \mathcal{A}_{\beta}^{n+1} \right\}.$$

Note that, for any $x \in \mathbb{Z}_{\beta}^+$, there exists a unique admissible word $v_n \dots v_0 \in \mathcal{L}_{\beta}$ such that $x = \sum_{k=0}^n v_k \beta^k$,

whereas there may exist $x \in \Sigma_{\beta}$, $n \in \mathbb{N}$ and two distinct words $v, w \in \mathcal{A}_{\beta}^*$ such that $x = \sum_{k=0}^n v_k \beta^k =$

$\sum_{k=0}^n w_k \beta^k$. As an example, the words 100 and 011 represent the same real number in Fibonacci base, since $\beta^2 = \beta + 1$. The word 011 is defined on \mathcal{A}_β^* , but it does not belong to \mathcal{L}_β . Hence, if β is such that $\mathbb{Z}_\beta^+ = \Sigma_\beta^+$, we can omit the admissibility condition in order to generate the set of β -integers.

The set of Parry numbers for which $\mathbb{Z}_\beta^+ = \Sigma_\beta^+$ is exactly the set of numbers which define confluent numeration systems, which were introduced and studied by Frougny [17]. Confluent numeration systems characterize numeration systems for which any finite expansion on the alphabet \mathcal{A}_β can be renormalized as an admissible word which represents a β -integer. Therefore confluent Parry numbers, that is, Parry numbers which define a confluent numeration system, are the numbers for which there is no carry propagating to the right. We will use later the following proposition, which gathers different results from [19] and [17].

Proposition 2.1. [19, 17] *Let β be a Parry number. The following assertions are equivalent.*

1. *One has $d_\beta(1) = 0.k_1^{d-1}k_d$, with $k_1, k_d \in \mathbb{N}$ and $k_1 \geq k_d$,*
2. *the language \mathcal{L}_σ is stable under mirror image,*
3. *there exist infinitely many prefixes of ω , the fixed point of σ_β , that are palindromes,*
4. *one has $\mathbb{Z}_\beta^+ = \Sigma_\beta^+$.*

When any of these assertions holds, β is said to be a confluent Parry number.

It is proved in [20] that confluent Parry numbers are Pisot numbers such that $\text{Fin}(\beta) = \mathbb{Z}[\beta^{-1}]$, that is, the finiteness property holds true. Moreover, the class of confluent Parry numbers defines a class of β -substitutions whose associated dynamical system has purely discrete spectrum [40], and the complexity of their substitutive language is affine [19].

2.2 The symmetric property for confluent Parry numbers

We focus on a geometric property for the Rauzy fractal \mathcal{T} defined by a Pisot number β , indeed, we see in this section that $\overline{\tau(\Sigma_\beta^+)}$ is stable under a central symmetry map defined below. Note that, since $\mathbb{Z}_\beta^+ \subset \Sigma_\beta^+$, we get $\mathcal{T} \subset \overline{\tau(\Sigma_\beta^+)}$. Moreover, according to Proposition 2.1, the set of confluent Parry number characterizes the Parry numbers for which $\mathcal{T} = \overline{\tau(\Sigma_\beta^+)}$.

We will see in Section 3.1 that the relations developed here enable a covering of \mathcal{T} with balls having arbitrarily small positive radius, in a more accurate way than in [9]. In particular, this covering is accurate enough to allow the computation, in several cases of cubic Pisot numbers, of the maximal length of the finite β -fractional parts for the sums of two β -integers.

Definition 2.2. *Let $d \in \mathbb{N}^*$, and let \mathcal{E} be a bounded set of $\mathbb{R}^r \times \mathbb{C}^s \simeq \mathbb{R}^{d-1}$. The set \mathcal{E} is said to be centrally symmetric if there exists $\vec{C} \in \mathbb{R}^r \times \mathbb{C}^s$ such that \mathcal{E} is stable under the symmetry $S_{\vec{C}} : \mathbb{R}^r \times \mathbb{C}^s \rightarrow \mathbb{R}^r \times \mathbb{C}^s$, $\vec{X} \mapsto 2\vec{C} - \vec{X}$. In this case, \vec{C} is called center of the set \mathcal{E} .*

Lemma 2.3. *Let β be a Pisot number. Then $\overline{\tau(\Sigma_\beta^+)}$ is a centrally symmetric set, whose center \vec{C} satisfies $p_{\mathcal{H}_j}(\vec{C}) = \frac{[\beta]}{2(1-\alpha_j)}$, for all $j \in J$.*

Proof. First, note that $\tau(\Sigma_\beta^+)$ is a bounded set of $\mathbb{R}^r \times \mathbb{C}^s \simeq \mathbb{R}^{d-1}$, and that $p_{\mathcal{H}_j}(\overline{\tau(\Sigma_\beta^+)}) = \overline{\tau_j(\Sigma_\beta^+)}$. Let $\vec{X} \in \tau(\Sigma_\beta^+)$. There exists a sequence $(\varepsilon_n)_{n \in \mathbb{N}} \in \mathcal{A}_\beta^{\mathbb{N}}$ such that $p_{\mathcal{H}_j}(\vec{X}) = \sum_{n \in \mathbb{N}} \varepsilon_n \alpha_j^n$ for all $j \in J$. Let $(\delta_n)_{n \in \mathbb{N}} \in \mathcal{A}_\beta^{\mathbb{N}}$ be defined by $\delta_n = [\beta] - \varepsilon_n$ for all $n \in \mathbb{N}$. We define $\vec{Y} \in \mathbb{R}^r \times \mathbb{C}^s$ by $p_{\mathcal{H}_j}(\vec{Y}) = \sum_{n \in \mathbb{N}} \delta_n \alpha_j^n$ for all $j \in J$. Then $p_{\mathcal{H}_j}(\vec{Y}) = \frac{[\beta]}{1-\alpha_j} - p_{\mathcal{H}_j}(\vec{X})$ for all $j \in J$. Moreover, \vec{Y} can be described as the limit of a sequence $(\vec{Y}_n)_{n \in \mathbb{N}} \in \tau(\Sigma_\beta^+)$, whose elements are defined by $p_{\mathcal{H}_j}(\vec{Y}_n) = \sum_{k=0}^n \delta_k \alpha_j^k$ for all $j \in J$. Since $\tau(\Sigma_\beta^+)$ is a compact set, we deduce that $\vec{Y} \in \mathcal{T}$. Hence, if we define $\vec{C} \in \mathbb{R}^r \times \mathbb{C}^s$ by $p_{\mathcal{H}_j}(\vec{C}) = \frac{[\beta]}{2(1-\alpha_j)}$ for all $j \in J$, we obtain that $\overline{\tau(\Sigma_\beta^+)}$ is stable under the symmetry of center \vec{C} , which ends the proof. \square

Definition 2.4. The symmetrized tile is defined as $\mathcal{T}' = \overline{\tau(\Sigma_{\beta}^+)}$.

One directly deduces from Proposition 2.1 and Lemma 2.3 the following:

Proposition 2.5. Let β be a confluent Parry number. Then \mathcal{T} is a centrally symmetric set.

Remark 5. When β is the sofic Parry number defined by the polynomial $X^2 - (k_1 + 1)X + k_1 - k_2$ with $k_1 > k_2 > 0$, one has $d_{\beta}(1) = 0.k_1k_2^{\infty}$, $\mathcal{T} = [0, \frac{k_1\beta^2}{\beta^2 - k_2^2}]$ and $\mathcal{T}' = [0, \frac{k_1\beta}{\beta - k_2}]$. Hence \mathcal{T} is a centrally symmetric set with $\mathcal{T} \subsetneq \mathcal{T}'$. We do not know whether there exist others Parry numbers for which \mathcal{T} is centrally symmetric with $\mathcal{T} \subsetneq \mathcal{T}'$.

The following proposition characterizes centrally symmetrical Rauzy fractals that contain their center \vec{C} . Since there exist Rauzy fractals that are not connected, it is not clear whether $\vec{C} \in \mathcal{T}$ for any confluent Parry number.

Proposition 2.6. Let β be a confluent Parry number. If either k_1 or $k_1 - k_d$ is even, then $\vec{C} \in \mathcal{T}$.

Proof. Let β be a confluent Parry number. We consider the two cases k_1 even and k_1 odd. When k_1 is even, $\frac{k_1}{2} \in \mathcal{A}_{\beta}$ and $(\sum_{i=0}^n \frac{k_1}{2}\beta^i)_{n \in \mathbb{N}}$ is a sequence of β -integers. For any $j \in J$, the image under τ_j of the elements of this sequence is $(\sum_{i=0}^n \frac{k_1}{2}\alpha_j^i)_{n \in \mathbb{N}}$, which tends to $\frac{k_1}{2} \frac{1}{1-\alpha_j}$. Since $k_1 = [\beta]$, we get $\frac{k_1}{2} \frac{1}{1-\alpha_j} = p_{\mathcal{H}_j}(\vec{C})$ according to Lemma 2.3. Since \mathcal{T} is closed, we deduce that $\vec{C} \in \mathcal{T}$.

Suppose now that k_1 is odd, with β root of $X^d - k_1 \sum_{i=1}^{d-1} X^i - k_d$. If $k_1 - k_d$ is even, we have $\sum_{i=0}^{d-1} \frac{k_1}{2}\beta^i = \frac{1}{2}\beta^d + \frac{k_1 - k_d}{2}$. One has $\sum_{i \in \mathbb{N}} \frac{k_1}{2}\alpha_j^i = \sum_{l \in \mathbb{N}} (\sum_{i=0}^d \frac{k_1}{2}\alpha_j^{i+l(d+1)}) = \sum_{l \in \mathbb{N}} (\frac{k_1+1}{2}\alpha_j^d + \frac{k_1 - k_d}{2})\alpha_j^{l(d+1)}$. Let $(\vec{C}_n)_{n \in \mathbb{N}} \in (\mathbb{R}^{d-1})^{\mathbb{N}}$ be defined by $p_{\mathcal{H}_j}(\vec{C}_n) = \sum_{l=0}^n (\frac{k_1+1}{2}\alpha_j^d + \frac{k_1 - k_d}{2})\alpha_j^{l(d+1)}$ for all $j \in J$ and for all $n \in \mathbb{N}$. Since $k_1 - k_d$ and $k_1 + 1$ are even, with $k_1 \geq k_d \geq 1$, $\frac{k_1+1}{2}$ and $\frac{k_1 - k_d}{2} \in \mathcal{A}_{\beta}$. Additionally, the word $(\frac{k_1+1}{2}0^{d-1}\frac{k_1 - k_d}{2})^n$ is admissible for all $n \in \mathbb{N}$. This means that the sequence of β -integers $(p_n)_{n \in \mathbb{N}}$, defined by $p_n = \sum_{l=0}^n (\frac{k_1+1}{2}\beta^d + \frac{k_1 - k_d}{2})\beta^{l(d+1)}$, satisfies $\tau(p_n) = \vec{C}_n$ for all $n \in \mathbb{N}$, hence $\vec{C}_n \in \mathcal{T}$ for all $n \in \mathbb{N}$. Using again Lemma 2.3 and the compactness of \mathcal{T} , we deduce that $\vec{C} \in \mathcal{T}$. \square

Remark 6. Since $p_{\mathcal{H}_j}(\vec{C}) = \frac{[\beta]}{2(1-\alpha_j)}$ for any $j \in J$, $\vec{C} \neq \vec{0}$.

By the way, let us give a geometric interpretation of Proposition 2.5. Assume that β is a confluent Parry number, with $d_{\beta}(1) = 0.k_1^{d-1}k_d$. According to Proposition 2.1, the associated substitutive language \mathcal{L}_{σ} defined by the substitution σ_{β} is stable under the mirror image map. Moreover, there exist infinitely many palindromic prefixes for ω , the fixed point of σ . Indeed, let $(\omega_{v_n})_{n \in \mathbb{N}}$ be the sequence defined by $\omega_{v_n} = h^n(a_1)$ for all $n \in \mathbb{N}$, where the map h is defined by

$$h : \mathcal{L}_{\sigma} \longrightarrow \mathcal{L}_{\sigma}, v \longmapsto \sigma_{\beta}(v)a_1^{k_1}.$$

One checks that h preserves the set of palindromes of \mathcal{L}_{σ} , which implies that $(\omega_{v_n})_{n \in \mathbb{N}}$ is a sequence of palindromic prefixes of ω . Let $v \in \mathcal{A}$ and $x = p_{\mathcal{D}}(f(v))$, where f denotes the abelianization map. We remind that, by definition of the map $p_{\mathcal{D}}$ defined in Section 1.4, one has $p_{\mathcal{D}}(f(a_1)) = 1$ and $p_{\mathcal{D}}(\sigma(w)) = \beta.p_{\mathcal{D}}(f(w))$ for all $w \in \mathcal{L}_{\sigma}$. Thus, one has $p_{\mathcal{D}}(f(h(v))) = \beta x + k_1$, hence for all $n \in \mathbb{N}$, $p_{\mathcal{D}}(f(h^n(a_1))) = g^n(1)$, where $g : \mathbb{R} \rightarrow \mathbb{R}$ is the affine map defined by $g(x) = \beta x + k_1$.

Since ω_{v_n} is a palindrome, the set $E_n = \{f(\omega_k), k \in [0, \dots, v_n]\}$ is a centrally symmetrical set of \mathbb{Z}^d . Hence $p_{\mathcal{D}}(E_n)$ and $p_{\mathcal{H}}(E_n)$ respectively are centrally symmetrical sets of \mathbb{R} and \mathbb{R}^{d-1} . Let \vec{C}_n denote the center of $p_{\mathcal{H}}(E_n)$. Then for all $j \in J$, $p_{\mathcal{H}_j}(\vec{C}_n) = \frac{g_j^{v_n}(1)}{2}$, where g_j is the affine map: $x \longmapsto \alpha_j x + k_1$. Let c_j be the fixed point of g_j for all $j \in J$. Since β is a Pisot number, g_j is a contracting map for all $j \in J$, hence the sequence $(p_{\mathcal{H}_j}(\vec{C}_n))_{n \in \mathbb{N}}$ tends to $\frac{c_j}{2}$. Let \vec{C} be the limit of the sequence $(\vec{C}_n)_{n \in \mathbb{N}}$.

Since $p_{\mathcal{H}_j}(\vec{C}) = \frac{c_j}{2}$, and since $k_1 = [\beta]$, the center of \mathcal{T} satisfies $p_{\mathcal{H}_j}(\vec{C}) = \frac{[\beta]}{2(1-\alpha_j)}$ for all $j \in J$. We retrieve the result of Lemma 2.3, since $\mathcal{T} = \mathcal{T}'$. Note that, rather than the foregoing construction of \mathcal{T} as the limit of centrally symmetric sets, constructed by considering prefixes of the fixed point ω of σ_β , it is also possible to describe \mathcal{T} as the image of the closure of $p_{\mathcal{H}}(f(\omega' \cdot \omega'))$ under a translation map, where $\omega' \cdot \omega'$ is a two-sided palindrome whose factors belong to \mathcal{L}_σ . This construction, and the previous one, cannot be completed when β is a Parry number which is not confluent, since \mathcal{L}_σ contains only finitely many palindromes in this case.

There is no reason for the tile \mathcal{T} to be a centrally symmetric set when β is a Parry number which is not confluent. See for instance Figure 2 and Figure 3, which respectively represent, for the numeration system defined by $d_\beta(1) = 0.101$, first, the Rauzy fractal \mathcal{T} , then the sets $\mathcal{T} \cup S_{\vec{C}}(\mathcal{T})$ and \mathcal{T}' with different levels of shading.




Fig. 2: \mathcal{T} defined by $d_\beta(1) = 0.101$




Fig. 3: \mathcal{T}' defined by $d_\beta(1) = 0.101$

3 Computation of L_\oplus in several cases of cubic Pisot numbers

In this section, we are interested in computing L_\oplus , or at least computing a better upper bound for L_\oplus than those determined in [9]. First, we focus in 3.1 on several algebraic inequalities, provided by Propositions 3.4 and 3.5, that are satisfied by the image under τ of the finite β -fractional parts that are obtained as the β -fractional parts of the sum of two β -integers. Whereas these relations do not allow us to compute directly the set of these finite β -fractional parts, we can discard, thanks to an algorithm defined in Section 3.2, the β -fractional parts which do not satisfy these inequalities. In particular, this allows us to compute L_\oplus for several cases of Pisot units, since, among the remaining β -fractional parts that were not discarded by the algorithm A , we find an element of $\text{Fin}(\beta)$, whose β -fractional part is of length L_\oplus , which is the sum of two β -integers.

3.1 Fractional parts of sums or differences of β -integers

Let β be a Pisot number, and let $(w_i)_{i \in \mathbb{N}} \in \mathcal{A}_\beta^\mathbb{N}$. Let $(v_i)_{i \in \mathbb{N}}$ be defined for all $i \in \mathbb{N}$ by $v_i = [\beta] - w_i$. Since $(v_i)_{i \in \mathbb{N}} \in \mathcal{A}_\beta^\mathbb{N}$, $\mathcal{T}' = \overline{\tau(\Sigma_\beta^+)}$ contains both \mathcal{T} and $S_{\vec{C}}(\mathcal{T}) = \bigcup_{\vec{X} \in \mathcal{T}} \{2\vec{C} - \vec{X}\}$, where \vec{C} is the center of \mathcal{T}' . We deduce from $\mathcal{T} \subset \mathcal{T}'$ that inequalities satisfied by elements of \mathcal{T}' are satisfied by elements of \mathcal{T} as well. Note that there is no reason that $\mathcal{T} \cup S_{\vec{C}}(\mathcal{T}) = \mathcal{T}'$, see Figure 3.

First, let us recall some results of [9], where a method is described in order to compute upper bounds for L_\oplus and L_\ominus . Let α_j denote a Galois conjugate of β such that $|\alpha_j| < 1$. We define for all $n \in \mathbb{N}^*$ the real numbers

$$M_n(j) = \max_{x \in \mathbb{Z}_\beta^+} \{|\tau_j(x)|, x < \beta^n\}, H_n(j) = \frac{M_n(j)}{1 - |\alpha_j|^n} \text{ and } H(j) = \sup_{x \in \mathbb{Z}_\beta^+} |\tau_j(x)|.$$

Let us give a geometric interpretation for $H_n(j)$. Indeed, $H_n(j)$ is the upper bound of the modulus of complex numbers $z \in \mathcal{T}$ such that, for all $j \in J$, $p_{\mathcal{H}_j}(z)$ admits a periodic α_j -expansion of period n . Hence, for all $j \in J$, $H_l(j) \geq H_k(j)$ for all $k, l \in \mathbb{N}$ such that $l|k$, and $H_k(j) \geq H(j)$ for all $k \in \mathbb{N}^*$.

The following inequality, used for instance in the proof of Proposition 3.2, holds.

Proposition 3.1. For all $x \in \mathbb{Z}_{\beta}$ and $n \in \mathbb{N}^*$, $|\tau_j(x)| < H_n(j)$.

Proposition 3.1 allows us to obtain the following result, which provides upper bounds for L_{\oplus} and L_{\odot} .

Proposition 3.2. [9] Let $\beta > 1$ be an algebraic integer. If there exists a Galois conjugate α_j of β such that $K(j) = \inf_{z \in \mathbb{Z}_{\beta}^+ \setminus \beta \mathbb{Z}_{\beta}^+} |\tau_j(z)| > 0$, then

$$L_{\oplus} < \frac{\ln 2 + \ln H(j) - \ln K(j)}{-\ln |\alpha_j|} \text{ and } L_{\odot} < \frac{2 \ln H(j) - \ln K(j)}{-\ln |\alpha_j|}.$$

In particular, assume that $d_{\beta}(1) = 0.k_1k_2k_3$, and that β has two complex Galois conjugates. Then β is a Pisot cubic number; the finiteness property holds [3], which implies due to [2] that 0 is an inner point of \mathcal{T} . Hence $K(j) > 0$, and the required conditions of Proposition 3.2 are satisfied.

Proposition 3.2 is a consequence of Proposition 3.1. Indeed, let x and $y \in \mathbb{Z}_{\beta}$ with $x + y \in \text{Fin}(\beta)$. Since $l_f(x + y) \leq L_{\oplus}$, one gets $|\tau_j(x)| + |\tau_j(y)| < 2H(j)$ and $|\tau_j(x + y)| > |\alpha_j|^{L_{\oplus}} K(j)$. Hence the upper bound for L_{\oplus} , and the case of multiplication is performed in a similar way. We focus now on the search of better inequalities satisfied by elements of $\text{Fin}(\beta)$, since this will allow us to deduce better upper bounds for L_{\oplus} and L_{\odot} . Inequalities improving Proposition 3.1 are detailed below in Propositions 3.4 and 3.5.

Since $\mathbb{Z}_{\beta}^+ \subset \Sigma_{\beta}^+$, any inequality satisfied by elements of $\tau(\Sigma_{\beta}^+)$ is satisfied by $\tau(\mathbb{Z}_{\beta}^+)$ as well. Hence the study of $\tau(\Sigma_{\beta}^+)$ will provide additional inequalities satisfied by elements of \mathcal{T} . First, we define for all $n \in \mathbb{N}^*$ the real numbers

$$M'_n(j) = \max_{x \in \Sigma_{\beta}^+} \{|\tau_j(x)|, x < \beta^n\}, H'_n(j) = \frac{M'_n(j)}{1 - |\alpha_j|^n} \text{ and } H'(j) = \sup_{x \in \Sigma_{\beta}^+} |\tau_j(x)|.$$

Similarly as in [9], we obtain using the triangle inequality the following property.

Lemma 3.3. For all $x \in \mathbb{Z}_{\beta}$, $|\tau_j(x)| < H'_n(j)$ and $|\tau_j(x) - 2p_{\mathcal{H}_j}(\vec{C})| < H'_n(j)$.

Proof. The first assertion is deduced from Proposition 3.1, whereas the second assertion is a consequence of $S_{\vec{C}}(\mathcal{T}) \subset \mathcal{T}'$. \square

Note that, since $H'(j) \geq H(j)$, the first inequality of Lemma 3.3 is a weaker condition than the inequality of Proposition 3.1. However, it is more convenient to compute $H'(j)$, since the definitions of $H'_n(j)$ and $H'(j)$ do not use the admissibility condition on words.

We obtain additional algebraic inequalities, satisfied by finite β -fractional parts of the sums of two β -integers, thanks to the following proposition.

Proposition 3.4. Let $x, y \in \mathbb{Z}_{\beta}^+$ with $x \geq y$. If $x + y \in \text{Fin}(\beta)$, then:

1. $|\tau_j(\{x + y\}_{\beta}) - 2p_{\mathcal{H}_j}(\vec{C})| < 3H'(j)$;
2. $|\tau_j(\{x + y\}_{\beta})| < (2 + |\alpha_j|)H(j)$.

If $x - y \in \text{Fin}(\beta)$, then:

1. $|\tau_j(\{x - y\}_{\beta}) + 2p_{\mathcal{H}_j}(\vec{C})| < 3H'(j)$;
2. $|\tau_j(\{x - y\}_{\beta})| < (2 + |\alpha_j|)H(j)$.

Proof. First, let us consider the case of addition of positive β -integers. Let $x, y \in \mathbb{Z}_{\beta}^+$ such that $x + y \in \text{Fin}(\beta)$. The first assertion is a consequence of Lemma 3.3, using the fact that, since x, y and $[x + y]_{\beta}$ are β -integers, we get $|\tau_j(x) - 2p_{\mathcal{H}_j}(\vec{C})| < H'(j)$, $|\tau_j(y) - 2p_{\mathcal{H}_j}(\vec{C})| < H'(j)$ and $|\tau_j([x + y]_{\beta}) - 2p_{\mathcal{H}_j}(\vec{C})| < H'(j)$. Let us prove the second assertion.

If 0 is a suffix of the expansion of either x, y , or $[x + y]_{\beta}$, then one of these three β -integers, say x , satisfies $|\tau_j(x)| < |\alpha_j|H(j)$, which improves the inequality of the first assertion. On the other hand, suppose that

none of the β -expansions of x , y and $[x+y]_\beta$ admits 0 as a suffix. Denote respectively by x_0 , y_0 and z_0 the suffixes of x , y and $[x+y]_\beta$. Then, one of the following non-negative β -integers $x' = x - \min\{x_0, y_0\}$, $y' = y - \min\{x_0, y_0\}$ or $z' = y - \min\{x_0, y_0\}$ admits 0 as a suffix, and both $x' + y = z' + \{x+y\}_\beta$ and $x + y' = z' + \{x+y\}_\beta$ hold. Among the β -integers x, x', y, y' , one of them belongs to $\beta\mathbb{Z}_\beta^+$. This proves that there exist $x', y' \in \mathbb{Z}_\beta^+$, with either x' or $y' \in \beta\mathbb{Z}_\beta^+$, such that $\{x+y\}_\beta = \{x'+y'\}_\beta$, hence $|\tau_j(\{x+y\}_\beta)| < (2 + |\alpha_j|)H(j)$.

The case of the subtraction of two positive β -integers is handled in the same way. \square

These relations improve those found in [9]. We obtain Proposition 3.5 in a similar way, which enables the computation of L_\oplus in Section 3.2 for several cases of cubic Pisot units.

Remark 7. Proposition 3.4 means that $3H'(j)$ and $(2 + |\alpha_j|)H(j)$ are both upper bounds for $|\tau_j(\{x+y\}_\beta)|$ when x and y are β -integers such that $[x+y]_\beta \in \text{Fin}(\beta)$. When β is a confluent Parry number, $H(j) = H'(j)$, which means that $(2 + |\alpha_j|)H(j)$ provides a better upper bound for $|\tau_j(\{x+y\}_\beta)|$ than $3H'(j)$. However, when β is the positive root of the polynomial $X^3 - X^2 - 1$, we check by pure computation that $3H'(j) < (2 + |\alpha_j|)H(j)$. Hence, in the general case of Pisot numbers, we do not know which of the two upper bounds $3H'(j)$ and $(2 + |\alpha_j|)H(j)$ for $|\tau_j(\{x+y\}_\beta)|$ is better than the other one.

The following proposition is an improvement of Proposition 3.4, in the sense that it sums up the inequalities concerning the elements of \mathcal{T} . In particular, we obtain as a result an accurate covering of \mathcal{T} by balls having arbitrarily small positive radius. As a consequence, we construct the algorithm A described in Section 3.2, which computes a set of admissible words whose β -fractional parts are of a bounded length. This set contains the finite β -fractional parts of sums of two positive β -integers.

Proposition 3.5. Let $x, y \in \mathbb{Z}_\beta^+$ with $x > y$, $x = \sum_{k \in \mathbb{N}} \varepsilon_k \beta^k$ and $y = \sum_{k \in \mathbb{N}} \delta_k \beta^k$. Suppose that $x+y \in \text{Fin}(\beta)$. Let $(\nu_k)_{k \in \mathbb{Z}}$ be the β -expansion of $x+y$. Then, for all $N \in \mathbb{N}^*$:

$$\left| \sum_{k < 0} \nu_k \alpha_j^k + \sum_{k=0}^{N-1} (\nu_k - \varepsilon_k - \delta_k) \alpha_j^k - \alpha_j^N p_{\mathcal{H}_j}(\vec{C}) \right| < \frac{3H'(j)(1 + |\alpha_j|)}{2} |\alpha_j|^N.$$

Proof. First, we know due to Lemma 2.3 that \vec{C} is the center of the centrally symmetric set \mathcal{T}' , with $p_{\mathcal{H}_j}(\vec{C}) = \frac{[\beta]}{2(1-\alpha_j)} = \sum_{k \geq 0} \frac{[\beta]}{2} \alpha_j^k$ for all $j \in J$.

Let $x, y \in \mathbb{Z}_\beta^+$ such that $\{x+y\}_\beta \in \text{Fin}(\beta)$. Let $(\varepsilon_k)_{k \in \mathbb{Z}}$, $(\delta_k)_{k \in \mathbb{Z}}$ and $(\nu_k)_{k \in \mathbb{Z}}$ respectively be the β -expansion of x , y and $x+y$. We remind that, by hypothesis, $\varepsilon_k = \delta_k = 0$ for all $k < 0$, and that $(\varepsilon_k - \delta_k - \nu_k)_{k \in \mathbb{Z}}$ consists of only finitely many non-zeros elements. Since $\{x+y\}_\beta + [x+y]_\beta - x - y = 0$, one gets for all $N \in \mathbb{N}^*$:

$$\begin{aligned} \sum_{k < 0} \nu_k \alpha_j^k + \sum_{k=0}^{N-1} (\nu_k - \varepsilon_k - \delta_k) \alpha_j^k &= - \sum_{k \geq N} (\nu_k - \varepsilon_k - \delta_k) \alpha_j^k \\ &= - \alpha_j^N \sum_{k \in \mathbb{N}} (\nu_{k+N} - \varepsilon_{k+N} - \delta_{k+N}) \alpha_j^k. \end{aligned}$$

Since for all $j \in J$, one has $p_{\mathcal{H}_j}(\vec{C}) = \sum_{k \in \mathbb{N}} \frac{[\beta]}{2} \alpha_j^k = \frac{[\beta]}{2(1-\alpha_j)}$, we obtain:

$$\sum_{k \geq N} (\nu_k - \varepsilon_k - \delta_k) \alpha_j^k = \alpha_j^N \left(\sum_{k \in \mathbb{N}} (\nu_{k+N} - \varepsilon_{k+N} - \delta_{k+N} + \frac{[\beta]}{2}) \alpha_j^k - p_{\mathcal{H}_j}(\vec{C}) \right).$$

Moreover, $\nu_{k+N} - \varepsilon_{k+N} - \delta_{k+N} + \frac{[\beta]}{2} \in [-3\frac{[\beta]}{2}, 3\frac{[\beta]}{2}]$ for all $k \in \mathbb{N}$. Hence

$$\left| \sum_{k < 0} \nu_k \alpha_j^k + \sum_{k=0}^{N-1} (\nu_k - \varepsilon_k - \delta_k) \alpha_j^k - \alpha_j^N p_{\mathcal{H}_j}(\vec{C}) \right| < \frac{3}{2} |\alpha_j|^N R,$$

where $R = \sup\{|\sum_{k \in \mathbb{N}} v_k \alpha_j^k|, (v_k)_{k \in \mathbb{N}} \in [-[\beta], [\beta]]^{\mathbb{N}}\}$.

Since $\overline{\{\sum_{k \in \mathbb{N}} v_k \alpha_j^k, (v_k)_{k \in \mathbb{N}} \in [-[\beta], [\beta]]^{\mathbb{N}}\}}$ is a convex set, and using a similar argument as in the proof of Proposition 3.4, we get:

$$R = \max_{x', y' \in \Sigma_{\beta}^+} |\tau_j(x' - y')| \leq (1 + |\alpha_j|)H'(j).$$

□

Note that we do not use the admissibility in the proof of Propositions 3.4 and 3.5. This means that, whereas we are looking for inequalities concerning elements which belong to \mathcal{T} , these inequalities hold in fact for elements of \mathcal{T}' .

The relation given by Proposition 3.5 produces a good approximation of \mathcal{T}' , in the sense that it enables an accurate covering of \mathcal{T}' with balls having arbitrarily small positive radius. More precisely, the coordinates on \mathcal{H}_j of the centers of these balls are $\sum_{k=0}^{N-1} (\nu_k - \varepsilon_k - \delta_k) \alpha_j^k - p_{\mathcal{H}_j}(\vec{C})$ when ν, ε and δ describe the set of admissible words of length N , and their radii are $\frac{3}{2} |\alpha_j|^N R$.

It could be possible that the set \mathcal{T} is only a “small” subset of \mathcal{T}' . In this case, the inequality satisfied by elements of \mathcal{T} in Proposition 3.5 is not accurate enough, and we need to improve the relation satisfied by elements of \mathcal{T} in order to compute the exact value of L_{\oplus} , or at least a better upper bound for L_{\oplus} than those given by Proposition 3.2. Note that the class of confluent Parry numbers is in some sense optimal, since $\mathcal{T} = \mathcal{T}'$ corresponds to the case where a good approximation of \mathcal{T}' is a good approximation of \mathcal{T} as well.

3.2 Algorithmic computation of L_{\oplus} in several cubic Pisot cases

In this section, we are interested in the set of finite β -fractional parts that are obtained when one sums two positive β -integers. We introduce L_{\oplus}^+ as the maximal length of the finite β -fractional parts of the sum of two positive β -integers. Note that L_{\oplus} , the maximal length of the finite β -fractional parts of the sum of two β -integers, can also be defined as the maximal length of the finite β -fractional parts that we obtain after adding or subtracting two positive β -integers. Hence, if one can both compute L_{\oplus}^+ and the maximal length of the finite β -fractional parts of the difference of two positive β -integers, we deduce the value of L_{\oplus} .

Let \mathcal{L}'_{β} be the set which consists of words $(\nu_k - \varepsilon_k - \delta_k)_{k \in \llbracket 1, \dots, n \rrbracket}$, where the words $(\nu_k)_{k \in \llbracket 1, \dots, n \rrbracket}$, $(\varepsilon_k)_{k \in \llbracket 1, \dots, n \rrbracket}$ and $(\delta_k)_{k \in \llbracket 1, \dots, n \rrbracket}$ belong to \mathcal{L}_{β} . Note that the words of \mathcal{L}'_{β} are defined on the alphabet $\llbracket -2[\beta], \dots, [\beta] \rrbracket$. Let $x, y \in \mathbb{Z}_{\beta}^+$. Then the expansion in base β of $x + y - [x + y]_{\beta}$ constructed digit by digit belongs to \mathcal{L}'_{β} , and it represents an element of $[0, 1[$.

In order to obtain a better upper bound for L_{\oplus} than those computed in [9], we define the following algorithm A .

1. We set for B an upper bound for L_{\oplus} , for instance the upper bound computed in Proposition 3.2, and we set $N = 1$.
2. We check whether there exists words $v = v_B \dots v_1 \in \mathcal{L}_{\beta}$ and $w = w_0 \dots w_{N-1} \in \mathcal{L}'_{\beta}$ such that the inequality given by Proposition 3.5 holds, that is, if we have for all $j \in J$

$$\left| \sum_{k=1}^B v_k \alpha_j^{-k} + \sum_{k=0}^{N-1} w_k \alpha_j^k - \alpha_j^N p_j(\vec{C}) \right| < \frac{3H'(j)(1 + |\alpha_j|)}{2} |\alpha_j|^N. \quad (3)$$

3. When (3) is satisfied for no $v \in \mathcal{L}_{\beta}$ and no $w \in \mathcal{L}'_{\beta}$, we decrease by 1 the value of B , we set $N = 1$ and we return to point 2.
4. When there exists $v \in \mathcal{L}_{\beta}$ and $w \in \mathcal{L}'_{\beta}$ such that (3) holds, we increase by 1 the value of N and we return to point 2.

Since $B \in \mathbb{N}$, there eventually exists an admissible word v such that (3) is satisfied for any $w \in \mathcal{L}_\beta^-$. This algorithm lowers the value of B , that is, it provides a better upper bound for L_\oplus^+ . However, we cannot check if this algorithm computes the value of L_\oplus^+ . Additionally, even if the algorithm computes the exact value of L_\oplus^+ after finitely many steps, we do not know when the exact value of L_\oplus^+ is reached. In this case, it suffices then to find two β -integers x and y such that $d_\beta(x+y)$ has a fractional part of the required length to deduce the exact value of L_\oplus^+ . We finally perform the same computations for the case of the difference of two positive β -integers in order to obtain L_\oplus .

Let us focus now on the case of cubic Pisot numbers. Let β be the positive root of the polynomial $P(X) = X^3 - k_1X^2 - k_2X - 1$, where $k_1, k_2 \in \mathbb{N}^2$ satisfy $k_1 \geq \max\{1, k_2\}$. Suppose moreover that P has two complex conjugate roots, that we denote by α and $\bar{\alpha}$. For k_2 large enough, this condition is satisfied when $k_2 \leq 2\sqrt{k_1}$, and is not satisfied otherwise.

Due to [3], we know that the finiteness property (\mathcal{F}) holds in this case. It is also proved in this article that, when β is a Pisot unit which satisfies (\mathcal{F}), then 0 is an inner point of \mathcal{T} . Hence $K = \min_{z \in \mathbb{Z}_\beta^+ \setminus \beta\mathbb{Z}_\beta^+} |\tau(z)| >$

0, and the required conditions of Proposition 3.2 hold for the class of Parry numbers we consider. We respectively denote by $L_\oplus(k_1, k_2)$ and $L_\oplus^+(k_1, k_2)$ the value of L_\oplus and L_\oplus^+ for the associated numeration system, defined by $d_\beta(1) = 0.k_1k_21$.

Proposition 3.6. *The algorithm A computes $L_\oplus(k_1, k_2)$ when $k_1 \leq 3$. We get:*

1. $L_\oplus(1, 0) = 11$ and $L_\oplus^+(1, 0) = 10$,
2. $L_\oplus(1, 1) \leq 6$,
3. $L_\oplus(2, 0) = L_\oplus^+(2, 0) = 7$,
4. $L_\oplus(2, 1) = 5$ and $L_\oplus^+(2, 1) = 4$,
5. $L_\oplus(2, 2) = L_\oplus^+(2, 2) = 5$,
6. $L_\oplus(3, 0) = L_\oplus^+(3, 0) = 4$,
7. $L_\oplus(3, 1) = L_\oplus^+(3, 1) = 4$,
8. $L_\oplus(3, 2) = 4$ and $L_\oplus^+(3, 2) = 3$,
9. $L_\oplus(3, 3) = L_\oplus^+(3, 3) = 5$.

Note that the inequality $L_\oplus(1, 1) \leq 6$ was already known, see [9].

Example 2. *When $d_\beta(1) = 0.101$, one has $\beta^{13} + \beta^9 + \beta^6 + \beta^3 + 1 - (\beta^{11} + \beta^8 + \beta^4 + \beta) = \beta^{11} + \beta^8 + \beta^4 + \beta + \beta^{-3} + \beta^{-6} + \beta^{-11}$ and $(\beta^4 + 1) + (\beta^3 + 1) = \beta^5 + \beta^2 + \beta^{-1} + \beta^{-5} + \beta^{-10}$. When $d_\beta(1) = 0.201$, one has $(\beta^5 + \beta^4 + \beta + 2) + (\beta^5 + \beta + 2) = \beta^6 + \beta^3 + 2\beta^2 + \beta^{-1} + 2\beta^{-2} + \beta^{-5} + \beta^{-7}$.*

From a practical point of view, we noticed that, except in the Tribonacci case, the algorithm A computes $L_\oplus(k_1, k_2)$ in a finite number of steps for any example of a pair (k_1, k_2) among those that satisfy the needed conditions, that is, for which β is the positive root of the polynomial $X^3 - k_1X^2 - k_2X - 1$ and admits two complex Galois conjugates.

3.3 The particular case of Tribonacci

We end our study by the Tribonacci case, that is, β is the real root of the polynomial $X^3 - X^2 - X - 1$. Since β admits two complex Galois conjugates, we set from now on $\alpha = \alpha_1$ and $\tau = \tau_1$. Note that this definition of τ coincides with the previous definition of τ given in Section 1.4. The previous method does not allow us to determine the exact value of $L_\oplus(1, 1)$, since the admissible word 100011, which represents the β -fractional part $\beta^{-1} + \beta^{-5} + \beta^{-6}$ is not discarded by the algorithm described in Section 3.2. Hence we do not know whether there exist two β -integers x and y such that $\{x+y\}_\beta = \beta^{-1} + \beta^{-5} + \beta^{-6}$. This explains why the particular case of Tribonacci needs additional computations.

We prove in this section that $L_\oplus(1, 1) = 5$, thanks to the following lemma.

Lemma 3.7. *Let β be the Tribonacci number, that is, $d_{\beta}(1) = 0.111$. Suppose that there exist $x, y \in \mathbb{Z}_{\beta}^{+}$ such that $l_f(x + y) = 6$. Then, we have:*

1. $\{x + y\}_{\beta} = \beta^{-1} + \beta^{-5} + \beta^{-6}$,
2. the real numbers $\frac{x-1}{\beta^3}$, $\frac{y-1}{\beta^3}$ and $\frac{[x+y]_{\beta} - \beta^2 - \beta}{\beta^3}$ are β -integers.

Proof. The first assertion is a consequence of Proposition 3.6, since we prove by pure computation that the relation (3) only holds for v an admissible word of length 6, namely when $v = 100011$. Additionally, we obtain only one possibility for the suffix of length 3 of $w \in \mathcal{L}_{\beta}$, which is $(-1)(-1)2$. This suffix corresponds to the suffix of length 3 of the expansion in base β of $x + y - [x + y]_{\beta}$ constructed digit by digit. This implies that the β -expansions of the β -integers x , y and $[x + y]_{\beta}$ must respectively admit 001, 001 and 110 as suffixes, which proves the second assertion. \square

Proposition 3.8. *When $d_{\beta}(1) = 0.111$, then $L_{\oplus}^{+} = 5$.*

Proof. Suppose that there exist x_0 and $y_0 \in \mathbb{Z}_{\beta}^{+}$ such that $\{x_0 + y_0\}_{\beta} = \beta^{-1} + \beta^{-5} + \beta^{-6}$. This means that the set $\mathcal{S}_6 = \{(x, y, z) \in (\mathbb{Z}_{\beta}^{+})^3, x + y = z + \beta^{-1} + \beta^{-5} + \beta^{-6}\}$ is not empty. Let $z_0 = [x_0 + y_0]_{\beta}$. Let $(x_0, y_0, z_0) \in \mathcal{S}_6$ such that $l_i(z_0)$, the length of the β -fractional part of z_0 , is minimal. Due to Lemma 3.7, there exists $(x_1, y_1, z_1) \in (\mathbb{Z}_{\beta}^{+})^3$ such that $x_0 = 1 + \beta^3 x_1$, $y_0 = 1 + \beta^3 y_1$ and $z_0 = \beta + \beta^2 + \beta^3 z_1$. Thus, we get:

$$\begin{aligned} \beta^3(x_1 + y_1) &= x_0 + y_0 - 2 \\ &= z_0 + \beta^{-1} + \beta^{-5} + \beta^{-6} - 2 \\ &= \beta^3 z_1 + \beta^{-1} + \beta^{-5} + \beta^{-6} + \beta + \beta^2 - 2 \\ &= \beta^3 z_1 + \beta^2 + \beta^{-2} + \beta^{-3}. \end{aligned}$$

We deduce that $x_1 + y_1 = z_1 + \beta^{-1} + \beta^{-5} + \beta^{-6}$, hence the 3-uple (x_1, y_1, z_1) belongs to \mathcal{S}_6 . However, $l_i(z_1) \leq l_i(z_0) - 3$, which contradicts the minimality of $l_i(z_0)$. Hence \mathcal{S}_6 is empty, and $L_{\oplus}^{+} < 6$. \square

Theorem 3.9. *When $d_{\beta}(1) = 0.111$, then $L_{\oplus} = 5$.*

Proof. Proposition 3.8 proves that $L_{\oplus}^{+} = 5$. In order to obtain the result on \mathbb{Z}_{β} instead of \mathbb{Z}_{β}^{+} , it suffices to note that, since $\mathbb{Z}_{\beta} = \pm\mathbb{Z}_{\beta}^{+}$, we can suppose that $x + y > 0$, thus $[x + y]_{\beta} \in \mathbb{Z}_{\beta}^{+}$ without loss of generality. Moreover, if $x + y > 0$, then x or y is positive, and the case where they are both positive is exactly the case previously studied. Thus, we only have to check whether there exist $x, -y \in \mathbb{Z}_{\beta}^{+}$, $x - y > 0$ such that $l_f(x - y) = 6$. However, we check by an exhaustive computation that there exists $N \in \mathbb{N}$ for which the inequality (3) used in the algorithm A does not hold for any admissible word v of length 6. Since $L_{\oplus} \geq 5$, this proves that $L_{\oplus} = L_{\oplus}^{+} = 5$. \square

Remark 8. *Let us explain why the algorithm A described in Section 3.2 does not improve the relation $5 \leq L_{\oplus} \leq 6$ given by [9] in the Tribonacci case. This is due to the fact that the inequality provided by Proposition 3.5, a priori satisfied by images under τ_j of β -integers, also extends to elements of \mathcal{T} . For instance, the complex number $\frac{1}{1-\alpha^3}$ belongs to \mathcal{T} , since it is the limit of the sequence $(\sum_{k=0}^n \alpha^{3k})_{n \in \mathbb{N}}$, which consists of images of β -integers under τ . Hence $\frac{1}{1-\alpha^3} + \frac{1}{1-\alpha^3} = \frac{2}{1-\alpha^3}$ is the sum of two elements which belong to \mathcal{T} . Moreover, $\frac{2}{1-\alpha^3}$ is the limit of the sequence $((\sum_{k=0}^n \alpha^{3k+1} + \alpha^{3k+2}) + \alpha^{-1} + \alpha^{-5} + \alpha^{-6})_{n \in \mathbb{N}}$, which consists of images of elements of $\text{Fin}(\beta)$, having $\beta^{-1} + \beta^{-5} + \beta^{-6}$ as β -fractional part, under τ . This means that, whereas there do not exist $x, y \in \mathbb{Z}_{\beta}$ such that $\{x + y\}_{\beta} = \beta^{-1} + \beta^{-5} + \beta^{-6}$, there exists a complex number, namely $\frac{2}{1-\alpha^3}$, which is the sum of two elements of \mathcal{T} and admits an α -expansion whose fractional part is of length 6.*

For any $x \in \text{Fin}(\beta) \cap [0, 1[$, let $d_{\beta}(x) = 0.v$ and $T.v = \overline{\tau(\{y \in \text{Fin}(\beta), \{y\}_{\beta} = x\})}$. Due to Messaoudi [27], the fact that the complex numbers $\frac{2}{1-\alpha^3}$ and $\frac{1}{1-\alpha^3}$ both have three α -expansions means that they belong to the fractal boundary of three different tiles $T.x$. By computation, one gets $\{\frac{2}{1-\alpha^3}\} \subset \mathcal{T}_{.100011} \cap \mathcal{T}_{.001} \cap \mathcal{T}_{.011}$ and $\{\frac{1}{1-\alpha^3}\} \subset \mathcal{T} \cap \mathcal{T}_{.001} \cap \mathcal{T}_{.011}$. Figure 4 gives a geometrical representation of these relations.




Fig. 4: Tiles \mathcal{T} , $\mathcal{T}_{.011}$, $\mathcal{T}_{.001}$ and $\mathcal{T}_{.100011}$

It would be interesting to compute the value of L_{\odot} for the Tribonacci case. Due to [9], we know that $4 \leq L_{\odot} \leq 5$. Similarly as in the case of addition, one checks by pure computation that there exists only one real number x such that $l_f(x) = 5$, namely $x = \beta^{-1} + \beta^{-3} + \beta^{-5}$, which could be the β -fractional part of the product of two β -integers. We do not know whether there exist two β -integers x and y such that $\{xy\}_{\beta} = \beta^{-1} + \beta^{-3} + \beta^{-5}$; nevertheless we find that the complex numbers $\frac{1+\alpha}{1-\alpha^3}$, $\frac{1}{1-\alpha^3}$ and $\frac{\alpha^2}{1-\alpha^3}$ belong to \mathcal{T} with $\frac{1+\alpha}{1-\alpha^3} \cdot \frac{1}{1-\alpha^3} = \frac{\alpha^2}{1-\alpha^3} + \alpha^{-1} + \alpha^{-3} + \alpha^{-5}$. Hence there exists an α -expansion of $\frac{1+\alpha}{1-\alpha^3} \cdot \frac{1}{1-\alpha^3}$, the product of two elements which belong to \mathcal{T} , which admits an α -fractional part whose fractional part is $\alpha^{-1} + \alpha^{-3} + \alpha^{-5}$. However, these informations are not sufficient to determine whether there exist $x, y \in \mathbb{Z}_{\beta}$ such that $\{xy\}_{\beta} = \beta^{-1} + \beta^{-3} + \beta^{-5}$.

Acknowledgements

I am indebted to Valérie Berthé and Pierre Arnoux for their advices and corrections. I also thank Julien Cassaigne, Petr Ambrož, Zuzana Másáková, Edita Pelantová, Shigeki Akiyama and Anne Siegel for many interesting talks, remarks and ideas developed in this paper. I strongly thank the referees for their valuable comments and corrections.

References

- [1] S. Akiyama. Pisot numbers and greedy algorithm. In *Number theory (Eger, 1996)*, pages 9–21. de Gruyter, Berlin, 1998.
- [2] S. Akiyama. Self affine tiling and Pisot numeration system. In *Number theory and its applications (Kyoto, 1997)*, volume 2 of *Dev. Math.*, pages 7–17. Kluwer Acad. Publ., Dordrecht, 1999.
- [3] S. Akiyama. Cubic Pisot units with finite beta expansions. In *Algebraic number theory and Diophantine analysis (Graz, 1998)*, pages 11–26. de Gruyter, Berlin, 2000.
- [4] S. Akiyama. On the boundary of self affine tilings generated by Pisot numbers. *J. Math. Soc. Japan*, 54(2):283–308, 2002.
- [5] S. Akiyama, F. Bassino, and C. Frougny. Automata for arithmetic Meyer sets. *Lecture Notes in Computer Science*, 2976:252–261, 2004.
- [6] S. Akiyama and A. Pethő. On canonical number systems. *Theoret. Comput. Sci.*, 270(1-2):921–933, 2002.
- [7] S. Akiyama, H. Rao, and W. Steiner. A certain finiteness property of Pisot number systems. *J. Number Theory*, 107(1):135–160, 2004.
- [8] S. Akiyama and J. M. Thuswaldner. A survey on topological properties of tiles related to number systems. *Geom. Dedicata*, 109:89–105, 2004.

- [9] P. Ambrož, C. Frougny, Z. Másáková, and E. Pelantová. Arithmetics on number systems with irrational bases. *Bull. Belgian Math. Soc. Simon Stevin*, 10:641–659, 2003.
- [10] D. Barache, B. Champagne, and J.-P. Gazeau. Pisot-cyclotomic quasilattices and their symmetry semigroups. In *Quasicrystals and discrete geometry (Toronto, ON, 1995)*, volume 10 of *Fields Inst. Monogr.*, pages 15–66. Amer. Math. Soc., Providence, RI, 1998.
- [11] J. Bernat. Arithmetic automaton for Perron numbers. *Accepté par Discrete Mathematics and Theoretical Computer Science*, 2005.
- [12] A. Bertrand. Développement en base de Pisot et répartition modulo 1. *C. R. Acad. Sc.*, 385:419–421, 1977.
- [13] Č. Burdík, C. Frougny, J. P. Gazeau, and R. Krejcar. Beta-integers as natural counting systems for quasicrystals. *J. Phys. A*, 31(30):6449–6472, 1998.
- [14] V. Canterini. Connectedness of geometric representation of substitutions of Pisot type. *Bull. Belg. Math. Soc. Simon Stevin*, 10(1):77–89, 2003.
- [15] V. Canterini and A. Siegel. Geometric representation of substitutions of Pisot type. *Trans. Amer. Math. Soc.*, 353(12):5121–5144 (electronic), 2001.
- [16] S. Fabre. Substitutions et β -systèmes de numération. *Theoret. Comput. Sci.*, 137(2):219–236, 1995.
- [17] C. Frougny. Confluent linear numeration systems. *Theoret. Comput. Sci.*, 106(2):183–219, 1992.
- [18] C. Frougny. On-line addition in real base. In *Mathematical foundations of computer science 1999 (Szklarska Porpolhka eba)*, volume 1672 of *Lecture Notes in Comput. Sci.*, pages 1–11. Springer, Berlin, 1999.
- [19] C. Frougny, Z. Másáková, and E. Pelantová. Complexity of infinite words associated with beta-expansions. *Theor. Inform. Appl.*, 38(2):163–185, 2004.
- [20] C. Frougny and B. Solomyak. Finite beta-expansions. *Ergodic Theory Dynam. Systems*, 12(4):713–723, 1992.
- [21] J.-P. Gazeau and J.-L. Verger-Gaugry. Geometric study of the set of β -integers for a Perron number and mathematical quasicrystals. *J. Théor. Nombres Bordeaux*, 16:1–25, 2004.
- [22] L.-S. Guimond, Z. Másáková, and E. Pelantová. Combinatorial properties of infinite words associated with cut-and-project sequences. *J. Théor. Nombres Bordeaux*, 15:697–725, 2003.
- [23] L.-S. Guimond, Z. Másáková, and E. Pelantová. Arithmetics of beta-expansions. *Acta Arith.*, 112(1):23–40, 2004.
- [24] S. Ito and M. Kimura. On Rauzy fractal. *Japan J. Indust. Appl. Math.*, 8(3):461–486, 1991.
- [25] J. C. Lagarias. Geometric models for quasicrystals I. Delone sets of finite type. *Discrete Comput. Geom.*, 21(2):161–191, 1999.
- [26] M. Lothaire. *Algebraic Combinatorics On Words*. Cambridge University Press, 2002.
- [27] A. Messaoudi. Frontière du fractal de Rauzy et système de numération complexe. *Acta Arith.*, 95(3):195–224, 2000.
- [28] Y. Meyer. *Algebraic numbers and harmonic analysis*. North-Holland Publishing Co., Amsterdam, 1972. North-Holland Mathematical Library, Vol. 2.
- [29] Y. Meyer. Quasicrystals, Diophantine approximation and algebraic numbers. In *Beyond quasicrystals (Les Houches, 1994)*, pages 3–16. Springer, Berlin, 1995.

- [30] R. V. Moody. Meyer sets and their duals. In *The mathematics of long-range aperiodic order (Waterloo, ON, 1995)*, volume 489 of *NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci.*, pages 403–441. Kluwer Acad. Publ., Dordrecht, 1997.
- [31] W. Parry. On the β -expansions of real numbers. *Acta Math. Acad. Sci. Hungar.*, 11:401–416, 1960.
- [32] N. Pytheas Fogg. *Substitutions in Dynamics, Arithmetics and Combinatorics (Berthé V., Ferenczi S., Mauduit C. and Siegel A.)*. Springer Lecture Notes in Mathematics, 1794, Berlin, 2002.
- [33] G. Rauzy. Nombres algébriques et substitutions. *Bull. Soc. Math. France*, 110(2):147–178, 1982.
- [34] G. Rauzy. Rotations sur les groupes, nombres algébriques et substitutions. *Séminaire de théorie des nombres*, 1987–1988.
- [35] A. Rényi. Representations for real numbers and their ergodic properties. *Acta Math. Acad. Sci. Hungar.*, 8:477–493, 1957.
- [36] K. Scheicher and J. M. Thuswaldner. Canonical number systems, counting automata and fractals. *Math. Proc. Cambridge Philos. Soc.*, 133(1):163–182, 2002.
- [37] K. Schmidt. On periodic expansions of Pisot numbers and Salem numbers. *Bull. London Math. Soc.*, 12:269–278, 1980.
- [38] D. Shechtman, I. Blech, D. Gratias, and J. Cahn. Metallic phase with with long-range orientational order and no translational symmetry. *Phys. Rev. Lett.*, 53:1951–1953, 1984.
- [39] V. F. Sirvent and Y. Wang. Self-affine tiling via substitution dynamical systems and Rauzy fractals. *Pacific J. Math.*, 206(2):465–485, 2002.
- [40] B. Solomyak. Substitutions, adic transformations, and beta-expansions. In *Symbolic dynamics and its applications (New Haven, CT, 1991)*, volume 135 of *Contemp. Math.*, pages 361–372. Amer. Math. Soc., Providence, RI, 1992.
- [41] W. P. Thurston. Groups, tilings and finite state automata. *Summer 1989 AMS Colloquium lectures*, 1989.