
GEOMETRIC AND PROJECTIVE INSTABILITY FOR THE GROSS-PITAEVSKI EQUATION

by Laurent Thomann

ABSTRACT. — Using variational methods, we construct approximate solutions for the Gross-Pitaevski equation which concentrate on circles in \mathbb{R}^3 . These solutions will help to show that the L^2 flow is unstable for the usual topology and for the projective distance.

1. Introduction

In this paper we deal with the equations

$$(1) \quad \begin{cases} ih\partial_t u + h^2\Delta u - |x|^2 u = a_h h^2 |u|^2 u, & (t, x) \in \mathbb{R}^{1+3}, \\ u(0, x) = u_0(x) \in L^2(\mathbb{R}^3), \end{cases}$$

where $h > 0$ is a small parameter and a_h a constant which depends on h , that can be either positive (defocusing case) or negative (focusing case). In all the paper we assume that there exists a constant $A > 0$, independent of h , such that $|a_h| \leq A$.

This equation appears in the study of Bose-Einstein condensates; for more details see [7].

In the following we will refer to the definitions:

LAURENT THOMANN, Université Paris-Sud, Mathématiques, Bât 425, Tel 0169155785,
91405 Orsay Cedex. • *E-mail* : laurent.thomann@math.u-psud.fr
Url : <http://www.math.u-psud.fr/~thomann>

2000 Mathematics Subject Classification. — 35Q55; 35B35; 81Q05.

Key words and phrases. — non linear Schrödinger equation, instability .

DEFINITION 1.1. — (Geometric instability) We say that the Cauchy problem (1) is geometrically unstable if there exist $u_h^1, u_h^2 \in L^2(\mathbb{R}^3)$ solutions of (1) with initial data $u_h^1(0), u_h^2(0) \in L^2(\mathbb{R}^3)$ such that $\|u_h^1(0)\|_{L^2}, \|u_h^2(0)\|_{L^2} \leq C$ where C is a constant independent of h and $t_h > 0$ such that

$$\frac{\|(u_h^2 - u_h^1)(t_h)\|_{L^2}}{\|(u_h^2 - u_h^1)(0)\|_{L^2}} \longrightarrow +\infty \text{ when } h \longrightarrow 0.$$

DEFINITION 1.2. — (Projective instability) We say that the Cauchy problem (1) is projectively unstable if there exist $u_h^1, u_h^2 \in L^2(\mathbb{R}^3)$ solutions of (1) with initial data $u_h^1(0), u_h^2(0) \in L^2(\mathbb{R}^3)$ such that $\|u_h^1(0)\|_{L^2}, \|u_h^2(0)\|_{L^2} \leq C$ where C is a constant independent of h and $t_h > 0$ such that

$$\frac{d_{\text{pr}}(u_h^2(t_h), u_h^1(t_h))}{d_{\text{pr}}(u_h^2(0), u_h^1(0))} \longrightarrow +\infty \text{ when } h \longrightarrow 0.$$

Here d_{pr} denotes the complex projective distance defined by

$$d_{\text{pr}}(v_1, v_2) = \arccos\left(\frac{|\langle v_1, v_2 \rangle|}{\|v_1\|_{L^2} \|v_2\|_{L^2}}\right) \text{ for } v_1, v_2 \in L^2(\mathbb{R}^3).$$

NOTATIONS 1.3. — In this paper c, C denote constants the value of which may change from line to line. These constants will always be independent of h . We use the notations $a \sim b$, $a \lesssim b$, $a \gtrsim b$, if $\frac{1}{C}b \leq a \leq Cb$, $a \leq Cb$, $b \leq Ca$ respectively. We write $a \ll b$, $a \gg b$ if $a \leq Kb$, $a \geq Kb$ for some large constant K which is independent of h .

The first result of this paper is

THEOREM 1.4. — Let $h^{-1} \in \mathbb{N}$. In each of the following cases, there exist $c_0 > 0$ and $u_h^1, u_h^2 \in L^2(\mathbb{R}^3)$ solutions of (1) with initial data $\|u_h^2(0)\|_{L^2}, \|u_h^1(0)\|_{L^2} \rightarrow \kappa$ such that if $|a_h|\kappa^2 \leq c_0$, we have:

(i) Assume a is independent of h and $\kappa|a|t \gg 1$,

$$\frac{\|(u_h^2 - u_h^1)(t)\|_{L^2}}{\|(u_h^2 - u_h^1)(0)\|_{L^2}} \gtrsim |a|\kappa t.$$

(ii) Assume $|a_h|t_h \rightarrow +\infty$ when $h \rightarrow 0$ with $t_h \ll \log \frac{1}{h}$, then

$$\sup_{0 \leq t \leq t_h} \|(u_h^2 - u_h^1)(t)\|_{L^2} \gtrsim 1,$$

but

$$\|(u_h^2 - u_h^1)(0)\|_{L^2} \rightarrow 0.$$

In particular, the Cauchy problem (1) is geometrically unstable

Denote by $x = (x_1, x_2, x_3)$ the current point in \mathbb{R}^3 . In cylindrical coordinates ($x_1 = r \cos \theta, x_2 = r \sin \theta, x_3 = y$), the functions considered in Theorem 1.4 take the form

$$(2) \quad u_h(0, x) = \kappa_h h^{-\frac{1}{2}} e^{i \frac{k^2}{h} \theta} v_0 \left(\frac{r-k}{\sqrt{h}}, \frac{y}{\sqrt{h}} \right),$$

where $k \in \mathbb{N}$, $v_0 \in L^2(\mathbb{R}^2)$ and

$$(3) \quad u_h(t, x) = u_h(0, x) e^{-i \lambda_h t} + w_h(t, x),$$

with w_h a small error term in $L^2(\mathbb{R}^3)$, at least for times when instability effects occur.

The Ansatz (2) shows that the function u in (3) will concentrate on the circle ($x_1^2 + x_2^2 = k^2, x_3 = 0$) in \mathbb{R}^3 .

To prove Theorem 1.4, we consider two initial data of the form (2) associate with κ and κ' such that $|\kappa' - \kappa|$ is small, and therefore the initial data are close in the L^2 -norm, but we will see that the solutions do not remain close to each other after a time t .

The construction of two solutions to (1) of the form (2),(3) which concentrate on disjoint circles yield the following result

THEOREM 1.5. — *Let $h^{-1} \in \mathbb{N}$. There exist $c_0 > 0$ and $u_h^1, u_h^2 \in L^2(\mathbb{R}^3)$ solutions of (1) with initial data $\|u_h^2(0)\|_{L^2}, \|u_h^1(0)\|_{L^2} \rightarrow \kappa$ such that if $|a_h| \kappa^2 \leq c_0$ and $|a_h| t_h \rightarrow +\infty$ when $h \rightarrow 0$ with $t_h \ll \log \frac{1}{h}$, we have*

$$\sup_{0 \leq t \leq t_h} d_{pr}(u_h^2(t), u_h^1(t)) \gtrsim 1,$$

but

$$d_{pr}(u_h^2(0), u_h^1(0)) \rightarrow 0.$$

In particular, the Cauchy problem (1) is projectively unstable.

The part (i) of Theorem 1.4 shows that there is no Lipschitz dependence between the solutions of equation (1) and the initial data in the regime $\kappa a t \gg 1$, whereas the part (ii) and Theorem 1.5 assert that the dependence is not continuous, but for larger times. Both types of instabilities are nonlinear behaviour, but the first one is weaker than the second.

The instability results of Theorem 1.4 are not new in the case $a > 0$. R. Carles [3] shows the instability, for finite times, of the equation

$$i h \partial_t v + h^2 \Delta v - |x|^2 v = f(h^k |v|^2) v, \quad (t, x) \in \mathbb{R}^{1+n},$$

when $n \geq 2$, $1 < k < n$, and $f \in C^\infty(\mathbb{R}_+, \mathbb{R})$ with $f' > 0$.

In [1], N. Burq, P. Gérard and N. Tzvetkov have pointed out geometric instability for the cubic Schrödinger equation $i \partial_t u + \Delta_{\mathbb{S}^2} u = a |u|^2 u$ on \mathbb{S}^2 when $a > 0$. This phenomenon doesn't occur on $L^2(\mathbb{R}^3)$ for the equation $i \partial_t u + \Delta u = a |u|^2 u$ in $L^2(\mathbb{R}^3)$, it is therefore strongly related to the geometry of the operator and

of the manifold we work on. Here there is no semiclassical parameter in the equations, but we could obtain similar results in this latter case with a scaling argument, as these instability effects are local. There are stronger instability phenomena in H^s norm, for $0 < s < \frac{1}{2}$ or for s negative, for more details see [5] or [4] for the one dimensional case.

In [2], N. Burq and M. Zworski prove Theorem 1.4 in the case $a > 0$. To obtain geometric instability, they expand the solution on the Hilbertian basis given by the eigenfunctions of $-h^2\Delta + |x|^2$. The nonlinear term in (1) induces a phase shift in time for the groundstate and this yields the result. We will give a more precise description of the solution by solving a pertubated eigenvalue problem for the harmonic oscillator and this will also treat the focusing case. They also obtain projective instability for the equation

$$ih\partial_t u + h^2\Delta u - V(x)u = ah^2|u|^2u,$$

where V is a cylindrically symmetric potential with respect to the variable $y = x_3$, but they have to add the following assumption: Denote by $r = \sqrt{x_1^2 + x_2^2}$ then the function $(r, y) \mapsto V(r, y) + r^{-2}$ has two distinct absolute non-degenerate minima $(r_j, y_j), j = 1, 2$, and its Hessian at (r_j, y_j) are equal. We use a variational method to construct quasimodes which are localized on circles in \mathbb{R}^3 , which allows to remove such an hypothesis. This idea comes from an unpublished work from N. Burq, P. Gérard and N. Tzvetkov.

Thanks to the form $F(|u|^2)u$ of the nonlinearity in (1), we look for a solution u which writes $u(t, x) = e^{-i\lambda t}f(x)$. Then f has to satisfy

$$(-h^2\Delta + |x|^2)f = h\lambda f - a_h h^2|f|^2f.$$

In the case $a_h = 0$, f is an eigenvector of the operator $-h^2\Delta + |x|^2$ associate with the eigenvalue $h\lambda$. In the general case, the term $a_h h^2|f|^2f$ will be treated as a perturbation of the linear problem

$$(-h^2\Delta + |x|^2)f = h\lambda f.$$

In fact, we will find a development in powers of h of $h\lambda$ and f

$$h\lambda \sim \sum_{k \geq 0} \mu_k h^k, \quad f \sim \sum_{k \geq 0} f_k h^k,$$

by solving a cascade of equations. This will be done in cylindrical coordinates: Write $x = (x_1, x_2, x_3)$ and make the cylindrical change of variables $x_1 = r \cos \theta$, $x_2 = r \sin \theta$ and $x_3 = y$ with $(r, \theta, y) \in \mathbb{R}_+^* \times [0, 2\pi[\times \mathbb{R}$. Then the Laplace operator takes the form

$$\Delta = \frac{1}{r^2}\partial_\theta^2 + \partial_r^2 + \frac{1}{r}\partial_r + \partial_y^2.$$

Let κ be a positive constant and k a positive integer, we want to find a solution of (1) of the form

$$(4) \quad \tilde{u} = \kappa h^{-\frac{1}{2}} e^{-i\lambda t} e^{i\frac{k^2}{h}\theta} \tilde{v}(r, y, h),$$

where λ is a constant to be determined, and \tilde{v} a real function which therefore has to satisfy

$$-h^2(\partial_r^2 + \partial_y^2)\tilde{v} + \left(\frac{k^4}{r^2} + r^2 + y^2\right)\tilde{v} = \lambda h\tilde{v} - a_h h^2 \kappa^2 \tilde{v}^3 + h^2 \frac{1}{r} \partial_r \tilde{v}.$$

Notice that we have to choose $h^{-1} \in \mathbb{N}$ so that (4) makes sense for all $k \in \mathbb{N}$. We try to construct \tilde{v} which concentrates exponentially at the minimum of the potential $V = \frac{k^4}{r^2} + r^2 + y^2$, i.e. at $(r, y) = (k, 0)$.

Thus we make the change of variables $r = k + \sqrt{h}\rho$, $y = \sqrt{h}\sigma$ and set $\tilde{v}(r, y, h) = v\left(\frac{r-k}{\sqrt{h}}, \frac{y}{\sqrt{h}}, h\right)$.

We write the Taylor expansion of V in h :

$$\begin{aligned} \frac{k^4}{(k + \sqrt{h}\rho)^2} + (k + \sqrt{h}\rho)^2 + h\sigma^2 &= 2k^2 + (4\rho^2 + \sigma^2)h - \frac{4}{k}\rho^3 h^{\frac{3}{2}} \\ &\quad + \frac{5}{k^2}\rho^4 h^2 + R(\rho, h)h^{\frac{5}{2}}. \end{aligned}$$

Then v has to be solution of

$$(5) \quad \begin{aligned} Eq(v) := P_0 v - \frac{\lambda h - 2k^2}{h} v + a_h \kappa^2 v^3 - h^{\frac{1}{2}} \left(\frac{1}{k + \sqrt{h}\rho} \partial_\rho v + \frac{4}{k} \rho^3 v \right) \\ + \frac{5}{k^2} \rho^4 h v - h^{\frac{3}{2}} R v = 0, \end{aligned}$$

where $P_0 = -(\partial_\rho^2 + \partial_\sigma^2) + (4\rho^2 + \sigma^2)$. Now, write

$$\begin{aligned} v(\rho, \sigma, h) &= v_0(\rho, \sigma) + h^{\frac{1}{2}} v_1(\rho, \sigma) + h v_2(\rho, \sigma) + h^{\frac{3}{2}} w(\rho, \sigma, h) \\ \frac{\lambda h - 2k^2}{h} &= E_0 + h^{\frac{1}{2}} E_1 + h E_2 + h^{\frac{3}{2}} E_3(h). \end{aligned}$$

By identifying the powers of h we obtain the following equations:

$$(6) \quad P_0 v_0 = E_0 v_0 - a_h \kappa^2 v_0^3,$$

$$(7) \quad P_0 v_1 = E_0 v_1 + E_1 v_0 - 3a_h \kappa^2 v_0^2 v_1 + \frac{1}{k} \partial_\rho v_0 + \frac{4}{k} \rho^3 v_0,$$

$$(8) \quad \begin{aligned} P_0 v_2 &= E_0 v_2 + E_1 v_1 + E_2 v_0 - 3a_h \kappa^2 (v_0^2 v_2 + v_0 v_1^2) + \frac{1}{k} \partial_\rho v_1 \\ &\quad + \frac{4}{k} \rho^3 v_1 - \frac{1}{k^2} \rho \partial_\rho v_0 - \frac{5}{k^2} \rho^4 v_0. \end{aligned}$$

REMARK 1.6. — In the sequel we only mention the dependence in k , κ and a of the v_j and E_j when necessary. Moreover we write $a = a_h$.

2. Construction of the quasimodes

PROPOSITION 2.1. — *There exists a constant $c_0 > 0$ such that if $|a|\kappa^2 \leq c_0$, there exist $E_0 > 0$ and $v_0 \in L^2(\mathbb{R}^2)$ satisfying $v_0 \geq 0$ and $\|v_0\|_{L^2(\mathbb{R}^2)} = 1$, which solve (6).*

For $\psi \in \mathcal{S}'(\mathbb{R}^2)$, denote by $\hat{\psi}$ its Fourier transform, with the convention

$$\hat{\psi}(\zeta) = \int_{\mathbb{R}^2} e^{-i\zeta \cdot x} \psi(x) dx,$$

for $\psi \in L^1(\mathbb{R}^2)$.

We use a variational method based on Rellich's criterion.

PROPOSITION 2.2. — ([8], p 247) *The set*

$$S = \left\{ \psi \mid \int_{\mathbb{R}^2} |\psi(x)|^2 dx = 1, \int_{\mathbb{R}^2} (1 + |x|^2) |\psi(x)|^2 dx \leq 1, \int_{\mathbb{R}^2} (1 + |\zeta|^2) |\hat{\psi}(\zeta)|^2 d\zeta \leq 1 \right\},$$

is a compact subset of $L^2(\mathbb{R}^2)$.

Proof of Proposition 2.1. — We minimize the functional

$$J(u, a) = \int \left(|\nabla u|^2 + (4\rho^2 + \sigma^2)|u|^2 + \frac{1}{2}a\kappa^2|u|^4 \right),$$

on the space

$$H = \left\{ u \in H^1(\mathbb{R}^2), (\rho^2 + \sigma^2)^{\frac{1}{2}} u \in L^2(\mathbb{R}^2), \|u\|_{L^2} = 1 \right\}.$$

Now, on H we have the inequality

$$\|u\|_{L^4} \leq C \|u\|_{H^{\frac{1}{2}}} \leq C \|u\|_{L^2}^{\frac{1}{2}} \|\nabla u\|_{L^2}^{\frac{1}{2}} \leq C \|\nabla u\|_{L^2}^{\frac{1}{2}}.$$

Thus, there exists $c_0 > 0$ such that

$$\frac{1}{2}a\kappa^2 \int |u|^4 \leq \frac{1}{2} \int |\nabla u|^2,$$

as soon as $|a|\kappa^2 \leq c_0$, which we suppose from now.

Let $(u_n)_{n \geq 1}$ be a minimizing sequence. First, we can choose $u_n \geq 0$, because $|u_n|$ is also minimizing, as $|\nabla |u_n|| \leq |\nabla u_n|$. We have

$$\int \left(\frac{1}{2} |\nabla u_n|^2 + (4\rho^2 + \sigma^2) u_n^2 \right) \leq J(u_n, a\kappa^2) \leq C,$$

with C independent of a , κ and n . We are able to apply Rellich's criterion: there exists $v_0 \in H$ with $v_0 \geq 0$ such that, up to a subsequence, $u_n \rightarrow v_0$, and the lower semi-continuity of J ensures

$$J(v_0, a\kappa^2) = \inf_{u \in H} J(u, a\kappa^2).$$

Then there exists a Lagrange multiplier E_0 such that

$$P_0 v_0 = -(\partial_r^2 + \partial_y^2)v_0 + (4\rho^2 + \sigma^2)v_0 = E_0 v_0 - a\kappa^2 v_0^3,$$

and E_0 is given by

$$E_0 = \int (|\nabla v_0|^2 + (4\rho^2 + \sigma^2)v_0^2 + a\kappa^2 v_0^4).$$

□

PROPOSITION 2.3. — *Let $|a|\kappa^2 \leq c_0$. There exist constants $C, c > 0$ independent of a, κ such that for $0 \leq j \leq 2$*

$$(9) \quad \left| (I - \Delta)^{\frac{j}{2}} v_0(\rho, \sigma) \right| \leq C e^{-c(|\rho| + |\sigma|)}.$$

Proof. — We denote by $\xi = (\rho, \sigma)$, and we define $\varphi_\varepsilon(\xi) = e^{\frac{|\xi|}{1+\varepsilon|\xi|}}$. The function φ_ε is bounded and

$$(10) \quad |\nabla \varphi_\varepsilon| \leq \varphi_\varepsilon \quad \text{a.e.}$$

We multiply (6) by $\varphi_\varepsilon v_0$ and integrate over \mathbb{R}^2 :

$$\int \nabla(\varphi_\varepsilon v_0) \nabla v_0 + \int \varphi_\varepsilon |\xi|^2 v_0^2 \leq E_0 \int \varphi_\varepsilon v_0^2 + |a|\kappa^2 \int \varphi_\varepsilon v_0^4.$$

We compute $\nabla(\varphi_\varepsilon v_0) = v_0 \nabla \varphi_\varepsilon + \varphi_\varepsilon \nabla v_0$, and use (10) to obtain

$$\int (\varphi_\varepsilon |\nabla v_0|^2 + \varphi_\varepsilon |\xi|^2 v_0^2) \leq E_0 \int \varphi_\varepsilon v_0^2 + |a|\kappa^2 \int \varphi_\varepsilon v_0^4 + \int \varphi_\varepsilon v_0 |\nabla v_0|.$$

We set $w_0 = \varphi_\varepsilon^{\frac{1}{4}} v_0$, then

$$(11) \quad \nabla w_0 = \frac{1}{4} \varphi_\varepsilon^{-\frac{3}{4}} \nabla \varphi_\varepsilon v_0 + \varphi_\varepsilon^{\frac{1}{4}} \nabla v_0.$$

From the Gagliardo-Nirenberg inequality in dimension 2

$$\|w_0\|_{L^4}^4 \leq C \|w_0\|_{L^2}^2 \|\nabla w_0\|_{L^2}^2,$$

together with (11) we deduce

$$\int \varphi_\varepsilon v_0^4 \leq C \int (\varphi_\varepsilon^{\frac{1}{2}} v_0^2) \int \varphi_\varepsilon^{\frac{1}{2}} (v_0^2 + |\nabla v_0|^2).$$

As $\int v_0^2 = 1$ and $\int |\nabla v_0|^2 \leq C$, Jensen's inequality gives

$$(12) \quad \begin{aligned} \int \varphi_\varepsilon v_0^4 &\leq C \left(\int \varphi_\varepsilon v_0^2 \right)^{\frac{1}{2}} \left(\int \varphi_\varepsilon (v_0^2 + |\nabla v_0|^2) \right)^{\frac{1}{2}} \\ &\leq \frac{1}{16c_0} \int \varphi_\varepsilon |\nabla v_0|^2 + C \int \varphi_\varepsilon v_0^2. \end{aligned}$$

We also have

$$(13) \quad \int \varphi_\varepsilon v_0 |\nabla v_0| \leq \frac{1}{4} \int \varphi_\varepsilon |\nabla v_0|^2 + C \int \varphi_\varepsilon v_0^2.$$

Now, write for $R > 0$

$$\int \varphi_\varepsilon v_0^2 = \int_{|\xi| < R} \varphi_\varepsilon v_0^2 + \int_{|\xi| \geq R} \varphi_\varepsilon v_0^2 \leq e^R \int v_0^2 + \frac{1}{R^2} \int \varphi_\varepsilon |\xi|^2 v_0^2,$$

and deduce that for R big enough, independent of ε , there exists a constant C independent of ε satisfying

$$\int \varphi_\varepsilon (|\nabla v_0|^2 + |\xi|^2 v_0^2) \leq C.$$

Letting ε tend to 0 yields

$$(14) \quad e^{\frac{|\xi|}{2}} \nabla v_0 \in L^2 \quad \text{and} \quad e^{\frac{|\xi|}{2}} |\xi| v_0 \in L^2.$$

With the help of equation (6), compute

$$\begin{aligned} \Delta \left(v_0 e^{\frac{1}{4}(\rho+\sigma)} \right) &= a\kappa^2 v_0^3 e^{\frac{1}{4}(\rho+\sigma)} + (4\rho^2 + \sigma^2 - E_0) v_0 e^{\frac{1}{4}(\rho+\sigma)} \\ &\quad + \frac{1}{2} (1, 1) \cdot \nabla v_0 e^{\frac{1}{4}(\rho+\sigma)} + \frac{1}{16} v_0 e^{\frac{1}{4}(\rho+\sigma)}. \end{aligned}$$

According to (14), each term of the right hand side is in L^2 , excepted maybe the first one. But denote by $\tilde{v}_0 = v_0 e^{\frac{1}{12}(\rho+\sigma)}$, then (14) shows that $\tilde{v}_0 \in H^1(\mathbb{R}^2)$ and consequently $v_0 \in L^6(\mathbb{R}^2)$.

Hence, with the inequality $\|w\|_{L^\infty}^2 \lesssim \|w\|_{L^2} \|\Delta w\|_{L^2}$ applied to $w = v_0 e^{\frac{1}{4}(\rho+\sigma)}$ we deduce $v_0 \leq C e^{-\frac{1}{4}(\rho+\sigma)}$.

The same can be done with σ replaced with $-\sigma$ or ρ by $-\rho$. Therefore $v_0 \leq C e^{-\frac{1}{4}(|\rho|+|\sigma|)}$. Equation (6) and the previous estimate give

$$|\Delta v_0(\rho, \sigma)| \leq C e^{-c(|\rho|+|\sigma|)}.$$

To obtain the last estimation of Proposition 2.3, use the interpolation inequality

$$\|\nabla w\|_{L^\infty}^2 \leq \|w\|_{L^\infty} \|\Delta w\|_{L^\infty},$$

applied to $w = v_0 e^{c(\pm\rho \pm \sigma)}$. \square

We are now able to describe the behaviour of $E_0(a\kappa^2)$ and $v_0(a\kappa^2)$ when $a\kappa^2 \rightarrow 0$:

PROPOSITION 2.4. —

$$v_0(a\kappa^2) \longrightarrow \frac{2^{\frac{1}{4}}}{\pi^{\frac{1}{2}}} e^{-(\rho^2 + \frac{1}{2}\sigma^2)} \quad \text{in } L^2(\mathbb{R}^2) \quad \text{when } a\kappa^2 \longrightarrow 0,$$

and

$$(15) \quad E_0(a\kappa^2) = 3 + \frac{\sqrt{2}}{2\pi} a\kappa^2 + o(a\kappa^2).$$

Proof. — The function $u_0 = \frac{2^{\frac{1}{4}}}{\pi^{\frac{1}{2}}} e^{-(\rho^2 + \frac{1}{2}\sigma^2)}$ is the unique positive element in H that realises the infimum of $J(u, 0)$, and is the first eigenfunction of $P_0 = -\Delta + (4\rho^2 + \sigma^2)$ associate with the eigenvalue $E_0(0) = 3$. See [6], p 7 for details. For $|a|\kappa^2 \leq c_0$ we have

$$(16) \quad \|v_0(a\kappa^2)\|_{L^2} = 1, \quad \|\nabla v_0(a\kappa^2)\|_{L^2} \leq C, \quad \text{and} \quad \|\xi v_0(a\kappa^2)\|_{L^2} \leq C.$$

By Rellich's criterion, $(v(a\kappa^2))_{|a|\kappa^2 \leq c_0}$ is compact in H ; let \mathcal{A} be its adherence set. If $u \in \mathcal{A}$, there exists a sequence $b_n = a_n \kappa_n^2 \longrightarrow 0$ satisfying $v_0(b_n) \longrightarrow u$ in L^2 . As $v_0(b_n)$ realises the infimum of $J(v, b_n)$:

$$J(v_0(b_n), b_n) \leq J(u_0, b_n) = 3 + \frac{1}{2} b_n \int |u_0|^4,$$

therefore, $J(u, 0) \leq 3$. As $u \geq 0$, we conclude $u = u_0$, i.e. $\mathcal{A} = \{u_0\}$ and

$$v_0(a\kappa^2) \longrightarrow u_0 \quad \text{in } L^2(\mathbb{R}^2) \quad \text{when } a\kappa^2 \longrightarrow 0.$$

Moreover $|v(a\kappa^2)|, |u_0| \leq C$, then the convergence is also in L^4 . Now, the self-adjointness of P_0 gives

$$0 = \langle (P_0 - 3)u_0, v(a\kappa^2) \rangle = (E_0(a\kappa^2) - 3) \int v(a)u_0 - a\kappa^2 \int v^3(a)u_0,$$

then from $\int v(a\kappa^2)u_0 \longrightarrow \int u_0^2 = 1$ and $\int v(a\kappa^2)^3 u_0 \longrightarrow \int u_0^4 = \frac{\sqrt{2}}{2\pi}$ we conclude $E_0(a\kappa^2) = 3 + \frac{\sqrt{2}}{2\pi} a\kappa^2 + o(a\kappa^2)$. \square

PROPOSITION 2.5. — *Let $|a\kappa^2| \leq c_0$. There exist $E_1, E_2 \in \mathbb{R}$ and $v_1, v_2 \in L^2(\mathbb{R}^2)$ satisfying $v_1, v_2 \geq 0$ and $\|v_1\|_{L^2(\mathbb{R}^2)}, \|v_2\|_{L^2(\mathbb{R}^2)} \sim 1$, which solve (7) and (8).*

Moreover there exists $c > 0$ such that for $l = 1, 2$ and $0 \leq j \leq 2$

$$(17) \quad \left| (I - \Delta)^{\frac{j}{2}} v_l(\rho, \sigma) \right| \leq C e^{-c(|\rho| + |\sigma|)}.$$

Proof. — Equation (7) writes

$$(P(a\kappa^2) - E_0) v_1 = (-\partial_\rho^2 + \partial_\sigma^2 + V) v_1 = E_1 v_0 + \frac{1}{k} \partial_\rho v_0 + \frac{4}{k} \rho^3 v_0,$$

where we denote by $P(a\kappa^2) = P_0 + 3a\kappa^2 v_0^2$ and $V = 4\rho^2 + \sigma^2 + 3a\kappa^2 v_0^2 - E_0$. The potential V is so that $V \longrightarrow \infty$ as $|(\rho, \sigma)| \longrightarrow \infty$, then the spectrum

$\sigma(P(a))$ of $P(a\kappa^2)$ is purely discrete and the eigenvalues are given by the min-max principle (see [8] p. 120).

The first eigenvalue of $P(a\kappa^2)$ is therefore given by

$$\mu_0(a\kappa^2) = \inf_{u \in H} \int (|\nabla u|^2 + (4\rho^2 + \sigma^2)u^2 + 3a\kappa^2 v_0^2 u^2) - E_0(a\kappa^2),$$

and there exists $w_0 \in H$ with $w_0 \geq 0$ satisfying

$$(P(a\kappa^2) - E_0) w_0 = (P_0 - E_0(a\kappa^2) + 3a\kappa^2 v_0^2) w_0 = \mu_0(a) w_0,$$

and one shows, as in the proof of (2.4) that $w_0 \rightarrow u_0$ in $L^2 \cap L^4$.

Multiply (6) by u_0 and integrate

$$3a\kappa^2 \int v_0^2 w_0 u_0 + (3 - E_0(a\kappa^2)) \int w_0 u_0 = \mu_0(a\kappa^2) \int w_0 u_0,$$

then according to (15), $\mu_0(a\kappa^2) \sim \frac{\sqrt{2}}{\pi} a\kappa^2$ when $a\kappa^2 \rightarrow 0$. If $a > 0$ and $a\kappa^2$ is small enough we can conclude that $0 \notin \sigma(P(a))$.

Let's look at the case $a < 0$:

According to the min-max principle, the second eigenvalue of $P(a\kappa^2)$ is

$$\mu_1(a\kappa^2) = \inf_{u \in H, u \perp w_0} \int (|\nabla u|^2 + (4\rho^2 + \sigma^2)u^2 + 3a\kappa^2 v_0^2 u^2) - E_0(a\kappa^2),$$

and let w_1 realise the infimum.

We also have

$$5 = \inf_{u \in H, u \perp u_0} \int (|\nabla u|^2 + (4\rho^2 + \sigma^2)u^2) = \inf_{u \in H, u \perp u_0} J(u, 0),$$

realised for u_1 , the second normalised Hermite function. Now, define $\tilde{u} = \alpha w_1 + \beta w_0$ with α, β such that $\|\tilde{u}\|_{L^2} = \alpha^2 + \beta^2 = 1$ and $\alpha \int w_1 u_0 + \beta \int w_0 u_0 = 0$, then $\tilde{u} \in H$ and $\tilde{u} \perp u_0$. Notice that $|\alpha| \rightarrow 1$ and $\beta \rightarrow 0$ as $a\kappa^2 \rightarrow 0$.

One has $5 = J(u_1, 0) \leq J(\tilde{u}, 0)$, then we obtain $5 \leq \mu_1(a\kappa^2) + \varepsilon(a\kappa^2)$ with $\varepsilon(a\kappa^2) \rightarrow 0$ as $a\kappa^2 \rightarrow 0$, therefore $\mu_1(a\kappa^2) \geq 4$ for a small enough, and $0 \notin \sigma(P(a\kappa^2))$.

As a conclusion, for each choice of E_1 , equation (7) admits a solution $v_1 \in L^2$ as the second right hand side f is in L^2 . However, if we choose E_1 so that $f \perp v_0$, we also have $\|v_1\|_{L^2} \leq C$ uniformly in $|a|\kappa^2 \leq c_0$, as the eigenvalue $E_0(a\kappa^2)$ is simple.

The estimations (17) are obtained as in the proof of Proposition 2.3.

By the same argument we infer the existence of v_2 and E_2 which solve equation (8) and satisfy the estimates (17). \square

Take $\chi \in C_0^\infty(\mathbb{R})$ such that $\chi \geq 0$, $\text{supp}\chi \subset [\frac{1}{2}, \frac{3}{2}]$ and $\chi = 1$ on $[\frac{3}{4}, \frac{5}{4}]$. Set $v = \chi(\sqrt{h}\rho)(v_0 + h^{\frac{1}{2}}v_1 + hv_2)$, $\tilde{v}(r, y, h) = v(\frac{r-k}{\sqrt{h}}, \frac{y}{\sqrt{h}}, h)$ and $\lambda = \frac{2k^2}{h} + E_0 + h^{\frac{1}{2}}E_1 + hE_2$, and define

$$(18) \quad u_{app} = \kappa h^{-\frac{1}{2}} e^{-i\lambda t} e^{i\frac{k^2}{h}\theta} \tilde{v}.$$

Recall that, according to (15),

$$E_0(a\kappa^2) = 3 + \frac{\sqrt{2}}{2\pi} a\kappa^2 + o(a\kappa^2).$$

PROPOSITION 2.6. — *The function u_{app} defined by (18) satisfies*

$$(19) \quad ih\partial_t u_{app} + h^2 \Delta u_{app} - |x|^2 u_{app} = ah^2 |u_{app}|^2 u_{app} + R(h)$$

with

$$(20) \quad \|(|x|^2 + 1) R(h) \|_{L^2} \lesssim h^{\frac{5}{2}} \quad \text{and} \quad \| \Delta R(h) \|_{L^2} \lesssim h^{\frac{1}{2}}.$$

Proof. — By construction, $w = v_0 + h^{\frac{1}{2}} v_1 + h v_2$ satisfies $Eq(w) = h^{\frac{5}{2}} R_1(h)$ where Eq is defined by (5), and according to Propositions 2.4 and 2.6

$$|R_1(h)| \lesssim \left(\frac{1}{(k + \sqrt{h}\rho)^2} + |\rho|^3 \right) e^{-c_1(|\rho| + |\sigma|)},$$

and

$$(21) \quad | \Delta R_1(h) | \lesssim \left(\frac{1}{(k + \sqrt{h}\rho)^4} + |\rho|^3 \right) e^{-c_2(|\rho| + |\sigma|)}.$$

Now,

$$\begin{aligned} Eq(v) &= Eq(\chi(\sqrt{h}\rho)w) \\ &= \chi(\sqrt{h}\rho)Eq(w) - h\chi''(\sqrt{h}\rho)w - 2h^{\frac{1}{2}}\chi'(\sqrt{h}\rho)\partial_\rho w \\ &\quad + a\chi(\chi^2 - 1)w^3 \\ &= h^{\frac{5}{2}}\chi(\sqrt{h}\rho)R_1 + R_2 + R_3 + R_4 := R(h). \end{aligned}$$

Set $I = [\frac{1}{2}, \frac{3}{4}] \cup [\frac{5}{4}, \frac{3}{2}]$ and observe that $\text{supp}\chi' \subset I$, $\text{supp}\chi'' \subset I$, $\text{supp}\chi(\chi^2 - 1) \subset I$ and if $\sqrt{h}\rho \in I$ we have

$$|w|, |\partial_\rho w| \lesssim e^{-c/\sqrt{h}} e^{-c|\sigma|},$$

then it follows

$$(22) \quad \| \Delta^j R_p \|_{L^2} \lesssim e^{-c/\sqrt{h}},$$

for all $0 \leq j \leq 1$ and $2 \leq p \leq 4$. According to (21) we also have

$$\| \chi(\sqrt{h}\rho)R_1 \|_{L^2}^2 \lesssim \int (1 + |\rho|^6) e^{-2c_1(|\rho| + |\sigma|)} \leq C.$$

Therefore, coming back in variables (r, y, θ) , $\|R(h)\|_{L^2} \lesssim h^{\frac{5}{2}}$. Because of the fast decay of w we also have $\|(r^2 + y^2)R(h)\|_{L^2} \lesssim h^{\frac{5}{2}}$, hence $\|(|x|^2 + 1) R(h) \|_{L^2} \lesssim h^{\frac{5}{2}}$.

Differentiating u_{app} costs at most h^{-1} , then together with (21) and (22) we obtain $\|\Delta R(h)\|_{L^2} \lesssim h^{\frac{1}{2}}$. \square

PROPOSITION 2.7. — *Let $|a|\kappa^2 \leq c_0$ fixed, let u_{app} be given by (18) and let u be solution of*

$$(23) \quad \begin{cases} ih\partial_t u + h^2\Delta u - |x|^2 u = ah^2|u|^2 u, \\ u(0, x) = u_{app}(0, x), \end{cases}$$

then $\|(u - u_{app})(t_h)\|_{L^2} \rightarrow 0$ with $t_h \ll \log(\frac{1}{h})$, when $h \rightarrow 0$.

Proof. — Denote by $w = u - u_{app}$ and by $f = ah^2g + R(h)$ with $g = |u_{app} + w|^2(u_{app} + w) - |u_{app}|^2 u_{app}$, then

$$(24) \quad ih\partial_t w + h^2\Delta w - |x|^2 w = f.$$

We define

$$(25) \quad E(t) = \int \left(\frac{1}{2}(|x|^4 + 1)|w|^2 + h^4|\Delta w|^2 \right).$$

– Multiply (24) by $\frac{1}{2}(|x|^4 + 1)\bar{w}$, integrate and take the imaginary part:

$$(26) \quad \frac{1}{2}h \frac{d}{dt} \int \frac{1}{2}(|x|^4 + 1)|w|^2 = \text{Im} \int \frac{1}{2}(|x|^4 + 1)f\bar{w} + 2h^2 \text{Im} \int |x|^2 \bar{w} x \nabla w,$$

– Multiply $\Delta(24)$ by $h^4\Delta\bar{w}$, integrate and take the imaginary part:

$$(27) \quad \frac{1}{2}h \frac{d}{dt} \int h^4|\Delta w|^2 = h^4 \text{Im} \int \Delta f \Delta \bar{w} - 2h^4 \text{Im} \int \Delta w x \nabla \bar{w}.$$

With an integration by parts, we can show that

$$h^2 \int |x|^2 |\nabla w|^2 \lesssim \int |x|^4 |w|^2 + h^4 \int |\Delta w|^2,$$

therefore

$$(28) \quad h^2 \left| \int |x|^2 \bar{w} x \nabla w \right| \lesssim h \int |x|^4 |w|^2 + h^3 \int |x|^2 |\nabla w|^2 \lesssim hE,$$

and

$$(29) \quad h^4 \left| \int \Delta w x \nabla \bar{w} \right| \lesssim h^5 \int |\Delta w|^2 + h^3 \int |x|^2 |\nabla w|^2 \lesssim hE.$$

Then the inequalities (26)-(29) yield

$$(30) \quad h \frac{d}{dt} E(t) \lesssim \text{Im} \int \left(\frac{1}{2}(|x|^4 + 1)f\bar{w} + h^2|x|^2 \nabla f \nabla \bar{w} + h^4 \Delta f \Delta \bar{w} \right) + hE.$$

Using the expression of u_{app}

$$(31) \quad \begin{aligned} \|u_{app}\|_{L^2} &\lesssim 1, & \|u_{app}\|_{L^\infty} &\lesssim h^{-\frac{1}{2}}, \\ \|\nabla u_{app}\|_{L^2} &\lesssim h^{-1}, & \|\nabla u_{app}\|_{L^\infty} &\lesssim h^{-\frac{3}{2}}, \end{aligned}$$

and by definition of E

$$(32) \quad \|x\nabla w\|_{L^2} \lesssim h^{-1}E^{\frac{1}{2}}, \quad \|\Delta w\|_{L^2} \lesssim h^{-2}E^{\frac{1}{2}},$$

and the Gagliardo-Nirenberg inequalities in dimension 3 yield

$$(33) \quad \begin{aligned} \|w\|_{L^4} &\lesssim h^{-\frac{3}{4}}E^{\frac{1}{2}}, \quad \|\nabla w\|_{L^4} \lesssim h^{-\frac{7}{4}}E^{\frac{1}{2}}, \\ \|w\|_{L^\infty} &\lesssim \|w\|_{L^2}^{\frac{1}{4}}\|\Delta w\|_{L^2}^{\frac{3}{4}} \lesssim h^{-\frac{3}{2}}E^{\frac{1}{2}}. \end{aligned}$$

– First, the estimates (20) on $R(h)$ give

$$(34) \quad \begin{aligned} &\left| \int \left(\frac{1}{2}(|x|^4 + 1)R(h)\bar{w} + h^4\Delta R(h)\Delta\bar{w} \right) \right| \\ &\lesssim \|(|x|^2 + 1)R(h)\|_{L^2}E^{\frac{1}{2}} + h^2\|\Delta R(h)\|_{L^2}E^{\frac{1}{2}} \\ &\lesssim h^{\frac{5}{2}}E^{\frac{1}{2}}. \end{aligned}$$

– Then, as $g = |u_{app} + w|^2(u_{app} + w) - |u_{app}|^2u_{app}$, and according to (31) and (33)

$$(35) \quad \begin{aligned} \left| \operatorname{Im} \int (|x|^4 + 1)g\bar{w} \right| &\lesssim \int (|x|^4 + 1) (|u_{app}|^2|w|^2 + |u_{app}| |w|^3) \\ &\lesssim \|u_{app}\|_{L^\infty} (\|u_{app}\|_{L^\infty} + \|w\|_{L^\infty}) E \\ &\lesssim h^{-1}E + h^{-2}E^{\frac{3}{2}}. \end{aligned}$$

– Compute

$$\begin{aligned} |\Delta g| &\lesssim |u_{app}|^2|\Delta w| + |u_{app}|\nabla u_{app}|\nabla w| + |\nabla u_{app}|^2|w| \\ &\quad + |u_{app}|\Delta u_{app}|w| + |\Delta u_{app}||w|^2 + |w|^2|\Delta w| + |w|\nabla w|^2, \end{aligned}$$

hence

$$\begin{aligned} \|\Delta g\|_{L^2} &\lesssim \|u_{app}\|_{L^\infty}^2\|\Delta w\|_{L^2} + \|u_{app}\|_{L^\infty}\|\nabla u_{app}\|_{L^\infty}\|\nabla w\|_{L^2} \\ &\quad + \|\nabla u_{app}\|_{L^\infty}^2\|w\|_{L^2} + \|u_{app}\|_{L^\infty}\|\Delta u_{app}\|_{L^\infty}\|w\|_{L^2} \\ &\quad + \|\Delta u_{app}\|_{L^\infty}\|w\|_{L^4}^2 + \|w\|_{L^\infty}^2\|\Delta w\|_{L^2} + \|w\|_{L^2}\|\nabla w\|_{L^4}^2 \\ &\lesssim h^{-3}E^{\frac{1}{2}} + h^{-4}E + h^{-5}E^{\frac{3}{2}}, \end{aligned}$$

then

$$(36) \quad \begin{aligned} h^4 \left| \int \Delta g \Delta\bar{w} \right| &\lesssim h^4\|\Delta g\|_{L^2}\|\Delta w\|_{L^2} \\ &\lesssim h^{-1}E + h^{-2}E^{\frac{3}{2}} + h^{-3}E^2. \end{aligned}$$

Putting the estimates (34), (35), and (36) together with (30), we obtain

$$(37) \quad h \frac{d}{dt}E(t) \lesssim h^{\frac{5}{2}}E^{\frac{1}{2}} + hE + E^{\frac{3}{2}} + h^{-1}E^2.$$

Set $F = E^{\frac{1}{2}}$, then F satisfies $F(0) = 0$ and

$$(38) \quad h \frac{d}{dt} F(t) \lesssim h^{\frac{5}{2}} + hF + F^2 + h^{-1}F^3.$$

As long as $h^{-1}F^3 \lesssim hF$, i.e. for times such that $F \lesssim h$, we can write

$$\frac{d}{dt} F(t) \lesssim h^{\frac{3}{2}} + F.$$

Using Gronwall's inequality, $F \lesssim h^{\frac{3}{2}} e^{Ct}$. The non linear terms in (38) can be removed with the continuity argument for times t_h such that $e^{Ct_h} \lesssim h^{-\frac{1}{2}}$, i.e. $t_h \ll \log(\frac{1}{h})$ and one has $F(t_h) \rightarrow 0$ when $h \rightarrow 0$, hence the result. \square

We are now able to prove Theorem 1.4 and Theorem 1.5.

3. Geometric instability

Let $|a|\kappa^2 \leq c_0$. Consider the function u_{app} defined by (18) associate with κ with $k = 1$ (k will be equal to 1 in all this section).

$$u_{app} = \kappa h^{-\frac{1}{2}} e^{-i\lambda t} e^{i\frac{\theta}{h}} \tilde{v}.$$

Similarly, let the function u'_{app} defined by (18) associate with $\kappa' = \kappa + h^{\frac{1}{2}}$. Then there exists $\lambda' \in \mathbb{R}$ and $\tilde{v}' \in L^2(\mathbb{R}^3)$ such that

$$u'_{app} = (\kappa + h^{\frac{1}{2}}) h^{-\frac{1}{2}} e^{-i\lambda' t} e^{i\frac{\theta}{h}} \tilde{v}'.$$

define the functions $f, f' \in L(\mathbb{R}^3)$ by

$$(39) \quad f = h^{-\frac{1}{2}} e^{i\frac{\theta}{h}} \tilde{v}, \quad f' = h^{-\frac{1}{2}} e^{i\frac{\theta}{h}} \tilde{v}'.$$

Notice that by construction, $\|f\|_{L^2}, \|f'\|_{L^2} \sim 1$.

We now need the following

LEMMA 3.1. — *The functions defined by (39) satisfy*

$$(40) \quad \|f' - f\|_{L^2} \lesssim h^{\frac{1}{2}}.$$

Proof. — To construct f' , we have to solve the system (6)-(8) with $\kappa' = \kappa + h^{\frac{1}{2}}$. We reorganize this system by identifying the powers of h , and as equation (6) remains the same, we deduce (40). \square

Proof of Theorem 1.4 (i). — Denote by u (resp. u') the solution of (23) with initial condition $u_{app}(0)$ (resp. $u'_{app}(0)$). We have

$$(41) \quad \begin{aligned} \|(u' - u)(0)\|_{L^2} &= \|(u'_{app} - u_{app})(0)\|_{L^2} \\ &\leq \kappa \|f' - f\|_{L^2} + \kappa h^{\frac{1}{2}} \|f'\|_{L^2} \lesssim \kappa h^{\frac{1}{2}}, \end{aligned}$$

by Lemma 3.1. The triangle inequality gives

$$\begin{aligned}
\|(u'_{app} - u_{app})(t)\|_{L^2} &\geq \kappa \left| e^{i(\lambda' - \lambda)t} - 1 \right| \|f'\|_{L^2} - \kappa \|f' - f\|_{L^2} - \kappa h^{\frac{1}{2}} \|f'\|_{L^2} \\
(42) \qquad \qquad \qquad &\geq \kappa \left| e^{i(\lambda' - \lambda)t} - 1 \right| - C\kappa h^{\frac{1}{2}}.
\end{aligned}$$

As $(\lambda' - \lambda)t \sim \frac{\sqrt{2}}{2\pi} a \left((\kappa + h^{\frac{1}{2}})^2 - \kappa^2 \right) t \sim \frac{\sqrt{2}}{\pi} a\kappa t h^{\frac{1}{2}}$, with (42) we obtain, when $|a|\kappa t \gg 1$

$$\|(u'_{app} - u_{app})(t)\|_{L^2} \geq c|a|\kappa^2 t h^{\frac{1}{2}},$$

hence, using (41)

$$\frac{\|(u' - u)(t)\|_{L^2}}{\|(u' - u)(0)\|_{L^2}} \gtrsim |a|\kappa t.$$

which was the claim. \square

Proof of Theorem 1.4 (ii). — First notice that every parameter or function involved in this part depends on h even though we do not write the subscripts. We define

$$\begin{aligned}
u''_{app} &= (\kappa + \varepsilon_h) h^{-\frac{1}{2}} e^{-i\lambda'' t} e^{i\frac{\theta}{h}} \tilde{v}'' \\
(43) \qquad \qquad \qquad &:= (\kappa + \varepsilon_h) e^{-i\lambda'' t} f''.
\end{aligned}$$

with $\varepsilon_h \rightarrow 0$ when $h \rightarrow 0$, and denote by u'' the solution of (23) with initial condition $u''_{app}(0)$. Then

$$\begin{aligned}
\|(u'' - u)(0)\|_{L^2} &= \|(u''_{app} - u_{app})(0)\|_{L^2} \\
(44) \qquad \qquad \qquad &\leq \kappa \|f'' - f\|_{L^2} + \kappa \varepsilon_h \|f''\|_{L^2}.
\end{aligned}$$

The right hand side of (44) tends to 0 with h because $\|f'' - f\|_{L^2} \rightarrow 0$ and $\|f''\|_{L^2} \sim 1$. But when h is small enough

$$\begin{aligned}
\|(u''_{app} - u_{app})(t)\|_{L^2} &\geq \kappa \left| e^{i(\lambda'' - \lambda)t} - 1 \right| \|f''\|_{L^2} - \kappa \|f'' - f\|_{L^2} - \kappa \varepsilon_h \|f''\|_{L^2} \\
(45) \qquad \qquad \qquad &\geq \frac{1}{2} \kappa \left| e^{i(\lambda'' - \lambda)t} - 1 \right|.
\end{aligned}$$

Now use $(\lambda'' - \lambda)t_h \sim \frac{\sqrt{2}}{2\pi} a \left((\kappa + \varepsilon_h)^2 - \kappa^2 \right) t_h \sim C_0 a \kappa t_h \varepsilon_h$. Take $\varepsilon_h = (C_0 \kappa a t_h)^{-1/2}$ which tends to 0, then if $h \ll 1$, $|\lambda'' - \lambda|t_h \geq \pi$ and

$$\sup_{0 \leq t \leq t_h} \|(u''_{app} - u_{app})(t)\|_{L^2} \geq \kappa.$$

Now, according to Proposition 2.7, which can be used as we assume $t \ll \log \frac{1}{h}$, we have for h small enough

$$\sup_{0 \leq t \leq t_h} \|(u'' - u)(t)\|_{L^2} \geq \kappa.$$

This last inequality together with (44) proves the second part of Theorem 1.4. \square

4. Projective instability

We conserve the notations of the previous section, but here f_j and f'_j are constructed with $k = j$ in (4).

Define $U_{app} = \kappa e^{-i\lambda_1 t} f_1 + \kappa e^{-i\lambda_2 t} f_2$ and $U'_{app} = (\kappa + \varepsilon_h) e^{-i\lambda'_1 t} f'_1 + \kappa e^{-i\lambda_2 t} f_2$.

LEMMA 4.1. — *Let $V_{app} = U_{app}$ or $V_{app} = U'_{app}$, and v be solution of*

$$(46) \quad \begin{cases} ih\partial_t v + h^2\Delta v - |x|^2 v = ah^2|v|^2 v, \\ v(0, x) = V_{app}(0, x), \end{cases}$$

then $\|(v - V_{app})(t_h)\|_{L^2} \rightarrow 0$ with $t_h \ll \log(\frac{1}{h})$, when $h \rightarrow 0$.

Proof. — Write $V_{app} = v^1_{app} + v^2_{app}$ with $v^1_{app} = \kappa e^{-i\lambda_1 t} f_1$ or $v^1_{app} = (\kappa + \varepsilon_h) e^{-i\lambda'_1 t} f'_1$ and $v^2_{app} = \kappa e^{-i\lambda_2 t} f_2$. As the supports of v^1_{app} and v^2_{app} are disjoint we have

$$\begin{aligned} & ih\partial_t(v^1_{app} + v^2_{app}) + h^2\Delta(v^1_{app} + v^2_{app}) - |x|^2(v^1_{app} + v^2_{app}) \\ &= ah^2(|v^1_{app}|^2 v^1_{app} + |v^2_{app}|^2 v^2_{app}) + R^1(h) + R^2(h) \\ &= ah^2|v^1_{app} + v^2_{app}|^2(v^1_{app} + v^2_{app}) + R^1(h) + R^2(h), \end{aligned}$$

where for $j = 1, 2$, $R^j(h)$ is the error term given by Proposition 2.6 and therefore satisfies $\|(|x|^2 + 1)R^j(h)\|_{L^2} \lesssim h^{\frac{5}{2}}$ and $\|\Delta R^j(h)\|_{L^2} \lesssim h^{\frac{1}{2}}$. We conclude with the help of Proposition 2.7. \square

Proof of Theorem 1.5. — Consider the function u (resp. u') the solution of equation (46) with Cauchy data $U_{app}(0)$ (resp. $U'_{app}(0)$).

First notice that, for $t \geq 0$, $\|V_{app}(t)\|_{L^2}^2 \sim 2\kappa^2$. Compute

$$(47) \quad U_{app}(t)\overline{U'_{app}(t)} = \kappa(\kappa + \varepsilon_h) f_1 \overline{f'_1} e^{i(\lambda'_1 - \lambda_1)t} + \kappa^2 |f_2|^2.$$

Then for $t = 0$ we have

$$\int U_{app} \overline{U'_{app}}(0) \sim 2\kappa^2,$$

hence

$$d_{\text{pr}}(u(0), u'(0)) = d_{\text{pr}}(U_{app}(0), U'_{app}(0)) \rightarrow 0.$$

Let $t_h \ll \log \frac{1}{h}$, then as $(\lambda'_1 - \lambda_1)t_h \sim C_0 a \kappa \varepsilon_h t_h$, we now choose

$$\varepsilon_h = \frac{\pi}{C_0 a \kappa t_h},$$

then we have $(\lambda'_1 - \lambda_1)t_h \rightarrow \pi$, as $h \rightarrow 0$. Thus

$$\int U_{app} \overline{U'_{app}}(t_h) \rightarrow 0,$$

and

$$d_{\text{pr}}(U_{app}(t_h), U'_{app}(t_h)) \rightarrow \arccos(0) = \frac{\pi}{2}.$$

Finally, from Lemma 4.1 we deduce

$$d_{\text{pr}}(u(t_h), U_{\text{app}}(t_h)), d_{\text{pr}}(u'(t_h), U'_{\text{app}}(t_h)) \longrightarrow 0,$$

and therefore

$$\begin{aligned} d_{\text{pr}}(u(t_h), u'(t_h)) &\geq d_{\text{pr}}(U_{\text{app}}(t_h), U'_{\text{app}}(t_h)) - d_{\text{pr}}(u(t_h), U_{\text{app}}(t_h)) \\ &\quad - d_{\text{pr}}(u'(t_h), U'_{\text{app}}(t_h)) \\ &\geq \frac{\pi}{4}, \end{aligned}$$

for $h \ll 1$; hence the result. \square

BIBLIOGRAPHY

- [1] N. Burq, P. Gérard, and N. Tzvetkov. An instability property of the nonlinear Schrödinger equation on S^d . *Math. Res. Lett.*, 9(2-3):323–335, 2002.
- [2] N. Burq and M. Zworski. Instability for the semiclassical nonlinear Schrödinger equation. *Comm. Math. Phys.* 260 no. 1, 45–58, 2005.
- [3] R. Carles. Geometric optics and instability for semi-classical Schrödinger equations. To appear in *Arch. Ration. Mech.*
- [4] M. Christ, J. Colliander, and T. Tao. Asymptotics, frequency modulation, and low regularity ill-posedness for canonical defocusing equations. *Amer. J. Math.*, 125(6):1235–1293, 2003.
- [5] M. Christ, J. Colliander, and T. Tao. Ill-posedness for nonlinear Schrödinger and wave equation. To appear in *Ann. Inst. H. Poincaré Anal. Non Linéaire*.
- [6] B. Helffer. *Semi-classical analysis for the Schrödinger operator and applications*, volume 1336 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1988.
- [7] L. Pitaevskii and S. Stringari. *Bose-Einstein condensation*, volume 116 of *International Series of Monographs on Physics*. The Clarendon Press Oxford University Press, Oxford, 2003.
- [8] M. Reed and B. Simon. *Methods of modern mathematical physics. IV. Analysis of operators*. Academic Press [Harcourt Brace Jovanovich Publishers], New York, 1978.