

**COMPACTNESS AND GLOBAL ESTIMATES  
FOR THE GEOMETRIC PANEITZ EQUATION  
IN HIGH DIMENSIONS**

EMMANUEL HEBEY AND FRÉDÉRIC ROBERT

ABSTRACT. Given  $(M, g)$  a smooth compact Riemannian manifold of dimension  $n \geq 5$ , we investigate compactness for the fourth order geometrical equation  $P_g u = u^{2^\sharp - 1}$ , where  $P_g$  is the Paneitz operator, and  $2^\sharp = 2n/(n-4)$  is critical from the Sobolev viewpoint. We prove that the equation is compact when the Paneitz operator is of strong positive type.

In 1983, Paneitz [9] introduced a conformally fourth order operator defined on 4-dimensional Riemannian manifolds. Branson [3] generalized the definition to  $n$ -dimensional Riemannian manifolds,  $n \geq 5$ . While the conformal Laplacian is associated to the scalar curvature, the geometric Paneitz-Branson operator is associated to a notion of  $Q$ -curvature. The  $Q$ -curvature in dimension 4, and for conformally flat manifolds, is the integrand in the Gauss-Bonnet formula for the Euler characteristic. In this article we let  $(M, g)$  be a smooth compact conformally flat Riemannian  $n$ -manifold,  $n \geq 5$ , and consider the geometric Paneitz equation

$$P_g u = u^{2^\sharp - 1}, \quad (0.1)$$

where  $P_g$  is the Paneitz operator in dimension  $n \geq 5$ ,  $u$  is required to be positive, and  $2^\sharp = \frac{2n}{n-4}$  is the critical exponent for the Sobolev embedding. The Paneitz operator in dimension  $n \geq 5$  reads as

$$P_g u = \Delta_g^2 u - \operatorname{div}_g (A_g du) + \frac{n-4}{2} Q_g u,$$

where  $\Delta_g = -\operatorname{div}_g \nabla$  is the Laplace-Beltrami operator,  $Q_g$  is the  $Q$ -curvature of  $g$ ,  $A_g$  is the smooth symmetrical  $(2, 0)$ -tensor field given by

$$A_g = \frac{(n-2)^2 + 4}{2(n-1)(n-2)} S_g g - \frac{4}{n-2} Rc_g,$$

and  $Rc_g$  and  $S_g$  are respectively the Ricci curvature and scalar curvature of  $g$ . The Paneitz operator is conformally invariant in the sense that if  $\tilde{g} = u^{4/(n-4)} g$  is conformal to  $g$ , then  $P_{\tilde{g}}(f) = u^s P_g(uf)$  for all  $f \in C^\infty(M)$ , where  $s = 1 - 2^\sharp$ . From the viewpoint of conformal geometry, equation (0.1) turns out to be the natural fourth order analogue of the second order Yamabe equation. We refer to Chang [4] and Chang and Yang [5] for more details on the above definitions.

In what follows we let  $H_2^2(M)$  be the Sobolev space consisting of functions in  $L^2(M)$  with two derivatives in  $L^2$ . As shown in Hebey and Robert [7], up to passing to a subsequence, bounded sequences  $(u_\alpha)$  in  $H_2^2(M)$  of nonnegative solutions of (0.1) split into the sum of a nonnegative solution  $u^0$ , namely the weak limit of

---

*Date:* October 7, 2004.

the  $u_\alpha$ 's, a finite sum of  $k$  bubbles  $(B_\alpha^i)$ , obtained by rescaling positive solutions of the Euclidean equation  $\Delta^2 u = u^{2^* - 1}$ , and a rest  $R_\alpha$  which converges strongly to zero in  $H_2^2(M)$  as  $\alpha \rightarrow +\infty$ . This splitting provides exact asymptotics for the  $u_\alpha$ 's in the Sobolev setting. Following standard terminology, we say that equation (0.1) is compact if for any bounded sequence  $(u_\alpha)$  in  $H_2^2(M)$  of nonnegative solutions of (0.1), we necessarily have that  $k = 0$  in such decompositions. Regularity theory, see for instance Esposito-Robert [6], holds for (0.1). Then, thanks to Agmon-Douglis-Nirenberg-type estimates [1, 2], and estimates like the ones developed in Hebey, Robert and Wen [8], an equivalent definition for compactness is that bounded sequences in  $H_2^2(M)$  of nonnegative solutions of (0.1) are actually bounded in  $C^{4,\theta}(M)$ . A major stress in studying compactness is to understand large solutions. Namely, solutions with large energies which, in studying their possible blow-up, involve multi-bubbles which may interact one with another on the pointwise level.

Compactness for fourth order equations like (0.1) was recently studied in Hebey, Robert and Wen [8]. It was shown in [8] that equations like (0.1) are compact as long as they are not close to the geometric equation (0.1). In this note we investigate compactness for the geometric equation (0.1) and show how the analysis developed in [8] provides an answer to this question of whether (0.1) is compact or not. The result we get is the fourth order analogue of the result proved in Schoen [10] where the second order Yamabe equation was investigated. As in Hebey, Robert and Wen [8], and also Schoen [10], we assume in what follows that  $(M, g)$  is conformally flat (and hence, since  $n \geq 5$ , that the Weyl tensor of  $g$  is zero). We let  $G_g$  be the Green's function of  $P_g$ . The Green's function is unique if  $P_g$  is positive. By conformal invariance of the Paneitz operator, if  $\tilde{g} = u^{4/(n-4)}g$  is a conformal metric to  $g$ , then

$$G_{\tilde{g}}(x, y) = \frac{G_g(x, y)}{u(x)u(y)}. \quad (0.2)$$

It is known that if  $\tilde{g}$  is a flat metric around some  $x_0 \in M$ , then

$$G_{\tilde{g}}(x_0, x) = \frac{\lambda_n}{d_{\tilde{g}}(x_0, x)^{n-4}} + \mu_{\tilde{g}}(x_0, x), \quad (0.3)$$

where  $\lambda_n^{-1} = 2(n-2)(n-4)\omega_{n-1}$ ,  $\omega_{n-1}$  is the volume of the unit  $(n-1)$ -sphere, and the function  $x \rightarrow \mu_{\tilde{g}}(x_0, x)$  is smooth on  $M$ . Combining the above two equations, noting that conformal changes of metrics which leave a metric flat around one point come from conformal diffeomorphisms of the Euclidean space, we easily get (as in Schoen and Yau [11] for the conformal Laplacian) that if  $g$  and  $\tilde{g} = u^{4/(n-4)}g$  are conformal metrics, both being flat around  $x_0$ , then

$$\mu_{\tilde{g}}(x_0, x_0) = \frac{\mu_g(x_0, x_0)}{u(x_0)^2}. \quad (0.4)$$

In particular, by (0.4), the sign of  $\mu_g(x_0) = \mu_g(x_0, x_0)$  does not depend on the choice of the metric in  $[g]_{x_0}$ , where  $[g]_{x_0}$  stands for the set of conformal metrics to  $g$  which are flat around  $x_0$ .

In what follows we say that  $P_g$  is of *strong positive type* if  $P_g$  is positive,  $G_g$  is positive, and for any  $x \in M$  there exists  $\tilde{g} \in [g]_x$  such that  $\mu_{\tilde{g}}(x) > 0$ . For example, the Paneitz operator on quotients of the unit sphere is of strong positive

type. Positivity of the Paneitz operator was studied in Xu and Yang [12]. The main result of this note is:

**Theorem 0.1.** *The geometric equation (0.1) is compact on compact conformally flat manifolds of dimensions  $n \geq 5$  with Paneitz operator of strong positive type.*

Let  $(M, g)$  be a smooth compact conformally flat manifold of dimension  $n \geq 5$  with positive Paneitz operator  $P_g$ , and positive Green's function  $G_g$ . Given  $S \subset M$ , let  $[g]_S$  be the set of conformal metrics to  $g$  which are flat in a neighborhood of  $S$ . We prove Theorem 0.1 by proving that if  $(u_\alpha)$  is a bounded sequence in  $H_2^2(M)$  of nonnegative solutions of (0.1) which blows up with geometrical blow-up points  $S = \{x_1, \dots, x_N\}$ , then  $u_\alpha \rightarrow 0$  in  $H_2^2(M)$  as  $\alpha \rightarrow +\infty$ , and for any  $\tilde{g} \in [g]_S$ , there exist  $\lambda_{i,j} > 0$  such that for any  $i = 1, \dots, N$ ,

$$\mu_{\tilde{g}}(x_i) + \sum_{j \neq i} \lambda_{i,j} G_{\tilde{g}}(x_i, x_j) = 0. \quad (0.5)$$

Theorem 0.1 clearly follows from (0.5). We prove equation (0.5) in the rest of this note, using the material proved in Hebey, Robert and Wen [8].

## 1. PROOF OF THE RESULT

In what follows we prove (0.5), and thus Theorem 0.1. For that purpose we let  $(u_\alpha)$  be a bounded sequences in  $H_2^2(M)$  of nonnegative nontrivial solutions of (0.1). Since the Green's function  $G_g$  of  $P_g$  is positive, the  $u_\alpha$ 's are positive. In the sequel, everything is up to a subsequence. We know from Hebey and Robert [7] that there exist  $k \in \mathbb{N}$ ,  $u^0 \geq 0$  a nonnegative solution of (0.1), and  $k$  bubbles  $(B_\alpha^i)$ ,  $i = 1, \dots, k$ , such that

$$u_\alpha = u^0 + \sum_{i=1}^k B_\alpha^i + R_\alpha, \quad (1.1)$$

where  $R_\alpha \rightarrow 0$  in  $H_2^2(M)$  as  $\alpha \rightarrow +\infty$ . By contradiction, we assume that  $k \geq 1$ , and let  $S = \{x_1, \dots, x_N\}$  be the geometric blow-up points set consisting of the limits of the centers of the bubbles  $(B_\alpha^i)$ . Since bubbles may accumulate one on another (there are such examples for equations like (0.1), we refer to Hebey, Robert and Wen [8]),  $N$  might be less than  $k$ . By conformal invariance we may assume that  $g = \tilde{g}$  is flat around the points in  $S$ . Then the geometric Paneitz equation (0.1) reduces to  $\Delta_g^2 u_\alpha = u_\alpha^{2^* - 1}$  around the points in  $S$ .

A preliminary step in the proof of (0.5), we come back to the estimates proved in Hebey, Robert and Wen [8] and explain why they are still valid in the present context. A rough argument would be that blow-up phenomena are local in nature, while the Paneitz operator on conformally flat manifolds is locally, up to conformal changes of the metric, like the Paneitz operator on the sphere (and hence with positive constant coefficients as in [8]). More details are as follows. First we note that the standard procedure to get rescaling invariant pointwise estimates, as developed in [8] for fourth order operators, together with Agmon-Douglis-Nirenberg-type estimates for fourth order operators, give that there exists  $C > 0$  such that,

$$\left( \min_{1 \leq i \leq k} d_g(x_\alpha^i, x) \right)^{\frac{n-4}{2}} |u_\alpha(x) - u^0(x)| \leq C \quad (1.2)$$

for all  $\alpha$  and all  $x$ , where  $u^0$ , the weak limit of the  $u_\alpha$ 's, is as in (1.1), and where the  $x_\alpha^i$ 's,  $i = 1, \dots, k$ , are the centers of the bubbles ( $B_\alpha^i$ ) in (1.1). In particular, we get with (1.2) that  $u_\alpha \rightarrow u^0$  in  $C_{loc}^4(M \setminus S)$  as  $\alpha \rightarrow +\infty$ . In addition to (1.2), if  $\Phi_\alpha(x)$  stands for the left hand side in (1.2), we also have that

$$\lim_{R \rightarrow +\infty} \lim_{\alpha \rightarrow +\infty} \sup_{x \in M \setminus \Omega_\alpha(R)} \Phi_\alpha(x)^{\frac{n-4}{2}} = 0, \quad (1.3)$$

where, for  $R > 0$ ,  $\Omega_\alpha(R)$  is given by  $\Omega_\alpha(R) = \bigcup_{i=1}^k B_{x_\alpha^i}(R\mu_\alpha^i)$ , and the  $\mu_\alpha^i$ 's are the weights of the bubbles ( $B_\alpha^i$ ) in (1.2). Going on with the estimates in Hebey, Robert and Wen [8] we may assume, up to renumbering, and up to passing to a subsequence, that ( $B_\alpha^1$ ) is the bubble in (1.1) with the largest weight. Then we let the  $x_\alpha$ 's and  $\mu_\alpha$ 's be such that  $x_\alpha = x_\alpha^1$  and  $\mu_\alpha = \mu_\alpha^1$  for all  $\alpha$ . A preliminary remark is that the global splitting estimate  $\|u_\alpha\|_{p_1, p_2, \mu_\alpha^{-1}} \leq C$  in [8] easily follows from the positivity of the Green's function with only slight modifications of the arguments in [8]. The arguments in [8] used the decomposition of the fourth order operator into the product of two second order operators. We may use instead Agmon-Douglis-Nirenberg-type estimates and note that the positivity of the Green's function  $G_g$  implies that  $P_g$  satisfies the comparison principle. As an independent easy remark, since  $\Delta_g(\Delta_g u_\alpha) \geq 0$  around the points in  $S$ , there exists  $C > 0$  such that  $\Delta_g u_\alpha \geq -C$  in  $M$ , for all  $\alpha$ . Then, with such an estimate, and the analysis developed in [8], we easily get that the integral and asymptotic estimates in Hebey, Robert and Wen [8] follow from (1.1)–(1.3). In several places the computations in [8] simplify because of the simple nature of the geometric equation around the points in  $S$ . By the asymptotic estimates, if we let  $\tilde{u}_\alpha$  be the rescaled function obtained from  $u_\alpha$  by  $\tilde{u}_\alpha(x) = u_\alpha(\exp_{x_\alpha}(\sqrt{\mu_\alpha}x))$ , we get that there exist  $\delta > 0$ ,  $A > 0$ , and a biharmonic function  $\varphi \in C^4(B_0(2\delta))$  such that, up to a subsequence,

$$\tilde{u}_\alpha(x) \rightarrow \frac{A}{|x|^{n-4}} + \varphi(x) \quad (1.4)$$

in  $C_{loc}^3(B_0(2\delta) \setminus \{0\})$  as  $\alpha \rightarrow +\infty$ . Moreover, since we assumed that  $G_g > 0$ , so that either  $u^0 \equiv 0$  or  $u^0 > 0$  everywhere, we also have the important information that  $\varphi$  is positive in  $B_0(2\delta)$  if  $u^0 \not\equiv 0$ .

As an important step in the proof of (0.5), we claim that thanks to the asymptotics (1.4), and thanks to the property that  $\varphi$  is positive if  $u^0$  is nonzero, we necessarily have that  $u^0 \equiv 0$  when the  $u_\alpha$ 's blow up. In order to do this, we use the Pohozaev identity as in [8], and conformal invariance. The Pohozaev identity for fourth order equations reads as

$$\begin{aligned} & \int_{\Omega} (x^k \partial_k u) \Delta^2 u dx + \frac{n-4}{2} \int_{\Omega} u \Delta^2 u dx \\ &= \frac{n-4}{2} \int_{\partial\Omega} \left( -u \frac{\partial \Delta u}{\partial \nu} + \frac{\partial u}{\partial \nu} \Delta u \right) d\sigma \\ &+ \int_{\partial\Omega} \left( \frac{1}{2} (x, \nu) (\Delta u)^2 - (x, \nabla u) \frac{\partial \Delta u}{\partial \nu} + \frac{\partial (x, \nabla u)}{\partial \nu} \Delta u \right) d\sigma \end{aligned} \quad (1.5)$$

for all smooth bounded domains  $\Omega$  in  $\mathbb{R}^n$  and all  $u \in C^4(\overline{\Omega})$ , where  $\Delta$  is the Euclidean Laplacian,  $\nu$  is the outward unit normal of  $\partial\Omega$ , and  $d\sigma$  is the Euclidean volume element on  $\partial\Omega$ . We apply the Pohozaev identity (1.5) to the  $u_\alpha$ 's in the ball

$\Omega = B_0(\delta\sqrt{\mu_\alpha})$ . In the process we assimilate  $x_\alpha$  and 0 (thanks to the exponential map at  $x_\alpha$ ), and regard  $u_\alpha$  as a function in the Euclidean space. Noting that

$$\int_{B_\alpha} (x^k \partial_k u_\alpha) \Delta^2 u_\alpha dx + \frac{n-4}{2} \int_{B_\alpha} u_\alpha \Delta^2 u_\alpha dx = O\left(\int_{\partial B_\alpha} u_\alpha^{2^\sharp} d\sigma\right),$$

where  $B_\alpha = B_0(\delta\sqrt{\mu_\alpha})$ , and that  $\int_{\partial B_\alpha} u_\alpha^{2^\sharp} d\sigma = o(\mu_\alpha^{(n-4)/2})$  by (1.4), we get with (1.4), the Pohozaev identity, and the computations developed in [8], that

$$(n-2)(n-4)^2 \omega_{n-1} A \varphi(0) \mu_\alpha^{\frac{n-4}{2}} + o\left(\mu_\alpha^{\frac{n-4}{2}}\right) = 0,$$

where  $\omega_{n-1}$  is the volume of the unit  $(n-1)$ -sphere, and  $A$  and  $\varphi$  are as in (1.4). In particular,  $\varphi(0) = 0$ , and since  $\varphi > 0$  if  $u^0 \not\equiv 0$ , this proves the above claim that we necessarily have that  $u^0 \equiv 0$  when the  $u_\alpha$ 's blow up. With respect to the terminology in Hebey, Robert and Wen [8], this amounts to say that compactness reduces to pseudo-compactness for the geometric equation.

Going on with the proof of (0.5), and now that we know that  $u^0 \equiv 0$ , we need to add one important estimate to the estimates listed above we proved in [8]. We claim here that

$$\lambda_\alpha u_\alpha(x) \rightarrow \sum_{i=1}^N \lambda_i G_g(x_i, x) \quad (1.6)$$

in  $C_{loc}^4(M \setminus S)$  as  $\alpha \rightarrow +\infty$ , where  $\lambda_\alpha \rightarrow +\infty$  as  $\alpha \rightarrow +\infty$ ,  $S = \{x_1, \dots, x_N\}$  is the geometric blow-up points set of the  $u_\alpha$ 's,  $G_g$  is the Green's function of  $P_g$ , and the  $\lambda_i \geq 0$ ,  $i = 1, \dots, N$ , are such that  $\sum_{i=1}^N \lambda_i = 1$ . In order to prove (1.6), we use the positivity of  $P_g$  and  $G_g$  as follows. By the positivity of  $P_g$ , the lowest eigenvalue  $\lambda$  of  $P_g$  is positive. If  $\psi$  is an eigenfunction for  $\lambda$ , letting  $P_g u = |P_g \psi|$ , and writing that  $P_g u \geq P_g \psi$  and  $P_g u \geq -P_g \psi$ , we get with the positivity of  $G_g$  that  $u \geq |\psi|$ . Noting that  $u > 0$  since  $P_g u = \lambda |\psi|$  and  $G_g > 0$ , plugging  $u$  into the Rayleigh characterization of  $\lambda$ , it follows that either  $\psi < 0$  or  $\psi > 0$ . Without loss of generality we may assume that  $\psi > 0$ . Then the conformal metric  $\tilde{g} = \psi^{4/(n-4)} g$  is such that  $Q_{\tilde{g}} > 0$ , where  $Q_{\tilde{g}}$  is the  $Q$ -curvature of  $\tilde{g}$ . By conformal invariance,  $u = \psi^{-1} u_\alpha$  solves the equation  $P_{\tilde{g}} u = u^{2^\sharp-1}$ . Integrating over  $M$ , since  $Q_{\tilde{g}} > 0$ , we get that there exists  $C > 0$  such that  $\|u_\alpha\|_{L^1(M)} \leq C \|u_\alpha\|_{L^s(M)}^s$  where  $s = 2^\sharp - 1$ . By Agmon-Douglis-Nirenberg-type estimates, noting that (1.2) gives that the  $u_\alpha$ 's are bounded in  $C_{loc}^0(M \setminus S)$  as  $\alpha \rightarrow +\infty$ , we get that for any  $p > 1$ , and any  $\delta > 0$ , the  $L^\infty$ -norm of the  $u_\alpha$ 's in sets like  $M \setminus B_\delta$  is controlled by the  $L^p$ -norm of the  $u_\alpha$ 's in  $M$ , where  $B_\delta$  is the union over the  $x \in S$  of the geodesic balls  $B_x(\delta)$ . By the above estimate, using Hölder's inequality with  $1 \leq p \leq 2^\sharp$ , choosing  $p > 1$  close to 1, and since  $u_\alpha \rightarrow 0$  in  $L^q(M)$  for  $q < 2^\sharp$ , it easily follows that  $\|u_\alpha\|_{L^s(M \setminus B_\delta)} = o(\|u_\alpha\|_{L^s(M)})$  for all  $\delta > 0$ , where  $s$  is as above. In particular, the  $\lambda_i$ 's given by

$$\lambda_i = \lim_{\alpha \rightarrow +\infty} \frac{\int_{B_{x_i}(\delta)} u_\alpha^{2^\sharp-1} dv_g}{\int_M u_\alpha^{2^\sharp-1} dv_g}$$

are nonnegative, independent of  $\delta > 0$  small, and such that  $\sum \lambda_i = 1$ . In what follows we let  $\lambda_\alpha = \|u_\alpha\|_{2^\sharp-1}^{1-2^\sharp}$ . Then  $\lambda_\alpha \rightarrow +\infty$  as  $\alpha \rightarrow +\infty$ , while we can write

with the Green's representation formula that for  $x \in M \setminus B_\delta$ , and  $0 < \delta' \ll \delta$ ,

$$\begin{aligned} u_\alpha(x) &= \int_{B_{\delta'}} G_g(x, y) u_\alpha^{2^\sharp - 1}(y) dv_g(y) + \int_{M \setminus B_{\delta'}} G_g(x, y) u_\alpha^{2^\sharp - 1}(y) dv_g(y) \\ &= \left( \sum_{i=1}^N \lambda_i G_g(x_i, x) + o_{\delta'}(1) \right) \lambda_\alpha^{-1}, \end{aligned}$$

where  $\lim_{\delta' \rightarrow 0} \lim_{\alpha \rightarrow +\infty} o_{\delta'}(1) = 0$ . In particular,  $\lambda_\alpha u_\alpha(x) \rightarrow \sum_{i=1}^N \lambda_i G_g(x_i, x)$  in  $C_{loc}^0(M \setminus S)$  as  $\alpha \rightarrow +\infty$ , an equation from which we easily get that (1.6) is true.

With (1.6) we can now end the proof of (0.5). If  $f$  stands for the function in the right hand side of (1.6), then  $\Delta^2 f = 0$  in a set like  $\Omega = \bigcup_{i=1}^N B_{x_i}(\delta_0) \setminus S$  where  $\delta_0 > 0$ . We apply the Pohozaev identity (1.5) to the  $u_\alpha$ 's in  $B_{x_i}(\delta)$  for  $\delta > 0$  small and  $i$  in  $\{1, \dots, N\}$ . In the process we assimilate  $x_i$  and 0 (thanks to the exponential map at  $x_i$ ), and regard  $u_\alpha$  as a function in the Euclidean space. Noting that

$$\int_B (x^k \partial_k u_\alpha) \Delta^2 u_\alpha dx + \frac{n-4}{2} \int_B u_\alpha \Delta^2 u_\alpha dx = O\left(\int_{\partial B} u_\alpha^{2^\sharp} d\sigma\right),$$

where  $B = B_0(\delta)$ , it follows from (1.6) and the Pohozaev identity that

$$\begin{aligned} &\frac{n-4}{2} \int_{\partial B_0(\delta)} \left( -f \frac{\partial \Delta f}{\partial \nu} + \frac{\partial f}{\partial \nu} \Delta f \right) d\sigma \\ &+ \int_{\partial B_0(\delta)} \left( \frac{1}{2} (x, \nu) (\Delta f)^2 - (x, \nabla f) \frac{\partial \Delta f}{\partial \nu} + \frac{\partial (x, \nabla f)}{\partial \nu} \Delta f \right) d\sigma = 0. \end{aligned} \quad (1.7)$$

By (0.3), and (1.6), we can write that

$$f(x) = \frac{\hat{\lambda}_i}{|x - x_i|^{n-4}} + R_i(x)$$

for  $x \neq x_i$  close to  $x_i$ , where  $\lambda_i$  is as in (1.6),  $\hat{\lambda}_i = \lambda_n \lambda_i$ ,  $\lambda_n$  is as in (0.3),  $R_i$  is smooth around  $x_i$ , and

$$R_i(x_i) = \lambda_i \mu_g(x_i) + \sum_{j \neq i} \lambda_j G_g(x_j, x_i).$$

Plugging these equations into (1.7), letting  $\delta \rightarrow 0$ , we get that  $R_i(x_i) = 0$ . This equation holds for all  $i = 1, \dots, N$ , and we assumed that  $G_g > 0$ . It follows that  $\lambda_i > 0$  for all  $i$ . This ends the proof of (0.5) and of Theorem 0.1.

As a remark, Theorem 0.1 still holds if we replace in (0.1) the critical exponent  $2^\sharp$  by  $2^\sharp - p_\alpha$  where  $p_\alpha \geq 0$  is such that  $p_\alpha \rightarrow 0$  as  $\alpha \rightarrow +\infty$ . In this case the left hand side in (0.5) is not anymore zero but nonpositive. Needless to say, compactness of the subcritical equations provides a minimizing solution of (0.1).

## REFERENCES

- [1] Agmon, S., Douglis, A. and Nirenberg, L., Estimates near the boundary for solutions of elliptic partial differential equations satisfying general boundary conditions. I, *Comm. Pure Appl. Math.*, 12, 623–727, 1959.
- [2] Agmon, S., Douglis, A. and Nirenberg, L., Estimates near the boundary for solutions of elliptic partial differential equations satisfying general boundary conditions. II, *Comm. Pure Appl. Math.*, 17, 35–92, 1964.
- [3] Branson, T.P., Group representations arising from Lorentz conformal geometry, *J. Funct. Anal.*, 74, 199–291, 1987.

- [4] Chang, S.Y.A., On Paneitz operator - a fourth order differential operator in conformal geometry, Harmonic Analysis and Partial Differential Equations, Essays in honor of Alberto P. Calderon, Eds: M. Christ, C. Kenig and C. Sadosky, Chicago Lectures in Mathematics, 1999, 127–150.
- [5] Chang, S.Y.A. and Yang, P.C., On a fourth order curvature invariant, *Comp. Math.* 237, Spectral Problems in Geometry and Arithmetic, Ed: T. Branson, AMS, 9–28, 1999.
- [6] Esposito, P., Robert, F., Mountain pass critical points for Paneitz-Branson operators, *Calc. Var. Partial Differential Equations*, 15, 2002, 493–517.
- [7] Hebey, E., and Robert, F., Coercivity and Struwe’s compactness for Paneitz type operators with constant coefficients, *Calc. Var. Partial Differential Equations*, 13, 491–517, 2001.
- [8] Hebey, E., Robert, F., and Wen, Y., Compactness and global estimates for a fourth order equation of critical Sobolev growth arising from conformal geometry, *Preprint*, 2004.
- [9] Paneitz, S., A quartic conformally covariant differential operator for arbitrary pseudo-Riemannian manifolds, *Preprint*, 1983.
- [10] Schoen, R., Variational theory for the total scalar curvature functional for Riemannian metrics and related topics, Topics in Calculus of Variations (Montecatini Terme, 1987), *Lecture Notes in Math.*, vol. 1365, Springer-Verlag, Berlin, pp. 120–154, 1989.
- [11] Schoen, R., and Yau, S.T., Conformally flat manifolds, Kleinian groups and scalar curvature, *Invent. Math.*, 92, 47–71, 1988.
- [12] Xu, X., and Yang, P., Positivity of Paneitz operators, *Disc. Cont. Dynamical Systems*, 7, 2001, 329–342.

EMMANUEL HEBEY, UNIVERSITÉ DE CERGY-PONTOISE, DÉPARTEMENT DE MATHÉMATIQUES,  
SITE DE SAINT-MARTIN, 2 AVENUE ADOLPHE CHAUVIN, 95302 CERGY-PONTOISE CEDEX, FRANCE  
*E-mail address:* Emmanuel.Hebey@math.u-cergy.fr

FRÉDÉRIC ROBERT, LABORATOIRE J.A.DIEUDONNÉ, UNIVERSITÉ DE NICE SOPHIA-ANTIPOLIS,  
PARC VALROSE, 06108 NICE CEDEX 2, FRANCE  
*E-mail address:* frobert@math.unice.fr