

Polynomials with only rational roots

L. Hajdu

University of Debrecen

Journées Arithmétiques 2023

Université de Lorraine, Nancy

3-7 July 2023

Plan of the talk

- Introduction and motivation
- New results
 - sharp bounds for the degree in terms of the height
 - sharp bound for the degree if the coeffs are coprime to 6
 - finiteness and full description for fixed degree if the coeffs are composed of a fixed finite set of primes
- Open problems

The new results presented are joint with **R. Tijdeman** and **N. Varga**.

Introduction and motivation

Polynomials in $\mathbb{Z}[x]$ with only rational roots are the simplest examples of decomposable polynomials and forms. Such polynomials play an important role in the theory of Diophantine equations. (See e.g. results of **Evertse and Győry**.)

There is also an extensive literature on polynomials with restricted coefficients, in particular, with coefficients belonging to one of the sets $\{-1, 1\}$, $\{0, 1\}$ or $\{-1, 0, 1\}$. (See e.g. results concerning Littlewood polynomials and Newman polynomials.)

The set of polynomials $f(x) \in \mathbb{Z}[x]$ with all coefficients in $\{-1, 0, 1\}$, constant term non-zero and only rational roots is very restricted. The degree of f is at most 3, an example is

$$f(x) = x^3 - x^2 - x + 1 = (x - 1)^2(x + 1).$$

Theorem 1 (Tijdeman, Varga, H. (202?))

Let $f(x) \in \mathbb{Z}[x]$ be a polynomial of degree n with only non-zero rational roots and height bounded by $H \geq 2$. Then we have both

$$n \leq \left(\frac{2}{\log 2} + o(1) \right) \log H \quad (H \rightarrow \infty) \quad (1)$$

and

$$n \leq \frac{5}{\log 2} \log H. \quad (2)$$

Further, the constants $2/\log 2$ and $5/\log 2$ in (1) and (2), respectively, are best possible.

Sharp bounds for the degree in terms of the height

Remarks. For any $f \in \mathbb{Z}[x]$ of degree n , the height of $g := x^m f(x)$ is the same as that of f , while $\deg(g) = m + n$. So the assumption that the roots of f are non-zero is clearly necessary.

Several authors have considered upper bounds for the number r of real roots of $f(x) \in \mathbb{R}[x]$. (See e.g. results of **Bloch and Pólya**, **E. Schmidt**, **Schur**, **Erdős and Turán**, **Littlewood and Offord**, **Borwein**, **Erdélyi and Kós**.)

For example, a result of **Schur** implies for polynomials $f(x) \in \mathbb{Z}[x]$ with only real roots that

$$n \leq (4 + o(1)) \log H \quad (H \rightarrow \infty).$$

Proof of Theorem 1

On the one hand, let $f(x) = \sum_{j=0}^n a_j x^j$. Then

$$|f(i)| \leq \left| \sum_{j \text{ is even}} |a_j| + i \sum_{j \text{ is odd}} |a_j| \right| \leq \sqrt{\frac{1}{2}n^2 + n + 1} H. \quad (3)$$

On the other hand, we may write $f(x) = \prod_{j=1}^n (q_j x - p_j)$ with $p_j, q_j \in \mathbb{Z}_{\neq 0}$ for all j . Then

$$|f(i)| = \left| \prod_{j=1}^n (q_j i - p_j) \right| = \prod_{j=1}^n \sqrt{q_j^2 + p_j^2} \geq (\sqrt{2})^n. \quad (4)$$

Therefore,

$$n \log 2 \leq \log \left(\frac{1}{2}n^2 + n + 1 \right) + 2 \log H. \quad (5)$$

From this (1) easily follows.

Proof of Theorem 1

For the height H of the polynomial $f(x) = (x^2 - 1)^{n/2}$ with even $n \geq 2$ by Stirling's formula we have $\log H = (1 + o(1))n \log 2/2$. This shows that the constant $2/\log 2$ in (1) is best possible.

To prove (2), observe that assuming $(5/\log 2) \log H < n$ from (5) we obtain

$$n \log 2 < \log \left(\frac{1}{2}n^2 + n + 1 \right) + \frac{2n \log 2}{5},$$

whence $n \leq 9$.

These cases can be checked relatively easily, and (2) holds. In particular, the polynomial

$$(x - 1)^3(x + 1)^2 = x^5 - x^4 - 2x^3 + 2x^2 + x - 1$$

shows that the constant $5/\log 2$ in (2) is best possible.

Theorem 2 (Tijdeman, Varga, H. (202?))

Every polynomial $f(x) \in \mathbb{Z}[x]$ with only rational roots of which no coefficient is divisible by 2 or 3 has degree at most 3.

The example

$$f(x) = x^3 - x^2 - x + 1 = (x - 1)^2(x + 1)$$

shows that degree 3 is possible.

Background of the proof of Theorem 2

The proof is based upon the following two lemmas.

Lemma 1 (Fine (1947))

Let n be a positive integer such that all the coefficients of $(x + 1)^n$ are odd. Then n is of the shape $2^\alpha - 1$ with some $\alpha \in \mathbb{Z}_{\geq 0}$.

Lemma 2 (Tijdeman, Varga, H. (202?))

Let a, b be non-negative integers. Put $n := a + b$. If none of the coefficients of $(x - 1)^a(x + 1)^b$ is divisible by 3, then n is of the shape $3^\beta - 1, 2 \cdot 3^\beta - 1, 3^\gamma + 3^\delta - 1$ or $2 \cdot 3^\gamma + 3^\delta - 1$ with $\beta \geq 0, \gamma > \delta \geq 0$.

Remark. For all the mentioned values in Lemma 2 there are polynomials without coefficients divisible by 3.

Background of the proof of Theorem 2

We may assume that f is monic.

Since the roots of f are odd, Lemma 1 shows that $n + 1$ is a power of 2.

Further, since the roots of f are not divisible by 3, by Lemma 2 we get that $n + 1$ is of the shape 3^β , $2 \cdot 3^\beta$, $3^\gamma + 3^\delta$ or $2 \cdot 3^\gamma + 3^\delta$.

The combination is possible only for $n = 0, 1, 3$.

Polynomials with coeffs having only prime factors coming from a fixed finite set

Theorem 3 (Tijdeman, Varga, H. (202?))

Let S be a finite set of primes with $|S| = s$ and n a positive integer.

There exists an explicitly computable constant $C = C(n, s)$ depending only on n and s and sets T_1, T_2 with $\max(|T_1|, |T_2|) \leq C$ of n -tuples of S -units and $(n-1)/2$ -tuples of S -units for n odd, respectively, such that if $f(x)$ is an S -polynomial of degree n having only rational roots q_1, \dots, q_n , then q_1, \dots, q_n satisfy one of the conditions (i) or (ii):

- (i) $(q_1, \dots, q_n) = u(r_1, \dots, r_n)$ with some $(r_1, \dots, r_n) \in T_1$ and S -unit u ,
- (ii) $n = 2t + 1$ is odd, and re-indexing q_1, \dots, q_n if necessary, we have $q_1 = u$ and $(q_2, \dots, q_n) = v(r_1, -r_1, \dots, r_t, -r_t)$ with some $(r_1, \dots, r_t) \in T_2$ and S -units u, v .

Further, the possibilities (i) and (ii) cannot be excluded.

Background of the proof of Theorem 3

We use the theory of S -unit equations. Let S be a finite set of primes, b_1, \dots, b_m non-zero rationals, and consider the equation

$$b_1x_1 + \dots + b_mx_m = 0 \quad \text{in } S\text{-units } x_1, \dots, x_m. \quad (6)$$

A solution (y_1, \dots, y_m) of (6) is called non-degenerate if

$$\sum_{i \in I} b_i y_i \neq 0 \quad \text{for each non-empty subset } I \text{ of } \{1, \dots, m\}.$$

Two solutions (y_1, \dots, y_m) and (z_1, \dots, z_m) are called proportional, if there is an S -unit u such that $(z_1, \dots, z_m) = u(y_1, \dots, y_m)$.

Lemma 3 (Amoroso and Viada (2009))

Equation (6) has at most $(8m - 8)^{4(m-1)^4(m+s)}$ non-degenerate, non-proportional solutions, where $s = |S|$.

Sketch of the proof of Theorem 3

Write $f(x) = \sum_{j=0}^n a_j x^j$, having only rational roots q_1, \dots, q_n .

By our assumption, a_0, a_1, \dots, a_n are integral S -units.

We have

$$A_j = \sigma_j(q_1, \dots, q_n) \quad (1 \leq j \leq n)$$

where $A_j = (-1)^j a_{n-j}/a_n$ and σ_j is the j -th elementary symmetric polynomial (of degree j) of q_1, \dots, q_n .

This is a system of S -unit equations, which by a careful analysis, with delicate considerations lead to the statement by Lemma 3.

Sketch of the proof of Theorem 3

Finally, we show that the possibilities (i) and (ii) cannot be excluded.

If r_1, \dots, r_n is a set of rational roots of an S -polynomial of degree n , then clearly, the same is true for ur_1, \dots, ur_n for any S -unit u , showing the necessity of (i).

On the other hand, let r_1^2, \dots, r_t^2 be the rational roots of the S -polynomial $(x - r_1^2) \cdots (x - r_t^2)$. Then in the polynomial $(x^2 - r_1^2) \cdots (x^2 - r_t^2)$, all the coefficients of the even powers of x are S -units (while the coefficients of the odd powers of x equal 0). Thus for any S -units u, v , all the coefficients of the polynomial

$$(x + u)(x - vr_1)(x + vr_1) \cdots (x - vr_t)(x + vr_t)$$

are S -units. This shows that (ii) cannot be excluded either.

Problem 1. *Is it true that for any primes p and q there exists an $n_1 = n_1(p, q)$ such that every polynomial $f(x) \in \mathbb{Z}[x]$ with only rational roots of which no coefficient is divisible by p or q has degree at most n_1 ?*

Theorem 2 shows that the answer is 'yes' for the pair of primes $(p, q) = (2, 3)$.

A weaker statement is a restriction to S -polynomials.

Problem 2. *Is it true that for any finite set S of primes there exists an $n_2 = n_2(S)$ such that every S -polynomial $f(x) \in \mathbb{Z}[x]$ with only rational roots has degree at most n_2 ?*

Theorem 2 yields an affirmative answer for sets S of primes with $2, 3 \notin S$.

The last problem is raised by Lemmas 1 and 2.

Problem 3. *Is it true that for every prime p there exists a constant $c(p)$ such that if $f(x) \in \mathbb{Z}[x]$ has only rational roots and none of the coefficients of f is divisible by p , then $\deg(f) + 1$ in its p -adic expansion has at most $c(p)$ non-zero digits? In particular, can one take $c(p) = p - 1$?*

Lemmas 1 and 2 show that the answer is ‘yes’ with $c(p) = p - 1$ for $p = 2, 3$. Note that an affirmative answer to Problem 3 through a deep result of **Stewart** would yield positive answers to Problems 1 and 2, as well.

Thank you very much
for your attention!